

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : A61K 51/04	A2	(11) International Publication Number: WO 00/35492 (43) International Publication Date: 22 June 2000 (22.06.00)
(21) International Application Number: PCT/US99/30315 (22) International Filing Date: 17 December 1999 (17.12.99) (30) Priority Data: 60/112,732 18 December 1998 (18.12.98) US (71) Applicant: DU PONT PHARMACEUTICALS COMPANY [US/US]; Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19807 (US). (72) Inventors: HARRIS, Thomas, David; 56 Zion Hill Road, Salem, NH 03079 (US). RAJODADHYE, Milind; 21 Honeysuckle Road, Westford, MA 01886 (US). (74) Agent: OBRIEN, Maureen, P.; Du Pont Pharmaceuticals Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		(81) Designated States: AL, AU, BR, CA, CN, CZ, EE, HU, IL, IN, JP, KR, LT, LV, MK, MX, NO, NZ, PL, RO, SG, SI, SK, TR, UA, VN, ZA, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: VITRONECTIN RECEPTOR ANTAGONIST PHARMACEUTICALS		
(57) Abstract <p>The present invention describes novel compounds of formula: $(Q)_d-L_n-C_h$, useful for the diagnosis and treatment of cancer, methods of imaging tumors in a patient, and methods of treating cancer in a patient. The present invention also provides novel compounds useful for monitoring therapeutic angiogenesis treatment and destruction of new angiogenic vasculature. The present invention further provides novel compounds useful for imaging atherosclerosis, restenosis, cardiac ischemia, and myocardial reperfusion injury. The present invention still further provides novel compounds useful for the treatment of rheumatoid arthritis. The pharmaceuticals are comprised of a targeting moiety that binds to a receptor that is upregulated during angiogenesis, and optional linking group, and a therapeutically effective radioisotope or diagnostically effective imageable moiety. The imageable moiety is a gamma ray or positron emitting radioisotope, a magnetic resonance imaging contrast agent, and X-ray contrast agent, or an ultrasound contrast agent.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE

VITRONECTIN RECEPTOR ANTAGONIST PHARMACEUTICALS

FIELD OF THE INVENTION

5 The present invention provides novel pharmaceuticals
useful for the diagnosis and treatment of cancer, methods
of imaging tumors in a patient, and methods of treating
cancer in a patient. The pharmaceuticals are comprised
10 of a targeting moiety that binds to the vitronectin
receptor that is expressed in tumor vasculature, an
optional linking group, and a therapeutically effective
radioisotope or diagnostically effective imageable
moiety. The therapeutically effective radioisotope emits
a gamma ray or alpha particle sufficient to be cytotoxic.
15 The imageable moiety is a gamma ray or positron emitting
radioisotope, a magnetic resonance imaging contrast
agent, an X-ray contrast agent, or an ultrasound contrast
agent.

20

BACKGROUND OF THE INVENTION

Cancer is a major public health concern in the
United States and around the world. It is estimated that
over 1 million new cases of invasive cancer will be
25 diagnosed in the United States in 1998. The most
prevalent forms of the disease are solid tumors of the
lung, breast, prostate, colon and rectum. Cancer is
typically diagnosed by a combination of in vitro tests
and imaging procedures. The imaging procedures include
30 X-ray computed tomography, magnetic resonance imaging,
ultrasound imaging and radionuclide scintigraphy.
Frequently, a contrast agent is administered to the
patient to enhance the image obtained by X-ray CT, MRI
and ultrasound, and the administration of a
35 radiopharmaceutical that localizes in tumors is required
for radionuclide scintigraphy.

Treatment of cancer typically involves the use of
external beam radiation therapy and chemotherapy, either
alone or in combination, depending on the type and extent

of the disease. A number of chemotherapeutic agents are available, but generally they all suffer from a lack of specificity for tumors versus normal tissues, resulting in considerable side-effects. The effectiveness of these treatment modalities is also limited, as evidenced by the high mortality rates for a number of cancer types, especially the more prevalent solid tumor diseases. More effective and specific treatment means continue to be needed.

Despite the variety of imaging procedures available for the diagnosis of cancer, there remains a need for improved methods. In particular, methods that can better differentiate between cancer and other pathologic conditions or benign physiologic abnormalities are needed. One means of achieving this desired improvement would be to administer to the patient a metallopharmaceutical that localizes specifically in the tumor by binding to a receptor expressed only in tumors or expressed to a significantly greater extent in tumors than in other tissue. The location of the metallopharmaceutical could then be detected externally either by its imageable emission in the case of certain radiopharmaceuticals or by its effect on the relaxation rate of water in the immediate vicinity in the case of magnetic resonance imaging contrast agents.

This tumor specific metallopharmaceutical approach can also be used for the treatment of cancer when the metallopharmaceutical is comprised of a particle emitting radioisotope. The radioactive decay of the isotope at the site of the tumor results in sufficient ionizing radiation to be toxic to the tumor cells. The specificity of this approach for tumors minimizes the amount of normal tissue that is exposed to the cytotoxic agent and thus may provide more effective treatment with fewer side-effects.

Previous efforts to achieve these desired improvements in cancer imaging and treatment have centered on the use of radionuclide labeled monoclonal antibodies, antibody fragments and other proteins or

polypeptides that bind to tumor cell surface receptors. The specificity of these radiopharmaceuticals is frequently very high, but they suffer from several disadvantages. First, because of their high molecular weight, they are generally cleared from the blood stream very slowly, resulting in a prolonged blood background in the images. Also, due to their molecular weight they do not extravasate readily at the site of the tumor and then only slowly diffuse through the extravascular space to the tumor cell surface. This results in a very limited amount of the radiopharmaceutical reaching the receptors and thus very low signal intensity in imaging and insufficient cytotoxic effect for treatment.

Alternative approaches to cancer imaging and therapy have involved the use of small molecules, such as peptides, that bind to tumor cell surface receptors. An In-111 labeled somatostatin receptor binding peptide, In-111-DTPA-D-Phe¹-octeotide, is in clinical use in many countries for imaging tumors that express the somatostatin receptor (Baker, et al. Life Sci., 1991, 49, 1583-91 and Krenning, et al., Eur. J. Nucl. Med., 1993, 20, 716-31). Higher doses of this radiopharmaceutical have been investigated for potential treatment of these types of cancer (Krenning, et al., Digestion, 1996, 57, 57-61). Several groups are investigating the use of Tc-99m labeled analogs of In-111-DTPA-D-Phe¹-octeotide for imaging and Re-186 labeled analogs for therapy (Flanagan, et al., U.S. 5,556,939, Lyle, et al., U.S. 5,382,654, and Albert et al., U.S. 5,650,134).

Angiogenesis is the process by which new blood vessels are formed from pre-existing capillaries or post capillary venules; it is an important component of a variety of physiological processes including ovulation, embryonic development, wound repair, and collateral vascular generation in the myocardium. It is also central to a number of pathological conditions such as tumor growth and metastasis, diabetic retinopathy, and macular degeneration. The process begins with the

activation of existing vascular endothelial cells in response to a variety of cytokines and growth factors. Tumor released cytokines or angiogenic factors stimulate vascular endothelial cells by interacting with specific
5 cell surface receptors for the factors. The activated endothelial cells secrete enzymes that degrade the basement membrane of the vessels. The endothelial cells then proliferate and invade into the tumor tissue. The endothelial cells differentiate to form lumens, making
10 new vessel offshoots of pre-existing vessels. The new blood vessels then provide nutrients to the tumor permitting further growth and a route for metastasis.

Under normal conditions, endothelial cell proliferation is a very slow process, but it increases
15 for a short period of time during embryogenesis, ovulation and wound healing. This temporary increase in cell turnover is governed by a combination of a number of growth stimulatory factors and growth suppressing factors. In pathological angiogenesis, this normal
20 balance is disrupted resulting in continued increased endothelial cell proliferation. Some of the proangiogenic factors that have been identified include basic fibroblast growth factor (bFGF), angiogenin, TGF-alpha, TGF-beta, and vascular endothelium growth factor
25 (VEGF). While interferon-alpha, interferon-beta and thrombospondin are examples of angiogenesis suppressors.

The proliferation and migration of endothelial cells in the extracellular matrix is mediated by interaction with a variety of cell adhesion molecules (Folkman, J.,
30 Nature Medicine , 1995, 1, 27-31). Integrins are a diverse family of heterodimeric cell surface receptors by which endothelial cells attach to the extracellular matrix, each other and other cells. The integrin $\alpha\beta_1$ is a receptor for a wide variety for a wide variety of
35 extracellular matrix proteins with an exposed tripeptide Arg-Gly-Asp moiety and mediates cellular adhesion to its ligand: vitronectin, fibronectin, and fibrinogen, among others. The integrin $\alpha\beta_1$ is minimally expressed on normal

blood vessels, but is significantly upregulated on vascular cells within a variety of human tumors. The role of the $\alpha_v\beta_3$ receptors is to mediate the interaction of the endothelial cells and the extracellular matrix and facilitate the migration of the cells in the direction of the angiogenic signal, the tumor cell population. Angiogenesis induced by bFGF or TNF-alpha depend on the agency of the integrin $\alpha_v\beta_3$, while angiogenesis induced by VEGF depends on the integrin $\alpha_v\beta_3$ (Cheresh et. al., Science, 1995, 270, 1500-2). Induction of expression of the integrins $\alpha_v\beta_3$ and $\alpha_2\beta_1$ on the endothelial cell surface is another important mechanism by which VEGF promotes angiogenesis (Senger, et. al., Proc. Natl. Acad. Sci USA, 1997, 84, 13612-7).

Angiogenic factors interact with endothelial cell surface receptors such as the receptor tyrosine kinases EGFR, FGFR, PDGFR, Flk-1/KDR, Flt-1, Tek, tie, neuropilin-1, endoglin, endosialin, and Axl. The receptors Flk-1/KDR, neuropilin-1, and Flt-1 recognize VEGF and these interactions play key roles in VEGF-induced angiogenesis. The Tie subfamily of receptor tyrosine kinases are also expressed prominently during blood vessel formation.

Because of the importance of angiogenesis to tumor growth and metastasis, a number of chemotherapeutic approaches are being developed to interfere with or prevent this process. One of these approaches, involves the use of anti-angiogenic proteins such as angiostatin and endostatin. Angiostatin is a 38 kDa fragment of plasminogen that has been shown in animal models to be a potent inhibitor of endothelial cell proliferation. (O'Reilly et. al., Cell, 1994, 79, 315-328) Endostatin is a 20 kDa C-terminal fragment of collagen XVIII that has also been shown to be a potent inhibitor. (O'Reilly et. al., Cell, 1997, 88, 277-285) Systemic therapy with endostatin has been shown to result in strong anti-tumor activity in animal models. However, human clinical trials

of these two chemotherapeutic agents of biological origin have been hampered by lack of availability.

Another approach to anti-angiogenic therapy is to use

- 5 targeting moieties that interact with endothelial cell surface receptors expressed in the angiogenic vasculature to which are attached chemotherapeutic agents. Burrows and Thorpe (Proc. Nat. Acad. Sci, USA, 1993, 90, 8996-9000) described the use of an antibody-immunotoxin
- 10 conjugate to eradicate tumors in a mouse model by destroying the tumor vasculature. The antibody was raised against an endothelial cell class II antigen of the major histocompatibility complex and was then conjugated with the cytotoxic agent, deglycosylated ricin
- 15 A chain. The same group (Clin. Can. Res., 1995, 1, 1623-1634) investigated the use of antibodies raised against the endothelial cell surface receptor, endoglin, conjugated to deglycosylated ricin A chain. Both of these conjugates exhibited potent anti-tumor activity in mouse
- 20 models. However, both still suffer drawbacks to routine human use. As with most antibodies or other large, foreign proteins, there is considerable risk of immunologic toxicity which could limit or preclude administration to humans. Also, while the vasculature
- 25 targeting may improve the local concentration of the attached chemotherapeutic agents, the agents still must be cleaved from the antibody carrier and be transported or diffuse into the cells to be cytotoxic.

- Thus, it is desirable to provide anti-angiogenic
- 30 pharmaceuticals and tumor or new vasculature imaging agents which do not suffer from poor diffusion or transportation, possible immunologic toxicity, limited availability, and/or a lack of specificity.

- Another application of anti-angiogenic therapy is in
- 35 treating rheumatoid arthritis (RA). In RA, the ingrowth of a highly vascularized pannus is caused by the excessive production of angiogenic factors by the infiltrating macrophages, immune cells, or inflammatory cells. Therefore, it is desirable to have new

pharmaceuticals to destroy the highly vascularized pannus that results and thus treat the disease.

There is also a growing interest in therapeutic angiogenesis to improve blood flow in regions of the body that have become ischemic or poorly perfused. Several investigators are using growth factors administered locally to cause new vasculature to form either in the limbs or the heart. The growth factors VEGF and bFGF are the most common for this application. Recent publications include: Takeshita, S., et. al., J. Clin. Invest., 1994, 93, 662-670; and Schaper, W. and Schaper, J., Collateral Circulation: Heart, Brain, Kidney, Limbs, Kluwer Academic Publishers, Boston, 1993. The main applications that are under investigation in a number of laboratories are for improving cardiac blood flow and in improving peripheral vessel blood flow in the limbs. For example, Henry, T. et. al. (J. Amer. College Cardiology, 1998, 31, 65A) describe the use of recombinant human VEGF in patients for improving myocardial perfusion by therapeutic angiogenesis. Patients received infusions of rhVEGF and were monitored by nuclear perfusion imaging 30 and 60 days post treatment to determine improvement in myocardial perfusion. About 50% of patients showed improvement by nuclear perfusion imaging whereas 5/7 showed new collateralization by angiography. Thus, it is desirable to discover a method of monitoring improved cardiac blood flow which is targeted to new collateral vessels themselves and not, as in nuclear perfusion imaging, a regional consequence of new collateral vessels.

The detection, imaging and diagnosis of a number of cardiovascular diseases need to be improved, including restenosis, atherosclerosis, myocardial reperfusion injury, and myocardial ischemia, stunning or infarction. It has recently been determined that in all of these disease conditions, the integrin receptor $\alpha v \beta 3$ plays an important role.

For example, in the restenosis complication that occurs in ~30-50% of patients having undergone angioplasty or stent placement, neointimal hyperplasia and ultimate reocclusion is caused by aggressively
5 proliferating vascular smooth muscle cells that express $\alpha v\beta 3$. (Cardiovascular Res., 1997, 36, 408-428; DDT, 1997, 2, 187-199; Current Pharm. Design, 1997, 3, 545-584)

Atherosclerosis proceeds from an initial endothelial damage that results in the recruitment and subintimal
10 migration of monocytes at the site of the injury. Growth factors are released which induce medial smooth muscle cells to proliferate and migrate to the intimal layer. The migrating smooth muscle cells express $\alpha v\beta 3$.

In reperfusion injury, neutrophil transmigration is
15 integrin dependent and the integrins moderate initial infiltration into the viable border zone. The induction of $\alpha 5\beta 1$, $\alpha 4\beta 1$ and $\alpha v\beta 5$ in infiltrating neutrophils occurs within 3 to 5 hours after reperfusion as neutrophils move from the border zone to the area of necrosis.
20 (Circulation, 1999, 100, I-275)

Acute or chronic occlusion of a coronary artery is known to result in angiogenesis in the heart as native collateral vessels are recruited to attempt to relieve the ischemia. However, even a gradual occlusion usually
25 results in areas of infarction as the resulting angiogenesis is not sufficient to prevent damage. Cardiac angiogenesis has been associated with increased expression of the growth factors VEGF and FGF and the upregulation of the growth factor receptors flt-1 and
30 flk-1/KDR. (Drugs, 1999, 58, 391-396)

SUMMARY OF THE INVENTION

It is one object of the present invention to provide improved anti-angiogenic pharmaceuticals, comprised of a targeting moiety that binds to the vitronectin receptor
5 that is expressed in tumor neovasculature, an optional linking group, and a radioisotope. The vitronectin receptor binding compounds target the radioisotope to the tumor neovasculature. The beta or alpha-particle emitting radioisotope emits a cytotoxic amount of
10 ionizing radiation which results in cell death. The penetrating ability of radiation obviates the requirement that the cytotoxic agent diffuse or be transported into the cell to be cytotoxic.

It is another object of the present invention to
15 provide pharmaceuticals to treat rheumatoid arthritis. These pharmaceuticals comprise a targeting moiety that binds to a receptor that is upregulated during angiogenesis, an optional linking group, and a radioisotope that emits cytotoxic radiation (i.e., beta
20 particles, alpha particles and Auger or Coster-Kronig electrons). In rheumatoid arthritis, the ingrowth of a highly vascularized pannus is caused by the excessive production of angiogenic factors by the infiltrating macrophages, immune cells, or inflammatory cells.
25 Therefore, the radiopharmaceuticals of the present invention that emit cytotoxic radiation could be used to destroy the new angiogenic vasculature that results and thus treat the disease.

It is another object of the present invention to
30 provide imaging agents, comprised of vitronectin receptor binding compounds conjugated to an imageable moiety, such as a gamma ray or positron emitting radioisotope, a magnetic resonance imaging contrast agent, an X-ray contrast agent, or an ultrasound contrast agent. These
35 imaging agents are useful for imaging tumor neovasculature, therapeutic angiogenesis interventions in the heart, natural angiogenic processes in response to acute or chronic coronary vessel occlusion, restenosis

post-angioplasty, atherosclerosis and plaque formation, and reperfusion injury.

It is another object of the present invention to provide compounds useful for preparing the
5 pharmaceuticals of the present invention. These compounds are comprised of a non-peptide **quinolone** containing targeting moiety that binds to a receptor that is upregulated during angiogenesis or during
10 cardiovascular diseases, Q, an optional linking group, L_n , and a metal chelator or bonding moiety, C_h . The compounds may have one or more protecting groups attached to the metal chelator or bonding moiety. The protecting groups provide improved stability to the reagents for long-term storage and are removed either immediately
15 prior to or concurrent with the synthesis of the radiopharmaceuticals. Alternatively, the compounds of the present invention are comprised of a peptide or peptidomimetic targeting moiety that binds to a receptor that is upregulated during angiogenesis or during
20 cardiovascular diseases, Q, an optional linking group, L_n , and a surfactant, S_f .

The pharmaceuticals of the present invention may be used for diagnostic and/or therapeutic purposes. Diagnostic radiopharmaceuticals of the present invention
25 are pharmaceuticals comprised of a diagnostically useful radionuclide (i.e., a radioactive metal ion that has imageable gamma ray or positron emissions). Therapeutic radiopharmaceuticals of the present invention are pharmaceuticals comprised of a therapeutically useful
30 radionuclide, a radioactive metal ion that emits ionizing radiation such as beta particles, alpha particles and Auger or Coster-Kronig electrons.

The pharmaceuticals comprising a gamma ray or positron emitting radioactive metal ion are useful for
35 imaging tumors and by gamma scintigraphy or positron emission tomography. The pharmaceuticals comprising a gamma ray or positron emitting radioactive metal ion are also useful for imaging therapeutic angiogenesis, natural

angiogenic processes in response to acute or chronic coronary vessel occlusion, restenosis post-angioplasty, atherosclerosis and plaque formation, and reperfusion injury by gamma scintigraphy or positron emission tomography. The pharmaceuticals comprising a particle emitting radioactive metal ion are useful for treating cancer by delivering a cytotoxic dose of radiation to the tumors. The pharmaceuticals comprising a particle emitting radioactive metal ion are also useful for treating rheumatoid arthritis by destroying the formation of angiogenic vasculature. The pharmaceuticals comprising a paramagnetic metal ion are useful as magnetic resonance imaging contrast agents. The pharmaceuticals comprising one or more X-ray absorbing or "heavy" atoms of atomic number 20 or greater are useful as X-ray contrast agents. The pharmaceuticals comprising a microbubble of a biocompatible gas, a liquid carrier, and a surfactant microsphere, are useful as ultrasound contrast agents.


20

DETAILED DESCRIPTION OF THE INVENTION

[1] Thus, in a first embodiment, the present invention provides a novel compound, comprising: a targeting moiety and a chelator, wherein the targeting moiety is bound to the chelator, is a quinolone nonpeptide, and binds to a receptor that is upregulated during angiogenesis and the compound has 0-1 linking groups between the targeting moiety and chelator.


30

[2] In a preferred embodiment, the receptor is the integrin $\alpha_v\beta_3$ or $\alpha_v\beta_5$ and the compound is of the formula:



wherein, Q is a compound of Formula (II):

35




(II)

- 5 including stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, or pharmaceutically acceptable salt or prodrug forms thereof wherein:

R^{1e} is selected from:

10



15

A^e is $-CH_2-$ or $-N(R^{10e})-$;

A^{1e} and B^e are independently $-CH_2-$ or $-N(R^{10e})-$;

5

D^e is $-N(R^{10e})-$ or $-S-$;

E^e-F^e is $-C(R^{2e})=C(R^{3e})-$ or $-C(R^{2e})_2C(R^{3e})_2-$;

10 J^e is $-C(R^{2e})-$ or $-N-$;

K^e , L^e and M^e are independently $-C(R^{2e})-$ or $-C(R^{3e})-$;

R^{2e} and R^{3e} are independently selected from:

15 H, C_1-C_4 alkoxy, $NR^{11e}R^{12e}$, halogen, NO_2 , CN, CF_3 ,
 C_1-C_6 alkyl, C_3-C_6 alkenyl, C_3-C_7 cycloalkyl,
 C_3-C_7 cycloalkyl(C_1-C_4 alkyl), aryl(C_1-C_6 alkyl)-,
 (C_1-C_6 alkyl)carbonyl, (C_1-C_6 alkoxy)carbonyl,
 arylcarbonyl, and aryl substituted with 0-4 R^{7e} ,

20

alternatively, when R^{2e} and R^{3e} are substituents on adjacent atoms, they can be taken together with the carbon atoms to which they are attached to form a 5-7 membered carbocyclic or 5-7 membered heterocyclic aromatic or nonaromatic ring system, said carbocyclic or heterocyclic ring being substituted with 0-2 groups selected from C_1-C_4 alkyl, C_1-C_4 alkoxy, halo, cyano, amino, CF_3 and NO_2 ;

25

30 R^{2ae} is selected from:

H, C_1-C_{10} alkyl, C_2-C_6 alkenyl, C_3-C_{11} cycloalkyl,
 C_3-C_7 cycloalkyl(C_1-C_4 alkyl), aryl, aryl(C_1-C_4 alkyl)-,
 (C_2-C_7 alkyl)carbonyl, arylcarbonyl,
 (C_2-C_{10} alkoxy)carbonyl, C_3-C_7 cycloalkoxycarbonyl,
 35 C_7-C_{11} bicycloalkoxycarbonyl, aryloxy carbonyl,
 aryl(C_1-C_{10} alkoxy)carbonyl,
 C_1-C_6 alkylcarbonyloxy(C_1-C_4 alkoxy)carbonyl,
 arylcarbonyloxy(C_1-C_4 alkoxy)carbonyl, and

C₃-C₇ cycloalkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl;

R^{7e} is selected from:

- 5 H, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, aryl, aryl(C₁-C₄ alkyl)-, (C₁-C₄ alkyl)carbonyl, CO₂R^{18ae}, SO₂R^{11e}, SO₂NR^{10e}R^{11e}, OR^{10e}, and N(R^{11e})R^{12e};

U^e is selected from:

- 10 - (CH₂)_n^e-, - (CH₂)_n^eO(CH₂)_m^e-, - (CH₂)_n^eN(R¹²)(CH₂)_m^e-,
-NH(CH₂)_n^e-, - (CH₂)_n^eC(=O)(CH₂)_m^e-,
- (CH₂)_n^eS(O)_p^e(CH₂)_m^e-, - (CH₂)_n^eNHNH(CH₂)_m^e-,
-N(R^{10e})C(=O)-, -NHC(=O)(CH₂)_n^e-, -C(=O)N(R^{10e})-, and
-N(R^{10e})S(O)_p^e-;

- 15 G^e is N or CR^{19e};

W^e is -C(=O)-N(R^{10e})-(C₁-C₃ alkylene)-, in which the alkylene group is substituted by R^{8e} and by R^{9e}:

- 20 R^{8e} and R^{9e} are independently selected from:
H, CO₂R^{18be}, C(=O)R^{18be}, CONR¹⁷R^{18be},
C₁-C₁₀ alkyl substituted with 0-1 R^{6e},
C₂-C₁₀ alkenyl substituted with 0-1 R^{6e},
C₂-C₁₀ alkynyl substituted with 0-1 R^{6e},
25 C₃-C₈ cycloalkyl substituted with 0-1 R^{6e},
C₅-C₆ cycloalkenyl substituted with 0-1 R^{6e},
(C₁-C₁₀ alkyl)carbonyl,
C₃-C₁₀ cycloalkyl(C₁-C₄ alkyl)-,
phenyl substituted with 0-3 R^{6e},
30 naphthyl substituted with 0-3 R^{6e},
a 5-10 membered heterocyclic ring containing 1-3 N,
O, or S heteroatoms, wherein said heterocyclic ring may be saturated, partially saturated, or fully unsaturated, said heterocyclic ring being substituted with 0-2 R^{7e},
35 C₁-C₁₀ alkoxy substituted with 0-2 R^{7e},
hydroxy, nitro, -N(R^{10e})R^{11e}, -N(R^{16e})R^{17e}, aryl(C₀-C₆ alkyl)carbonyl, aryl(C₃-C₆ alkyl),

heteroaryl(C₁-C₆ alkyl), CONR^{18ae}R^{20e}, SO₂R^{18ae},
and SO₂NR^{18ae}R^{20e},

providing that any of the above alkyl, cycloalkyl,
aryl or heteroaryl groups may be unsubstituted
5 or substituted independently with 1-2 R^{7e};

R^{6e} is selected from:

H, C₁-C₁₀ alkyl, hydroxy, C₁-C₁₀ alkoxy, nitro, C₁-C₁₀
alkylcarbonyl, -N(R^{11e})R^{12e}, cyano, halo, CF₃,
10 CHO, CO₂R^{18be}, C(=O)R^{18be}, CONR^{17e}R^{18be},
OC(=O)R^{10e}, OR^{10e}, OC(=O)NR^{10e}R^{11e},
NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},
NR^{10e}C(=O)NR^{10e}R^{11e}, NR^{10e}SO₂NR^{10e}R^{11e},
NR^{10e}SO₂R^{21e}, S(O)_pR^{11e}, SO₂NR^{10e}R^{11e},
15 aryl substituted with 0-3 groups selected from
halogen, C₁-C₆ alkoxy, C₁-C₆ alkyl, CF₃,
S(O)_mMe, and -NMe₂,
aryl(C₁-C₄ alkyl)-, said aryl being substituted with
0-3 groups selected from halogen, C₁-C₆ alkoxy,
20 C₁-C₆ alkyl, CF₃, S(O)_pMe, and -NMe₂, and
a 5-10 membered heterocyclic ring containing 1-3 N, O, or
S heteroatoms, wherein said heterocyclic ring may be
saturated, partially saturated, or fully unsaturated,
said heterocyclic ring being substituted with 0-2 R^{7e};

25

R^{10e} is selected from:

H, CF₃, C₃-C₆ alkenyl, C₃-C₁₁ cycloalkyl, aryl,
(C₃-C₁₁ cycloalkyl)methyl, aryl(C₁-C₄ alkyl), and C₁-
C₁₀ alkyl substituted with 0-2 R^{6e};

30

R^{11e} is selected from:

H, hydroxy, C₁-C₈ alkyl, C₃-C₆ alkenyl, C₃-C₁₁
cycloalkyl, (C₃-C₁₁ cycloalkyl)methyl, C₁-C₆ alkoxy,
benzyloxy, aryl, heteroaryl, heteroaryl(C₁-C₄ alkyl)-
35 , aryl(C₁-C₄ alkyl), adamantylmethyl, and
C₁-C₁₀ alkyl substituted with 0-2 R^{4e};

R^{4e} is selected from:

- H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-, (C₁-C₁₀ alkyl)carbonyl, aryl, heteroaryl, aryl(C₁-C₆ alkyl)-, and heteroaryl(C₁-C₆ alkyl)-, wherein said aryl or
- 5 heteroaryl groups are substituted with 0-2 substituents independently selected from the group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, F, Cl, Br, CF₃, and NO₂,
- 10 alternatively, when R^{10e} and R^{11e} are both substituents on the same nitrogen atom (as in -NR^{10e}R^{11e}) they may be taken together with the nitrogen atom to which they are attached to form a heterocycle selected from:
- 15 3-azabicyclononyl, 1,2,3,4-tetrahydro-1-quinolinyl, 1,2,3,4-tetrahydro-2-isoquinolinyl, 1-piperidinyl, 1-morpholinyl, 1-pyrrolidinyl, thiamorpholinyl, thiazolidinyl, and 1-piperazinyl;
- said heterocycle being substituted with 0-3 groups selected from: C₁-C₆ alkyl, aryl, heteroaryl,
- 20 aryl(C₁-C₄ alkyl)-, (C₁-C₆ alkyl)carbonyl, (C₃-C₇ cycloalkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl, aryl(C₁-C₄ alkoxy)carbonyl, C₁-C₆ alkylsulfonyl, and arylsulfonyl;
- 25 R^{12e} is selected from:
- H, C₁-C₆ alkyl, triphenylmethyl, methoxymethyl, methoxyphenyldiphenylmethyl, trimethylsilylethoxymethyl, (C₁-C₆ alkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl, (C₁-C₆ alkyl)aminocarbonyl,
- 30 C₃-C₆ alkenyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-, aryl, heteroaryl(C₁-C₆ alkyl)carbonyl, heteroarylcabonyl, aryl(C₁-C₆ alkyl)-, (C₁-C₆ alkyl)carbonyl, arylcarbonyl, C₁-C₆ alkylsulfonyl, arylsulfonyl, aryl(C₁-C₆
- 35 alkyl)sulfonyl, heteroarylsulfonyl, heteroaryl(C₁-C₆ alkyl)sulfonyl, aryloxy carbonyl, and aryl(C₁-C₆ alkoxy)carbonyl, wherein said aryl groups are substituted with 0-2 substituents selected from the

group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, halo, CF₃, and nitro;

R^{16e} is selected from:

- 5 -C(=O)OR^{18ae}, -C(=O)R^{18be}, -C(=O)N(R^{18be})₂,
 -C(=O)NHSO₂R^{18ae}, -C(=O)NHC(=O)R^{18be},
 -C(=O)NHC(=O)OR^{18ae}, -C(=O)NHSO₂NHR^{18be}, -SO₂R^{18ae},
 -SO₂N(R^{18be})₂, and -SO₂NHC(=O)OR^{18be};

10 R^{17e} is selected from:

H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇
 cycloalkyl(C₁-C₄ alkyl)-, aryl, aryl(C₁-C₆ alkyl)-,
 and heteroaryl(C₁-C₆ alkyl);

15 R^{18ae} is selected from:

- C₁-C₈ alkyl optionally substituted with a bond to L_n,
 C₃-C₁₁ cycloalkyl optionally substituted with a
 bond to L_n, aryl(C₁-C₆ alkyl)- optionally
 substituted with a bond to L_n, heteroaryl(C₁-C₆
 20 alkyl)- optionally substituted with a bond to
 L_n, (C₁-C₆ alkyl)heteroaryl optionally
 substituted with a bond to L_n, biaryl(C₁-C₆
 alkyl) optionally substituted with a bond to
 L_n, heteroaryl optionally substituted with a
 25 bond to L_n, phenyl substituted with 3-4 R^{19e} and
 optionally substituted with a bond to L_n,
 naphthyl substituted with 0-4 R^{19e} and
 optionally substituted with a bond to L_n, and a
 bond to L_n, wherein said aryl or heteroaryl
 30 groups are optionally substituted with 0-4 R^{19e};

R^{18be} is H or R^{18ae};

R^{19e} is selected from:

- 35 H, halogen, CF₃, CO₂H, CN, NO₂, -NR^{11e}R^{12e}, OCF₃,
 C₁-C₈ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl,
 C₃-C₁₁ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-,
 aryl(C₁-C₆ alkyl)-, C₁-C₆ alkoxy, C₁-C₄

alkoxycarbonyl, aryl, aryl-O-, aryl-SO₂-,
heteroaryl, and heteroaryl-SO₂-, wherein said aryl
and heteroaryl groups are substituted with 0-4
groups selected from hydrogen, halogen, CF₃, C₁-C₃
5 alkyl, and C₁-C₃ alkoxy;

R^{20e} is selected from:
hydroxy, C₁-C₁₀ alkyloxy, C₃-C₁₁ cycloalkyloxy,
aryloxy, aryl(C₁-C₄ alkyl)oxy,
10 C₂-C₁₀ alkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
C₂-C₁₀ alkoxycarbonyloxy(C₁-C₂ alkyl)oxy-,
C₂-C₁₀ alkoxycarbonyl(C₁-C₂ alkyl)oxy-,
C₃-C₁₀ cycloalkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
C₃-C₁₀ cycloalkoxycarbonyloxy(C₁-C₂ alkyl)oxy-,
15 C₃-C₁₀ cycloalkoxycarbonyl(C₁-C₂ alkyl)oxy-,
aryloxycarbonyl(C₁-C₂ alkyl)oxy-,
aryloxycarbonyloxy(C₁-C₂ alkyl)oxy-,
arylcarbonyloxy(C₁-C₂ alkyl)oxy-,
C₁-C₅ alkoxy(C₁-C₅ alkyl)carbonyloxy(C₁-C₂ alkyl)oxy,
20 (5-(C₁-C₅ alkyl)-1,3-dioxa-cyclopenten-2-one-
yl)methyloxy,
(5-aryl-1,3-dioxa-cyclopenten-2-one-yl)methyloxy,
and
(R^{10e})(R^{11e})N-(C₁-C₁₀ alkoxy)-;


25 R^{21e} is selected from:
C₁-C₈ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl, (C₃-C₁₁
cycloalkyl)methyl, aryl, aryl(C₁-C₄ alkyl)-, and C₁-
C₁₀ alkyl substituted with 0-2 R^{7e};

30 R^{22e} is selected from:
-C(=O)-R^{18be}, -C(=O)N(R^{18be})₂, -C(=O)NHSO₂R^{18ae}, -
C(=O)NHC(=O)R^{18be}, -C(=O)NHC(=O)OR^{18ae}, and -
C(=O)NHSO₂NHR^{18be};

35 Y^e is selected from:

-COR^{20e}, -SO₃H, -PO₃H, -CONHNHSO₂CF₃, -CONHSO₂R^{18ae},
 -CONHSO₂NHR^{18be}, -NHCOCF₃, -NHCONHSO₂R^{18ae},
 -NHSO₂R^{18ae}, -OPO₃H₂, -OSO₃H, -PO₃H₂, -SO₂NHCOR^{18ae},
 -SO₂NHCO₂R^{18ae},

5



m^e is 0-2;

10 n^e is 0-4;

p^e is 0-2;

r^e is 0-2;

15

with the following proviso: n^e and m^e are chosen such
 that the number of atoms connecting R^{1e} and y^e is in
 the range of 8-14;

20 d is selected from 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

d' is 1-100;

L_n is a linking group having the formula:

25

((W)_h-(CR^{6R7})_g)_x-(Z)_k-((CR^{6aR7a})_{g'}-(W)_{h'})_{x'};

W is independently selected at each occurrence from the

30 group: O, S, NH, NHC(=O), C(=O)NH, NR⁸C(=O), C(=O)N
 R⁸, C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂,
 SO₂NH, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'}, (OCH₂CH₂CH₂)_{s''},
 (CH₂CH₂CH₂O)_t, and (aa)_t;

aa is independently at each occurrence an amino acid;

35

Z is selected from the group: aryl substituted with 0-3 R^{10} , C_{3-10} cycloalkyl substituted with 0-3 R^{10} , and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R^{10} ;

R^6 , R^{6a} , R^7 , R^{7a} , and R^8 are independently selected at each occurrence from the group: H, =O, COOH, SO_3H , PO_3H , C_1-C_5 alkyl substituted with 0-3 R^{10} , aryl substituted with 0-3 R^{10} , benzyl substituted with 0-3 R^{10} , and C_1-C_5 alkoxy substituted with 0-3 R^{10} , $NHC(=O)R^{11}$, $C(=O)NHR^{11}$, $NHC(=O)NHR^{11}$, NHR^{11} , R^{11} , and a bond to Ch ;

R^{10} is independently selected at each occurrence from the group: a bond to Ch , $COOR^{11}$, $C(=O)NHR^{11}$, $NHC(=O)R^{11}$, OH, NHR^{11} , SO_3H , PO_3H , $-OPO_3H_2$, $-OSO_3H$, aryl substituted with 0-3 R^{11} , C_1-5 alkyl substituted with 0-1 R^{12} , C_1-5 alkoxy substituted with 0-1 R^{12} , and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R^{11} ;

R^{11} is independently selected at each occurrence from the group: H, alkyl substituted with 0-1 R^{12} , aryl substituted with 0-1 R^{12} , a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-1 R^{12} , C_{3-10} cycloalkyl substituted with 0-1 R^{12} , polyalkylene glycol substituted with 0-1 R^{12} , carbohydrate substituted with 0-1 R^{12} , cyclodextrin substituted with 0-1 R^{12} , amino acid substituted with 0-1 R^{12} , polycarboxyalkyl substituted with 0-1 R^{12} , polyazaalkyl substituted with 0-1 R^{12} , peptide substituted with 0-1 R^{12} , wherein the peptide is comprised of 2-10 amino acids, 3,6-O-disulfo-B-D-galactopyranosyl, bis(phosphonomethyl)glycine, and a bond to Ch ;

R^{12} is a bond to C_h ;

k is selected from 0, 1, and 2;

5 h is selected from 0, 1, and 2;

h' is selected from 0, 1, and 2;

g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

s is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

10 s' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

s'' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;


t is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

x is selected from 0, 1, 2, 3, 4, and 5;

15 x' is selected from 0, 1, 2, 3, 4, and 5;

C_h is a metal bonding unit having a formula selected from the group:



25 $A^1, A^2, A^3, A^4, A^5, A^6, A^7,$ and A^8 are independently selected at each occurrence from the group: NR^{13} , $NR^{13}R^{14}$, S , SH , $S(Pg)$, O , OH , PR^{13} , $PR^{13}R^{14}$, $P(O)R^{15}R^{16}$, and a bond to L_n ;

E is a bond, CH, or a spacer group independently selected at each occurrence from the group: C₁-C₁₀ alkyl substituted with 0-3 R¹⁷, aryl substituted with 0-3 R¹⁷, C₃-₁₀ cycloalkyl substituted with 0-3 R¹⁷,
5 heterocyclo-C₁-₁₀ alkyl substituted with 0-3 R¹⁷, wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆-₁₀ aryl-C₁-₁₀ alkyl substituted with 0-3 R¹⁷, C₁-₁₀ alkyl-C₆-₁₀ aryl- substituted with 0-3 R¹⁷, and a
10 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁷;

15 R¹³ and R¹⁴ are each independently selected from the group: a bond to L_n, hydrogen, C₁-C₁₀ alkyl substituted with 0-3 R¹⁷, aryl substituted with 0-3 R¹⁷, C₁-₁₀ cycloalkyl substituted with 0-3 R¹⁷, heterocyclo-C₁-₁₀ alkyl substituted with 0-3 R¹⁷,
20 wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆-₁₀ aryl-C₁-₁₀ alkyl substituted with 0-3 R¹⁷, C₁-₁₀ alkyl-C₆-₁₀ aryl- substituted with 0-3 R¹⁷, a 5-10
25 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁷, and an electron, provided that when one of R¹³ or R¹⁴ is an electron, then the other is also an electron;


30 alternatively, R¹³ and R¹⁴ combine to form =C(R²⁰)(R²¹);

R¹⁵ and R¹⁶ are each independently selected from the group: a bond to L_n, -OH, C₁-C₁₀ alkyl substituted
35 with 0-3 R¹⁷, C₁-C₁₀ alkyl substituted with 0-3 R¹⁷, aryl substituted with 0-3 R¹⁷, C₃-₁₀ cycloalkyl substituted with 0-3 R¹⁷, heterocyclo-C₁-₁₀ alkyl substituted with 0-3 R¹⁷, wherein the heterocyclo

- group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆-10 aryl-C₁-10 alkyl substituted with 0-3 R¹⁷, C₁-10 alkyl-C₆-10 aryl- substituted with 0-3 R¹⁷, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁷;
- 10 R¹⁷ is independently selected at each occurrence from the group: a bond to L_n, =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R¹⁸, -C(=O)R¹⁸, -C(=O)N(R¹⁸)₂, -CHO, -CH₂OR¹⁸, -OC(=O)R¹⁸, -OC(=O)OR^{18a}, -OR¹⁸, -OC(=O)N(R¹⁸)₂, -NR¹⁹C(=O)R¹⁸, -NR¹⁹C(=O)OR^{18a}, -NR¹⁹C(=O)N(R¹⁸)₂,
 15 -NR¹⁹SO₂N(R¹⁸)₂, -NR¹⁹SO₂R^{18a}, -SO₃H, -SO₂R^{18a}, -SR¹⁸, -S(=O)R^{18a}, -SO₂N(R¹⁸)₂, -N(R¹⁸)₂, -NHC(=S)NHR¹⁸, =NOR¹⁸, NO₂, -C(=O)NHOR¹⁸, -C(=O)NHN(R¹⁸)R^{18a}, -OCH₂CO₂H, 2-(1-morpholino)ethoxy, C₁-C₅ alkyl, C₂-C₄ alkenyl, C₃-C₆ cycloalkyl, C₃-C₆ cycloalkylmethyl, C₂-C₆ alkoxyalkyl, aryl
 20 substituted with 0-2 R¹⁸, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O;
- 25 R¹⁸, R^{18a}, and R¹⁹ are independently selected at each occurrence from the group: a bond to L_n, H, C₁-C₆ alkyl, phenyl, benzyl, C₁-C₆ alkoxy, halide, nitro, cyano, and trifluoromethyl;
- 30 Pg is a thiol protecting group;
- R²⁰ and R²¹ are independently selected from the group: H, C₁-C₁₀ alkyl, -CN, -CO₂R²⁵, -C(=O)R²⁵, -C(=O)N(R²⁵)₂, C₂-C₁₀ 1-alkene substituted with 0-3 R²³, C₂-C₁₀
 35 1-alkyne substituted with 0-3 R²³, aryl substituted with 0-3 R²³, unsaturated 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3

R^{23} , and unsaturated C₃₋₁₀ carbocycle substituted with 0-3 R^{23} ;

alternatively, R^{20} and R^{21} , taken together with the
 5 divalent carbon radical to which they are attached form:



10 R^{22} and R^{23} are independently selected from the group: H, R^{24} , C₁₋₁₀ alkyl substituted with 0-3 R^{24} , C₂₋₁₀ alkenyl substituted with 0-3 R^{24} , C₂₋₁₀ alkynyl substituted with 0-3 R^{24} , aryl substituted with 0-3 R^{24} , a 5-10 membered heterocyclic ring system
 15 containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R^{24} , and C₃₋₁₀ carbocycle substituted with 0-3 R^{24} ;

alternatively, R^{22} , R^{23} taken together form a fused
 20 aromatic or a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O;

a and b indicate the positions of optional double bonds
 25 and n is 0 or 1;


R^{24} is independently selected at each occurrence from the group: =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R²⁵,
 -C(=O)R²⁵, -C(=O)N(R²⁵)₂, -N(R²⁵)₃⁺, -CH₂OR²⁵,
 30 -OC(=O)R²⁵, -OC(=O)OR^{25a}, -OR²⁵, -OC(=O)N(R²⁵)₂,
 -NR²⁶C(=O)R²⁵, -NR²⁶C(=O)OR^{25a}, -NR²⁶C(=O)N(R²⁵)₂,
 -NR²⁶SO₂N(R²⁵)₂, -NR²⁶SO₂R^{25a}, -SO₃H, -SO₂R^{25a}, -SR²⁵,
 -S(=O)R^{25a}, -SO₂N(R²⁵)₂, -N(R²⁵)₂, =NOR²⁵,

-C(=O)NHR²⁵, -OCH₂CO₂H, and 2-(1-morpholino)ethoxy;
and,

5 R²⁵, R^{25a}, and R²⁶ are each independently selected at each
occurrence from the group: hydrogen and C₁-C₆
alkyl;

and a pharmaceutically acceptable salt thereof.

10 [3] In a more preferred embodiment, the present
invention provides a compound wherein: Q is a compound of
Formula (IV):




5

(IV)

including stereoisomeric forms thereof, or mixtures of
stereoisomeric forms thereof, or pharmaceutically
acceptable salt or prodrug forms thereof wherein:

10

R^{1e} is selected from:



15

20

R^{2e} and R^{3e} are independently selected from:

H, C₁-C₄ alkoxy, NR^{11e}R^{12e}, halogen, NO₂, CN, CF₃,
C₁-C₆ alkyl, C₃-C₆ alkenyl, C₃-C₇ cycloalkyl,
C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl(C₁-C₆ alkyl)-,

25

(C₁-C₆ alkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl, arylcarbonyl, and aryl substituted with 0-4 R^{7e},

5 alternatively, when R^{2e} and R^{3e} are substituents on adjacent atoms, they can be taken together with the carbon atoms to which they are attached to form a 5-7 membered carbocyclic or 5-7 membered heterocyclic aromatic or nonaromatic ring system, said carbocyclic or heterocyclic ring being substituted
10 with 0-2 groups selected from C₁-C₄ alkyl, C₁-C₄ alkoxy, halo, cyano, amino, CF₃ and NO₂;

R^{2ae} is selected from:

15 H, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl, aryl(C₁-C₄ alkyl)-, (C₂-C₇ alkyl)carbonyl, arylcarbonyl, (C₂-C₁₀ alkoxy)carbonyl, C₃-C₇ cycloalkoxycarbonyl, C₇-C₁₁ bicycloalkoxycarbonyl, aryloxy carbonyl, aryl(C₁-C₁₀ alkoxy)carbonyl,
20 C₁-C₆ alkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl, arylcarbonyloxy(C₁-C₄ alkoxy)carbonyl, and C₃-C₇ cycloalkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl;

R^{7e} is selected from:

25 H, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, aryl, aryl(C₁-C₄ alkyl)-, (C₁-C₄ alkyl)carbonyl, CO₂R^{18ae}, SO₂R^{11e}, SO₂NR^{10e}R^{11e}, OR^{10e}, and N(R^{11e})R^{12e};

U^e is selected from:

30 -(CH₂)_n^e-, -(CH₂)_n^eO(CH₂)_m^e-, -NH(CH₂)_n^e-,
-N(R^{10e})C(=O)-, -NHC(=O)(CH₂)_n^e-, and -C(=O)N(R^{10e})-;

G^e is N or CR^{19e};

35 R^{8e} is selected from:

H, CO₂R^{18be}, C(=O)R^{18be}, CONR^{17e}R^{18be},
C₁-C₁₀ alkyl substituted with 0-1 R^{6e},
C₂-C₁₀ alkenyl substituted with 0-1 R^{6e},

- C₂-C₁₀ alkynyl substituted with 0-1 R^{6e},
 C₃-C₈ cycloalkyl substituted with 0-1 R^{6e},
 C₅-C₆ cycloalkenyl substituted with 0-1 R^{6e},
 (C₁-C₁₀ alkyl)carbonyl,
 5 C₃-C₁₀ cycloalkyl(C₁-C₄ alkyl)-,
 phenyl substituted with 0-3 R^{6e},
 naphthyl substituted with 0-3 R^{6e},
 a 5-10 membered heterocyclic ring containing 1-3 N,
 O, or S heteroatoms, wherein said heterocyclic
 10 ring may be saturated, partially saturated, or
 fully unsaturated, said heterocyclic ring being
 substituted with 0-2 R^{7e};

R^{9e} is selected from:

- 15 C₁-C₁₀ alkyl substituted with 0-1 R^{6e},
 C₁-C₁₀ alkoxy substituted with 0-2 R^{7e},
 H, nitro, N(R^{11e})R^{12e}, OC(=O)R^{10e}, OR^{10e},
 OC(=O)NR^{10e}R^{11e}, NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},
 NR^{10e}C(=O)NR^{10e}R^{11e}, NR^{10e}SO₂NR^{10e}R^{11e},
 20 NR^{10e}SO₂R^{21e}, hydroxy, OR^{22e}, -N(R^{10e})R^{11e}, -
 N(R^{16e})R^{17e}, aryl(C₀-C₆ alkyl)carbonyl, aryl(C₁-
 C₆ alkyl), heteroaryl(C₁-C₆ alkyl), CONR^{18ae}R^{20e},
 SO₂R^{18ae}, and SO₂NR^{18ae}R^{20e},
 providing that any of the above alkyl, cycloalkyl,
 25 aryl or heteroaryl groups may be unsubstituted
 or substituted independently with 1-2 R^{7e};

R^{6e} is selected from:

- 30 H, C₁-C₁₀ alkyl, hydroxy, C₁-C₁₀ alkoxy, nitro, C₁-C₁₀
 alkylcarbonyl, -N(R^{11e})R^{12e}, cyano, halo, CF₃,
 CHO, CO₂R^{18be}, C(=O)R^{18be}, CONR^{17e}R^{18be},
 OC(=O)R^{10e}, OR^{10e}, OC(=O)NR^{10e}R^{11e},
 NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},
 NR^{10e}C(=O)NR^{10e}R^{11e}, NR^{10e}SO₂NR^{10e}R^{11e},
 35 NR^{10e}SO₂R^{21e}, S(O)_pR^{11e}, SO₂NR^{10e}R^{11e},
 aryl substituted with 0-3 groups selected from
 halogen, C₁-C₆ alkoxy, C₁-C₆ alkyl, CF₃,
 S(O)_mMe, and -NMe₂,

aryl(C₁-C₄ alkyl)-, said aryl being substituted with
 0-3 groups selected from halogen, C₁-C₆ alkoxy,
 C₁-C₆ alkyl, CF₃, S(O)_pMe, and -NMe₂, and
 a 5-10 membered heterocyclic ring containing 1-3 N, O, or
 5 S heteroatoms, wherein said heterocyclic ring may be
 saturated, partially saturated, or fully unsaturated,
 said heterocyclic ring being substituted with 0-2 R^{7e};

R^{10e} is selected from:

10 H, CF₃, C₃-C₆ alkenyl, C₃-C₁₁ cycloalkyl, aryl,
 (C₃-C₁₁ cycloalkyl)methyl, aryl(C₁-C₄ alkyl), and C₁-
 C₁₀ alkyl substituted with 0-2 R^{6e};

R^{11e} is selected from:

15 H, hydroxy, C₁-C₈ alkyl, C₃-C₆ alkenyl, C₃-C₁₁
 cycloalkyl, (C₃-C₁₁ cycloalkyl)methyl, C₁-C₆ alkoxy,
 benzyloxy, aryl, heteroaryl, heteroaryl(C₁-C₄ alkyl)-
 , aryl(C₁-C₄ alkyl), adamantylmethyl, and
 C₁-C₁₀ alkyl substituted with 0-2 R^{4e};

20

R^{4e} is selected from:

H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇
 cycloalkyl(C₁-C₄ alkyl)-, aryl, heteroaryl, aryl(C₁-
 C₆ alkyl)-, and heteroaryl(C₁-C₆ alkyl)-, wherein
 25 said aryl or heteroaryl groups are substituted with
 0-2 substituents independently selected from the
 group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, F, Cl,
 Br, CF₃, and NO₂,

30 R^{12e} is selected from:

H, C₁-C₆ alkyl, triphenylmethyl, methoxymethyl,
 methoxyphenyldiphenylmethyl,
 trimethylsilylethoxymethyl, (C₁-C₆ alkyl)carbonyl,
 (C₁-C₆ alkoxy)carbonyl, (C₁-C₆ alkyl)aminocarbonyl,
 35 C₃-C₆ alkenyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-
 C₄ alkyl)-, aryl, heteroaryl(C₁-C₆ alkyl)carbonyl,
 heteroarylcarbonyl, aryl(C₁-C₆ alkyl)-,
 (C₁-C₆ alkyl)carbonyl, arylcarbonyl, C₁-C₆

alkylsulfonyl, arylsulfonyl, aryl(C₁-C₆ alkyl)sulfonyl, heteroarylsulfonyl, heteroaryl(C₁-C₆ alkyl)sulfonyl, aryloxy carbonyl, and aryl(C₁-C₆ alkoxy)carbonyl, wherein said aryl groups are substituted with 0-2 substituents selected from the group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, halo, CF₃, and nitro;

R^{16e} is selected from:
 10 -C(=O)OR^{18ae}, -C(=O)R^{18be}, -C(=O)N(R^{18be})₂, -SO₂R^{18ae}, and -SO₂N(R^{18be})₂;

R^{17e} is selected from:
 H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-, aryl, aryl(C₁-C₆ alkyl)-, and heteroaryl(C₁-C₆ alkyl);

R^{18ae} is selected from:
 C₁-C₈ alkyl optionally substituted with a bond to L_n,
 20 C₃-C₁₁ cycloalkyl optionally substituted with a bond to L_n, aryl(C₁-C₆ alkyl)- optionally substituted with a bond to L_n, heteroaryl(C₁-C₆ alkyl)- optionally substituted with a bond to L_n, (C₁-C₆ alkyl)heteroaryl optionally substituted with a bond to L_n, biaryl(C₁-C₆ alkyl) optionally substituted with a bond to L_n, heteroaryl optionally substituted with a bond to L_n, phenyl substituted with 3-4 R^{19e} and optionally substituted with a bond to L_n,
 25 naphthyl substituted with 0-4 R^{19e} and optionally substituted with a bond to L_n, and a bond to L_n, wherein said aryl or heteroaryl groups are optionally substituted with 0-4 R^{19e};

35 R^{18be} is H or R^{18ae};

R^{19e} is selected from:

H, halogen, CF₃, CO₂H, CN, NO₂, -NR^{11e}R^{12e}, OCF₃,
 C₁-C₈ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl,
 C₃-C₁₁ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-,
 aryl(C₁-C₆ alkyl)-, C₁-C₆ alkoxy, C₁-C₄
 5 alkoxy carbonyl, aryl, aryl-O-, aryl-SO₂-,
 heteroaryl, and heteroaryl-SO₂-, wherein said aryl
 and heteroaryl groups are substituted with 0-4
 groups selected from hydrogen, halogen, CF₃, C₁-C₃
 alkyl, and C₁-C₃ alkoxy;

10

R^{20e} is selected from:

hydroxy, C₁-C₁₀ alkyloxy, C₃-C₁₁ cycloalkyloxy,
 aryloxy, aryl(C₁-C₄ alkyl)oxy,
 C₂-C₁₀ alkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 15 C₂-C₁₀ alkoxy carbonyloxy(C₁-C₂ alkyl)oxy-,
 C₂-C₁₀ alkoxy carbonyl(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkoxy carbonyloxy(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkoxy carbonyl(C₁-C₂ alkyl)oxy-,
 20 aryloxy carbonyl(C₁-C₂ alkyl)oxy-,
 aryloxy carbonyloxy(C₁-C₂ alkyl)oxy-,
 arylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 C₁-C₅ alkoxy(C₁-C₅ alkyl)carbonyloxy(C₁-C₂ alkyl)oxy,
 (5-(C₁-C₅ alkyl)-1,3-dioxa-cyclopenten-2-one-
 25 yl)methyloxy,
 (5-aryl-1,3-dioxa-cyclopenten-2-one-yl)methyloxy,
 and
 (R^{10e})(R^{11e})N-(C₁-C₁₀ alkoxy)-;

30 R^{21e} is selected from:

C₁-C₈ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl, (C₃-C₁₁
 cycloalkyl)methyl, aryl, aryl(C₁-C₄ alkyl)-, and
 C₁-C₁₀ alkyl substituted with 0-2 R^{7e};

35 R^{22e} is selected from:

-C(=O)-R^{18be}, -C(=O)N(R^{18be})₂, -C(=O)NHSO₂R^{18ae},
 -C(=O)NHC(=O)R^{18be}, -C(=O)NHC(=O)OR^{18ae}, and
 -C(=O)NHSO₂NHR^{18be};

m^e is 0-2;

n^e is 0-4; and

5

p^e is 0-2;

with the following proviso: n^e and m^e are chosen such
that the number of atoms connecting R^1 and $-\text{COR}^{20e}$ in
10 Formula (IV) is in the range of 8-14;

d is selected from 1, 2, 3, 4, and 5;

d' is 1-50;

15

W is independently selected at each occurrence from the
group: O, NH, $\text{NHC}(=\text{O})$, $\text{C}(=\text{O})\text{NH}$, $\text{NR}^8\text{C}(=\text{O})$, $\text{C}(=\text{O})\text{NR}^8$,
 $\text{C}(=\text{O})$, $\text{C}(=\text{O})\text{O}$, $\text{OC}(=\text{O})$, $\text{NHC}(=\text{S})\text{NH}$, $\text{NHC}(=\text{O})\text{NH}$, SO_2 ,
20 $(\text{OCH}_2\text{CH}_2)_s$, $(\text{CH}_2\text{CH}_2\text{O})_s$, $(\text{OCH}_2\text{CH}_2\text{CH}_2)_s$, $(\text{CH}_2\text{CH}_2\text{CH}_2\text{O})_t$,
and $(\text{aa})_t$;

aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-1
25 R^{10} , C_3 -10 cycloalkyl substituted with 0-1 R^{10} , and a
5-10 membered heterocyclic ring system containing
1-4 heteroatoms independently selected from N, S,
and O and substituted with 0-1 R^{10} ;

30 R^6 , R^{6a} , R^7 , R^{7a} , and R^8 are independently selected at
each occurrence from the group: H, $=\text{O}$, COOH , SO_3H ,
 C_1 - C_5 alkyl substituted with 0-1 R^{10} , aryl
substituted with 0-1 R^{10} , benzyl substituted with 0-1
 R^{10} , and C_1 - C_5 alkoxy substituted with 0-1 R^{10} ,
35 $\text{NHC}(=\text{O})\text{R}^{11}$, $\text{C}(=\text{O})\text{NHR}^{11}$, $\text{NHC}(=\text{O})\text{NHR}^{11}$, NHR^{11} , R^{11} , and
a bond to C_h ;

k is 0 or 1;

s is selected from 0, 1, 2, 3, 4, and 5;
 s' is selected from 0, 1, 2, 3, 4, and 5;
 s" is selected from 0, 1, 2, 3, 4, and 5;
 t is selected from 0, 1, 2, 3, 4, and 5;

5

A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ are independently
 selected at each occurrence from the group: NR¹³,
 NR¹³R¹⁴, S, SH, S(Pg), OH, and a bond to L_n;

10 E is a bond, CH, or a spacer group independently selected
 at each occurrence from the group: C₁-C₁₀ alkyl
 substituted with 0-3 R¹⁷, aryl substituted with 0-3
 R¹⁷, C₃-₁₀ cycloalkyl substituted with 0-3 R¹⁷, and a
 5-10 membered heterocyclic ring system containing
 15 1-4 heteroatoms independently selected from N, S,
 and O and substituted with 0-3 R¹⁷;

R¹³, and R¹⁴ are each independently selected from the
 group: a bond to L_n, hydrogen, C₁-C₁₀ alkyl
 20 substituted with 0-3 R¹⁷, aryl substituted with 0-3
 R¹⁷, a 5-10 membered heterocyclic ring system
 containing 1-4 heteroatoms independently selected
 from N, S, and O and substituted with 0-3 R¹⁷, and
 an electron, provided that when one of R¹³ or R¹⁴ is
 25 an electron, then the other is also an electron;


alternatively, R¹³ and R¹⁴ combine to form =C(R²⁰)(R²¹);

R¹⁷ is independently selected at each occurrence from the
 30 group: a bond to L_n, =O, F, Cl, Br, I, -CF₃, -CN,
 -CO₂R¹⁸, -C(=O)R¹⁸, -C(=O)N(R¹⁸)₂, -CH₂OR¹⁸,
 -OC(=O)R¹⁸, -OC(=O)OR^{18a}, -OR¹⁸, -OC(=O)N(R¹⁸)₂,
 -NR¹⁹C(=O)R¹⁸, -NR¹⁹C(=O)OR^{18a}, -NR¹⁹C(=O)N(R¹⁸)₂,
 -NR¹⁹SO₂N(R¹⁸)₂, -NR¹⁹SO₂R^{18a}, -SO₃H, -SO₂R^{18a},
 35 -S(=O)R^{18a}, -SO₂N(R¹⁸)₂, -N(R¹⁸)₂, -NHC(=S)NHR¹⁸,
 =NOR¹⁸, -C(=O)NHN(R¹⁸)₂, -OCH₂CO₂H, and
 2-(1-morpholino)ethoxy;

R¹⁸, R^{18a}, and R¹⁹ are independently selected at each occurrence from the group: a bond to L_n, H, and C₁-C₆ alkyl;

5 R²⁰ and R²¹ are independently selected from the group: H, C₁-C₅ alkyl, -CO₂R²⁵, C₂-C₅ 1-alkene substituted with 0-3 R²³, C₂-C₅ 1-alkyne substituted with 0-3 R²³, aryl substituted with 0-3 R²³, and unsaturated 5-10 membered heterocyclic ring system containing 1-4
10 heteroatoms independently selected from N, S, and O and substituted with 0-3 R²³;

alternatively, R²⁰ and R²¹, taken together with the divalent carbon radical to which they are attached
15 form:



20 R²² and R²³ are independently selected from the group: H, and R²⁴;


alternatively, R²², R²³ taken together form a fused aromatic or a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected
25 from N, S, and O;

R²⁴ is independently selected at each occurrence from the group: -CO₂R²⁵, -C(=O)N(R²⁵)₂, -CH₂OR²⁵, -OC(=O)R²⁵, -OR²⁵, -SO₃H, -N(R²⁵)₂, and -OCH₂CO₂H; and,
30

R²⁵ is independently selected at each occurrence from the group: H and C₁-C₃ alkyl.

[4] In an even more preferred embodiment, the present invention provides a compound including stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, or pharmaceutically acceptable salt or prodrug forms thereof wherein:

R^{1e} is selected from:



R^{2e} and R^{3e} are independently selected from:

15 H, C₁-C₄ alkoxy, NR^{11e}R^{12e}, halogen, NO₂, CN, CF₃, C₁-C₆ alkyl, C₃-C₆ alkenyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl(C₁-C₆ alkyl)-, (C₁-C₆ alkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl, arylcarbonyl, and aryl substituted with 0-4 R^{7e},

20 alternatively, when R^{2e} and R^{3e} are substituents on adjacent atoms, they can be taken together with the carbon atoms to which they are attached to form a 5-7 membered carbocyclic or 5-7 membered heterocyclic aromatic or nonaromatic ring system,

25 said carbocyclic or heterocyclic ring being substituted with 0-2 groups selected from C₁-C₄ alkyl, C₁-C₄ alkoxy, halo, cyano, amino, CF₃ and NO₂;

R^{2ae} is selected from:

30 H, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl, aryl(C₁-C₄ alkyl)-, (C₂-C₇ alkyl)carbonyl, arylcarbonyl, (C₂-C₁₀ alkoxy)carbonyl, C₃-C₇ cycloalkoxycarbonyl, C₇-C₁₁ bicycloalkoxycarbonyl, aryloxycarbonyl,

35 aryl(C₁-C₁₀ alkoxy)carbonyl,

C₁-C₆ alkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl,
 arylcarbonyloxy(C₁-C₄ alkoxy)carbonyl, and
 C₃-C₇ cycloalkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl;

5 R^{7e} is selected from:

H, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, aryl, aryl(C₁-
 C₄ alkyl)-, (C₁-C₄ alkyl)carbonyl, CO₂R^{18ae}, SO₂R^{11e},
 SO₂NR^{10e}R^{11e}, OR^{10e}, and N(R^{11e})R^{12e};

10 U^e is selected from:

-(CH₂)_n^e-, -NH(CH₂)_n^e-, -N(R^{10e})C(=O)-, and
 -NHC(=O)(CH₂)_n^e;

G^e is N or CR^{19e};

15

R^{8e} is H;

R^{9e} is selected from:

20 H, nitro, N(R^{11e})R^{12e}, OC(=O)R^{10e}, OR^{10e},
 OC(=O)NR^{10e}R^{11e}, NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},
 NR^{10e}C(=O)NR^{10e}R^{11e}, NR^{10e}SO₂NR^{10e}R^{11e},
 NR^{10e}SO₂R^{21e}, hydroxy, OR^{22e}, -N(R^{10e})R^{11e},
 -N(R^{16e})R^{17e}, aryl(C₀-C₄ alkyl)carbonyl, aryl(C₁-
 C₄ alkyl), heteroaryl(C₁-C₄ alkyl), CONR^{18ae}R^{20e},
 25 SO₂R^{18ae}, and SO₂NR^{18ae}R^{20e},

providing that any of the above alkyl, cycloalkyl,
 aryl or heteroaryl groups may be unsubstituted
 or substituted independently with 1-2 R^{7e};

30 R^{10e} is selected from:

H, CF₃, C₃-C₆ alkenyl, C₃-C₆ cycloalkyl, aryl,
 (C₃-C₆ cycloalkyl)methyl, aryl(C₁-C₄ alkyl), and
 C₁-C₄ alkyl substituted with 0-2 R^{6e};

35 R^{6e} is selected from:

H, C₁-C₄ alkyl, hydroxy, C₁-C₄ alkoxy, nitro, C₁-C₄
 alkylcarbonyl, -N(R^{11e})R^{12e}, cyano, halo, CF₃,
 CHO, CO₂R^{18be}, C(=O)R^{18be}, CONR^{17e}R^{18be},

- OC(=O)R^{10e}, OR^{10e}, OC(=O)NR^{10e}R^{11e},
 NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},
 NR^{10e}C(=O)NR^{10e}R^{11e}, NR^{10e}SO₂NR^{10e}R^{11e},
 NR^{10e}SO₂R^{21e}, S(O)_pR^{11e}, SO₂NR^{10e}R^{11e},
- 5 aryl substituted with 0-3 groups selected from
 halogen, C₁-C₄ alkoxy, C₁-C₄ alkyl, CF₃,
 S(O)_mMe, and -NMe₂,
 aryl(C₁-C₄ alkyl)-, said aryl being substituted with
 0-3 groups selected from halogen, C₁-C₄ alkoxy,
- 10 C₁-C₄ alkyl, CF₃, S(O)_pMe, and -NMe₂, and
 a 5-10 membered heterocyclic ring containing 1-3 N,
 O, or S heteroatoms, wherein said heterocyclic ring
 may be saturated, partially saturated, or fully
 unsaturated, said heterocyclic ring being
- 15 substituted with 0-2 R^{7e};

- R^{11e} is selected from:
 H, hydroxy, C₁-C₄ alkyl, C₃-C₆ alkenyl, C₃-C₆
 cycloalkyl, (C₃-C₆ cycloalkyl)methyl, C₁-C₄ alkoxy,
- 20 benzyloxy, aryl, heteroaryl, heteroaryl(C₁-C₄ alkyl)-
 , aryl(C₁-C₄ alkyl), adamantylmethyl, and
 C₁-C₄ alkyl substituted with 0-2 R^{4e};

- R^{4e} is selected from:
- 25 H, C₁-C₄ alkyl, C₃-C₇ cycloalkyl, C₃-C₇
 cycloalkyl(C₁-C₄ alkyl)-, aryl, heteroaryl, aryl(C₁-
 C₄ alkyl)-, and heteroaryl(C₁-C₄ alkyl)-, wherein
 said aryl or heteroaryl groups are substituted with
 0-2 substituents independently selected from the
- 30 group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, F, Cl,
 Br, CF₃, and NO₂,

- R^{12e} is selected from:
- 35 H, C₁-C₄ alkyl, (C₁-C₄ alkyl)carbonyl, (C₁-C₄
 alkoxy)carbonyl, phenyl(C₁-C₄ alkyl)-,
 phenylsulfonyl, phenyloxycarbonyl, and phenyl(C₁-C₄
 alkoxy)carbonyl, wherein said phenyl groups are
 substituted with 0-2 substituents selected from the

group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, halo, CF₃, and nitro;

R^{16e} is selected from:

- 5 -C(=O)OR^{18ae} -C(=O)R^{18be}, -C(=O)N(R^{18be})₂, -SO₂R^{18ae},
and -SO₂N(R^{18be})₂;

R^{17e} is selected from:

- 10 H, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, C₃-C₆
cycloalkyl(C₁-C₄ alkyl)-, aryl, aryl(C₁-C₆ alkyl)-,
and heteroaryl(C₁-C₆ alkyl);

R^{18ae} is selected from:

- 15 C₁-C₈ alkyl optionally substituted with a bond to L_n,
C₃-C₁₁ cycloalkyl optionally substituted with a
bond to L_n, aryl(C₁-C₆ alkyl)- optionally
substituted with a bond to L_n, heteroaryl(C₁-C₆
alkyl)- optionally substituted with a bond to
20 L_n, (C₁-C₆ alkyl)heteroaryl optionally
substituted with a bond to L_n, biaryl(C₁-C₆
alkyl) optionally substituted with a bond to
L_n, heteroaryl optionally substituted with a
bond to L_n, phenyl substituted with 3-4 R^{19e} and
optionally substituted with a bond to L_n,
25 naphthyl substituted with 0-4 R^{19e} and
optionally substituted with a bond to L_n, and a
bond to L_n, wherein said aryl or heteroaryl
groups are optionally substituted with 0-4 R^{19e};

- 30 R^{18be} is H or R^{18ae};

R^{19e} is selected from:

- 35 H, halogen, CF₃, CO₂H, CN, NO₂, -NR^{11e}R^{12e}, OCF₃,
C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl,
C₃-C₆ cycloalkyl, C₃-C₆ cycloalkyl(C₁-C₄ alkyl)-,
aryl(C₁-C₄ alkyl)-, C₁-C₆ alkoxy, C₁-C₄
alkoxycarbonyl, aryl, aryl-O-, aryl-SO₂-,
heteroaryl, and heteroaryl-SO₂-, wherein said aryl

and heteroaryl groups are substituted with 0-4 groups selected from hydrogen, halogen, CF₃, C₁-C₃ alkyl, and C₁-C₃ alkoxy;


- 5 R^{20e} is selected from:
 hydroxy, C₁-C₆ alkyloxy, C₃-C₆ cycloalkyloxy,
 aryloxy, aryl(C₁-C₄ alkyl)oxy,
 C₂-C₁₀ alkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 C₂-C₁₀ alkoxy carbonyloxy(C₁-C₂ alkyl)oxy-,
 10 C₂-C₁₀ alkoxy carbonyl(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkoxy carbonyloxy(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkoxy carbonyl(C₁-C₂ alkyl)oxy-,
 aryloxy carbonyl(C₁-C₂ alkyl)oxy-,
 15 aryloxy carbonyloxy(C₁-C₂ alkyl)oxy-,
 arylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 C₁-C₅ alkoxy(C₁-C₅ alkyl)carbonyloxy(C₁-C₂ alkyl)oxy,
 (5-(C₁-C₅ alkyl)-1,3-dioxo-cyclopenten-2-one-
 yl)methyloxy,
 20 (5-aryl-1,3-dioxo-cyclopenten-2-one-yl)methyloxy,
 and
 (R^{10e})(R^{11e})N-(C₁-C₁₀ alkoxy)-;

- R^{21e} is selected from:
 25 C₁-C₄ alkyl, C₂-C₆ alkenyl, C₃-C₆ cycloalkyl, (C₃-C₆
 cycloalkyl)methyl, aryl, aryl(C₁-C₄ alkyl)-, and
 C₁-C₁₀ alkyl substituted with 0-2 R^{7e};

- R^{22e} is selected from:
 30 -C(=O)-R^{18be}, -C(=O)N(R^{18be})₂, -C(=O)NHSO₂R^{18ae},
 -C(=O)NHC(=O)R^{18be}, -C(=O)NHC(=O)OR^{18ae}, and
 -C(=O)NHSO₂NHR^{18be};

- m^e is 0-2;
 35 n^e is 0-4;
 p^e is 0-2;

C_h is



5

A^1 is selected from the group: OH, and a bond to L_n ;

A^2 , A^4 , and A^6 are each N;

10 A^3 , A^5 , and A^8 are each OH;


A^7 is a bond to L_n or NH-bond to L_n ;

E is a C_2 alkyl substituted with 0-1 R^{17} ;

15

R^{17} is =O;

alternatively, C_h is



20

A^1 is selected from the group: OH, and a bond to L_n ;

25 A^2 , A^3 and A^4 are each N;

A^5 , A^6 and A^8 are each OH;

A^7 is a bond to L_n ;

30

E is a C₂ alkyl substituted with 0-1 R¹⁷;

R¹⁷ is =O;

5 alternatively, C_h is $A^1 \begin{array}{c} \diagup \\ E-A^2 \end{array}$;

A¹ is NH₂ or N=C(R²⁰)(R²¹);

E is a bond;

10

A² is NHR¹³;

R¹³ is a heterocycle substituted with R¹⁷, the heterocycle being selected from pyridine and pyrimidine;

15

R¹⁷ is selected from a bond to L_n, C(=O)NHR¹⁸ and C(=O)R¹⁸;

R¹⁸ is a bond to L_n;

20

R²⁴ is selected from the group: -CO₂R²⁵, -OR²⁵, -SO₃H, and -N(R²⁵)₂; and,

25 R²⁵ is independently selected at each occurrence from the group: hydrogen and methyl.

[5] In another even more preferred embodiment, the present invention provides a compound including enantiomeric or diastereomeric forms thereof, or mixtures of enantiomeric or diastereomeric forms thereof, or pharmaceutically acceptable salt or prodrug forms thereof, wherein Q is selected from the group:

35 3-[7-[(imidazolin-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(3,5-dimethylisoxazol-4-ylsulfonylamino)propionic acid,

- 3-[7-[(imidazolin-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
- 5 3-[7-[(imidazolin-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butyloxycarbonylamino)propionic acid,
- 3-[7-[(imidazolin-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butylsulfonylamino)propionic acid,
- 10 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
- 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butyloxycarbonylamino)propionic acid,
- 15 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(phenylsulfonylamino)propionic acid,
- 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butylsulfonyl)aminopropionic acid,
- 20 3-[7-[(2-aminothiazol-4-yl)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
- 25 3-[7-[(imidazolin-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 30 3-[7-[(imidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(3,5-dimethylisoxazol-4-ylsulfonylamino)propionic acid,
- 35

- 3-[7-[(imidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
- 5 3-[7-[(imidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 10 3-[7-[(imidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((4-biphenyl)sulfonylamino)propionic acid,
- 15 3-[7-[(imidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(1-naphthylsulfonylamino)propionic acid,
- 20 3-[7-[(benzimidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 25 3-[7-[(4-methylimidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 30 3-[7-[(4,5-dimethylimidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 35 3-[7-[(4,5,6,7-tetrahydrobenzimidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 3-[7-[(pyridin-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 3-[7-(2-aminopyridin-6-yl)-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,

- 3-[7-[(7-azabenzimidazol-2-yl)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 5 3-[7-[(benzimidazol-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]propionic acid,
- 3-[7-[(pyridin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]propionic acid,
- 10 3-[7-[(imidazolin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]propionic acid,
- 3-[7-[(imidazol-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]propionic acid,
- 15 3-[7-[(imidazolin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
- 20 3-[7-[(imidazolin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butylloxycarbonylamino)propionic acid,
- 25 3-[7-[(imidazolin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(phenylsulfonylamino)propionic acid,
- 30 3-[7-[(imidazolin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butylsulfonylamino)propionic acid,
- 35 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
- 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-

- ylcarbonylamino]-2-(n-butylloxycarbonylamino)propionic acid,
3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(phenylsulfonylamino)propionic acid,
3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butylsulfonyl)aminopropionic acid,
3-[7-[(2-aminothiazol-4-yl)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(phenylsulfonylamino)propionic acid,
3-[7-[(2-aminothiazol-4-yl)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
3-[7-[(imidazolin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
3-[7-[(imidazol-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
3-[7-[(imidazol-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(phenylsulfonylamino)propionic acid,
3-[7-[(imidazol-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,6-dichlorophenyl)sulfonylamino)propionic acid,
3-[7-[(imidazol-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-

2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
3-[7-[(imidazol-2-ylamino)methyl]-1-(2-phenylethyl)-
6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-
2-((4-biphenyl)sulfonylamino)propionic acid,
3-[7-[(benzimidazol-2-ylamino)methyl]-1-(2-
phenylethyl)-6,8-difluoroquinoline-4-one-3-
ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
3-[7-[(4-methylimidazol-2-ylamino)methyl]-1-(2-
phenylethyl)-6,8-difluoroquinoline-4-one-3-
ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
3-[7-[(4,5-dimethylimidazol-2-ylamino)methyl]-1-(2-
phenylethyl)-6,8-difluoroquinoline-4-one-3-
ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
3-[7-[(4,5,6,7-tetrahydrobenzimidazol-2-
ylamino)methyl]-1-(2-phenylethyl)-6,8-
difluoroquinoline-4-one-3-ylcarbonylamino]-2-
((2,4,6-trimethylphenyl)sulfonylamino)propionic
acid,
3-[7-[(pyridin-2-ylamino)methyl]-1-(2-phenylethyl)-
6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-
2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
3-[7-(2-aminopyridin-6-yl)-1-(2-phenylethyl)-6,8-
difluoroquinoline-4-one-3-ylcarbonylamino]-2-
((2,4,6-trimethylphenyl)sulfonylamino)propionic
acid, and
3-[7-[(7-azabenzimidazol-2-yl)methyl]-1-(2-
phenylethyl)-6,8-difluoroquinoline-4-one-3-
ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid.



[6] In another even more preferred embodiment, the present invention provides a compound selected from the group:

- 2-(((4-(4-(((3-(2-(2-(3-((6-((1-aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)-propoxy)ethoxy)-ethoxy)propyl)amino)sulfonyl)phenyl)phenyl)-sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic acid;
- 10 3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((4-(4-(((3-(2-(2-(3-(2-(1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)cyclododecyl)acetylaminopropoxy)ethoxy)ethoxy)propyl)amino)sulfonyl)-phenyl)phenyl)sulfonyl)amino)propanoic acid;
- 20 2-(((4-(3-(N-(3-(2-(2-(3-((6-((1-aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)-propoxy)ethoxy)ethoxy)propyl)carbamoyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)propanoic;
- 25 3-((1-(3-((6-((1-aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)propyl)-7-((imidazole-2-ylamino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic acid;
- 30 3-((1-(3-((6-((1-aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)propyl)-7-((1-hydroxyimidazole-2-yl)amino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic acid;
- 35 3-((1-(3-(3-(N-(3-(2-(2-(3-((6-((1-aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)propoxy)ethoxy)-ethoxy)propyl)carbamoyl)propanoylamino)propyl)-7-((imidazole-2-ylamino)methyl)-4-oxo(3-
- 40


hydroquinolyl))carbonylamino)-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic acid;

2-(2-aza-2-(5-(N-(1,3-bis(3-(2-(2-(3-(3-(N-(3-(3-(N-(3-carboxy-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)-ethyl) carbamoyl)-7-((imidazole-2-ylamino)methyl)4-oxohydroquinolyl)propyl) carbamoyl)propanoylamino)propoxy)ethoxy)ethoxy)propyl) carbamoyl)(2-pyridyl))amino)vinyl)benzenesulfonic acid;



10





15




20






5




10




- 2-(((4-(3-(N-(3-(2-(2-(3-(2-(1,4,7,10-tetraaza-4,7,10-
5 tris(carboxymethyl)cyclododecylacetyl-amino)-6-
amino)hexanoylamino)propoxy)ethoxy)ethoxy)-
propyl)carbamoyl)propoxy)-2,6-
dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-
ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-
10 carbonylamino)propanoic acid ;
- 2-(((4-(3-(N-(3-(2-(2-(3-(2-(1,4,7,10-tetraaza-4,7,10-
tris(carboxymethyl)cyclododecylacetyl-amino)-6-(2-
15 (bis(phosphonomethyl)amino)acetyl-amino)hexanoylamino
)propoxy)ethoxy)ethoxy)propyl)carbamoyl)propoxy)-
2,6-dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-
2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-
carbonylamino)propanoic acid conjugate; and
- 20 2-(((4-(3-(N-(3-(2-(2-(3-(2-(2-((2-((2-
(bis(carboxymethyl)-
amino)ethyl)(carboxymethyl)amino)ethyl)(carboxymethy
l)amino)acetyl-amino)-3-sulfopropyl)propoxy)ethoxy)-
ethoxy)propyl)carbamoyl)propoxy)-2,6-
25 dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-
ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-
carbonylamino)propanoic acid;




5






;




;

5



; and



or a pharmaceutically acceptable salt form thereof.

5

[7] In a further preferred embodiment, the present invention provides a kit comprising a compound of Claim 2, or a pharmaceutically acceptable salt form thereof and a pharmaceutically acceptable carrier.

10

[8] In an even further preferred embodiment, the kit further comprises one or more ancillary ligands and a reducing agent.

15

[9] In a still further preferred embodiment, the ancillary ligands are tricine and TPPTS.

[10] In another still further preferred embodiment, the reducing agent is tin(II).

20

[11] In a second embodiment, the present invention provides a novel diagnostic or therapeutic metallopharmaceutical composition, comprising: a metal, a chelator capable of chelating the metal and a targeting moiety, wherein the targeting moiety is bound to the chelator, is a quinolone non-peptide and binds to a

25

receptor that is upregulated during angiogenesis and the compound has 0-1 linking groups between the targeting moiety and chelator.

- 5 [12] In a preferred embodiment, wherein the metallopharmaceutical is a diagnostic radiopharmaceutical, the metal is a radioisotope selected from the group: ^{99m}Tc , ^{95}Tc , ^{111}In , ^{62}Cu , ^{64}Cu , ^{67}Ga , and ^{68}Ga , and the linking group is present between the non-
10 peptide targeting moiety and chelator.

[13] In another preferred embodiment, the targeting moiety is a quinolone non-peptide and the receptor is $\alpha_v\beta_3$ or $\alpha_v\beta_5$.

15

[14] In another preferred embodiment, the radioisotope is ^{99m}Tc or ^{95}Tc , the radiopharmaceutical further comprises a first ancillary ligand and a second ancillary ligand capable of stabilizing the radiopharmaceutical.

20

[15] In another preferred embodiment, the radioisotope is ^{99m}Tc .

[16] In another preferred embodiment, the
25 radiopharmaceutical is selected from the group:

^{99m}Tc (2-(((4-(4-((3-(2-(2-(3-((6-(diazenido)(3-pyridyl))carbonylamino)propoxy)ethoxy)-ethoxy)propyl)amino)sulfonyl)phenyl)phenyl)-sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic acid)(tricine)(TPPTS);

35 ^{99m}Tc (2-(((4-(3-(N-(3-(2-(2-(3-((6-(diazenido)(3-pyridyl))carbonylamino)propoxy)ethoxy)-ethoxy)propyl)carbamoyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-



ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-
carbonylamino)propanoic acid) (tricine) (TPPDS);



- 5 ^{99m}Tc (3-((1-(3-((6-(diazenido)(3-pyridyl))carbonylamino)propyl)-7-((imidazole-2-ylamino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic acid) (tricine) (TPPTS);
- 10 ^{99m}Tc (3-((1-(3-((6-(diazenido)(3-pyridyl))carbonylamino)propyl)-7-((1-hydroxyimidazole-2-yl)amino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic acid) (tricine) (TPPTS);
- 15 ^{99m}Tc (3-((1-(3-(3-(N-(3-(2-(2-(3-((6-(diazenido)(3-pyridyl))carbonylamino)propoxy)ethoxy)-ethoxy)propyl)carbamoyl)propanoylamino)propyl)-7-((imidazole-2-ylamino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic acid) (tricine) (TPPTS);
- 20 ^{99m}Tc (2-(2-(5-(N-(1,3-bis(3-(2-(2-(3-(3-(N-(3-(3-(N-(3-carboxy-2-((2,4,6-trimethylphenyl)sulfonyl)amino)-ethyl)carbamoyl)-7-((imidazole-2-ylamino)methyl)4-oxohydroquinolyl)propyl)carbamoyl)propanoylamino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)(2-pyridyl)diazenido)) (tricine) (TPPTS);
- 25 ^{99m}Tc (3-([1-(3-{2-[(6-(diazenido)(3-pyridyl))carbonylamino](2R)-3-sulfopropyl}propyl)-7-[(imidazol-2-ylamino)methyl]-4-oxo(3-hydroquinolyl)]carbonylamino)(2S)-2-[(2,4,6-
- 30

trimethylphenyl)sulfonyl]amino)propanoic acid)
(tricine) (TPPTS).


[17] In another preferred embodiment, the radioisotope is
5 ^{111}In .


[18] In another preferred embodiment, the
radiopharmaceutical is selected from the group:






5





; and



5

[19] In another preferred embodiment wherein the metallopharmaceutical is a therapeutic radiopharmaceutical, the metal is a radioisotope selected from the group: ¹⁸⁶Re, ¹⁸⁸Re, ¹⁵³Sm, ¹⁶⁶Ho, ¹⁷⁷Lu, ¹⁴⁹Pm, ⁹⁰Y, ²¹²Bi, ¹⁰³Pd, ¹⁰⁹Pd, ¹⁵⁹Gd, ¹⁴⁰La, ¹⁹⁸Au, ¹⁹⁹Au, ¹⁶⁹Yb, ¹⁷⁵Yb, ¹⁶⁵Dy, ¹⁶⁶Dy, ⁶⁷Cu, ¹⁰⁵Rh, ¹¹¹Ag, and ¹⁹²Ir, and the


linking group is present between the non-peptide targeting moiety and chelator.




[20] In another preferred embodiment, the targeting moiety
5 is a quinolone non-peptide and the receptor is $\alpha_v\beta_3$ or $\alpha_v\beta_5$.

[21] In another preferred embodiment, the radioisotope is
10 ^{153}Sm .

[22] In another preferred embodiment, the radioisotope is
 ^{177}Lu .

[23] In another preferred embodiment, the
15 radiopharmaceutical is selected from the group:










5

[24] In another preferred embodiment, the radioisotope is ^{90}Y .


10

[25] In another preferred embodiment, the radiopharmaceutical is selected from the group;






5



; and




[26] In another preferred embodiment wherein the
 5 metallopharmaceutical is a MRI contrast agent, the metal
 is a paramagnetic metal ion selected from the group:
 Gd(III), Dy(III), Fe(III), and Mn(II), the targeting
 moiety is a quinolone nonpeptide and the linking group is
 present between the targeting moiety and chelator.

10

[27] In another preferred embodiment, the targeting moiety
 is quinolone non-peptide and the receptor is $\alpha_v\beta_3$ or $\alpha_v\beta_5$.

[28] In another preferred embodiment, the metal ion is
 15 Gd(III).

[29] In another preferred embodiment, the contrast agent
 is



[30] In another preferred embodiment wherein the metallopharmaceutical is a X-ray contrast agent, the metal is selected from the group: Re, Sm, Ho, Lu, Pm, Y, Bi, Pd, Gd, La, Au, Au, Yb, Dy, Cu, Rh, Ag, and Ir, the targeting moiety is a quinolone non-peptide, the receptor is $\alpha_v\beta_3$ or $\alpha_v\beta_5$, and the linking group is present between the targeting moiety and chelator.

10

[31] In another preferred embodiment, the present invention provides a novel method of treating rheumatoid arthritis in a patient comprising: administering a therapeutic radiopharmaceutical of Claim 19 capable of localizing in new angiogenic vasculature to a patient by injection or infusion.

15

[32] In another preferred embodiment, the present invention provides a novel method of treating cancer in a patient comprising: administering to a patient in need thereof a therapeutic radiopharmaceutical of Claim 19 by injection or infusion.

20

[33] In another preferred embodiment, the present invention provides a novel method of imaging therapeutic angiogenesis in a patient comprising: (1) administering a diagnostic radiopharmaceutical, a MRI contrast agent,

25

or a X-ray contrast agent of Claim 11 to a patient by injection or infusion; (2) imaging the area of the patient wherein the desired formation of new blood vessels is located.

5

[34] In another preferred embodiment, the present invention provides a novel method of imaging cancer in a patient comprising: (1) administering a diagnostic radiopharmaceutical of Claim 12 to a patient by injection
10 or infusion; (2) imaging the patient using planar or SPECT gamma scintigraphy, or positron emission tomography.

[35] In another preferred embodiment, the present
15 invention provides a novel method of imaging cancer in a patient comprising: (1) administering a MRI contrast agent of Claim 26; and (2) imaging the patient using magnetic resonance imaging.


20 [36] In another preferred embodiment, the present invention provides a novel method of imaging cancer in a patient comprising: (1) administering a X-ray contrast agent of Claim 30; and (2) imaging the patient using X-ray computed tomography.

25

[37] In a third embodiment, the present invention provides a novel compound, comprising: a targeting moiety and a surfactant, wherein the targeting moiety is bound to the surfactant, is a nonpeptide, and binds to a
30 receptor that is upregulated during angiogenesis and the compound has 0-1 linking groups between the targeting moiety and surfactant.


[38] In a preferred embodiment, the targeting moiety
35 comprises a quinolone non-peptide and the linking group is present between the targeting moiety and surfactant.

[39] In another preferred embodiment, the receptor is the integrin $\alpha_v\beta_3$ or $\alpha_v\beta_5$ and the compound is of the formula:



5

wherein, Q is a compound of Formula (II):




10

(II)

including stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, or pharmaceutically acceptable salt or prodrug forms thereof wherein:

15

R^{1e} is selected from:



A^e is -CH₂- or -N(R^{10e})-;

A^{1e} and B^e are independently -CH₂- or -N(R^{10e})-;

De is -N(R^{10e})- or -S-;

$$\text{E}^{\text{e}}\text{-Fe} \text{ is } -\text{C}(\text{R}^2\text{e})=\text{C}(\text{R}^3\text{e})- \text{ or } -\text{C}(\text{R}^2\text{e})_2\text{C}(\text{R}^3\text{e})_2-;$$

15 Je is $-C(R^{2e})-$ or $-N-$;

K^e , L^e and M^e are independently $-C(R^{2e})$ - or $-C(R^{3e})$ -;

R^{2e} and R^{3e} are independently selected from:

H, C₁-C₄ alkoxy, NR^{11e}R^{12e}, halogen, NO₂, CN, CF₃,
 C₁-C₆ alkyl, C₃-C₆ alkenyl, C₃-C₇ cycloalkyl,
 C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl(C₁-C₆ alkyl)-,
 (C₁-C₆ alkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl,
 5 arylcarbonyl, and aryl substituted with 0-4 R^{7e},

alternatively, when R^{2e} and R^{3e} are substituents on
 adjacent atoms, they can be taken together with the
 carbon atoms to which they are attached to form a 5-
 10 7 membered carbocyclic or 5-7 membered heterocyclic
 aromatic or nonaromatic ring system, said
 carbocyclic or heterocyclic ring being substituted
 with 0-2 groups selected from C₁-C₄ alkyl, C₁-C₄
 alkoxy, halo, cyano, amino, CF₃ and NO₂;

15

R^{2ae} is selected from:

H, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl,
 C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl, aryl(C₁-C₄
 alkyl)-, (C₂-C₇ alkyl)carbonyl, arylcarbonyl,
 20 (C₂-C₁₀ alkoxy)carbonyl, C₃-C₇ cycloalkoxycarbonyl,
 C₇-C₁₁ bicycloalkoxycarbonyl, aryloxycarbonyl,
 aryl(C₁-C₁₀ alkoxy)carbonyl,
 C₁-C₆ alkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl,
 arylcarbonyloxy(C₁-C₄ alkoxy)carbonyl, and
 25 C₃-C₇ cycloalkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl;

R^{7e} is selected from:

H, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, aryl, aryl(C₁-
 C₄ alkyl)-, (C₁-C₄ alkyl)carbonyl, CO₂R^{18ae}, SO₂R^{11e},
 30 SO₂NR^{10e}R^{11e}, OR^{10e}, and N(R^{11e})R^{12e};

U^e is selected from:

-(CH₂)_n^e-, -(CH₂)_n^eO(CH₂)_m^e-, -(CH₂)_n^eN(R¹²)(CH₂)_m^e-,
 -NH(CH₂)_n^e-, -(CH₂)_n^eC(=O)(CH₂)_m^e-,
 35 -(CH₂)_n^eS(O)_p^e(CH₂)_m^e-, -(CH₂)_n^eNHNNH(CH₂)_m^e-,
 -N(R^{10e})C(=O)-, -NHC(=O)(CH₂)_n^e-, -C(=O)N(R^{10e})-, and
 -N(R^{10e})S(O)_p^e-;

G^e is N or CR^{19e} ;

W^e is $-C(=O)-N(R^{10e})-(C_1-C_3 \text{ alkylene})-$, in which the alkylene group is substituted by R^{8e} and by R^{9e} .

5

R^{8e} and R^{9e} are independently selected from:

- H, CO_2R^{18be} , $C(=O)R^{18be}$, $CONR^{17}R^{18be}$,
- C_1-C_{10} alkyl substituted with 0-1 R^{6e} ,
- C_2-C_{10} alkenyl substituted with 0-1 R^{6e} ,
- 10 C_2-C_{10} alkynyl substituted with 0-1 R^{6e} ,
- C_3-C_8 cycloalkyl substituted with 0-1 R^{6e} ,
- C_5-C_6 cycloalkenyl substituted with 0-1 R^{6e} ,
- $(C_1-C_{10} \text{ alkyl})\text{carbonyl}$,
- C_3-C_{10} cycloalkyl(C_1-C_4 alkyl)-,
- 15 phenyl substituted with 0-3 R^{6e} ,
- naphthyl substituted with 0-3 R^{6e} ,
- a 5-10 membered heterocyclic ring containing 1-3 N, O, or S heteroatoms, wherein said heterocyclic ring may be saturated, partially saturated, or
- 20 fully unsaturated, said heterocyclic ring being substituted with 0-2 R^{7e} ,
- C_1-C_{10} alkoxy substituted with 0-2 R^{7e} ,
- hydroxy, nitro, $-N(R^{10e})R^{11e}$, $-N(R^{16e})R^{17e}$, aryl(C_0-C_6 alkyl)carbonyl, aryl(C_3-C_6 alkyl),
- 25 heteroaryl(C_1-C_6 alkyl), $CONR^{18ae}R^{20e}$, SO_2R^{18ae} , and $SO_2NR^{18ae}R^{20e}$,
- providing that any of the above alkyl, cycloalkyl, aryl or heteroaryl groups may be unsubstituted or substituted independently with 1-2 R^{7e} ;

30

R^{6e} is selected from:

- H, C_1-C_{10} alkyl, hydroxy, C_1-C_{10} alkoxy, nitro, C_1-C_{10} alkylcarbonyl, $-N(R^{11e})R^{12e}$, cyano, halo, CF_3 ,
- CHO , CO_2R^{18be} , $C(=O)R^{18be}$, $CONR^{17e}R^{18be}$,
- 35 $OC(=O)R^{10e}$, OR^{10e} , $OC(=O)NR^{10e}R^{11e}$,
- $NR^{10e}C(=O)R^{10e}$, $NR^{10e}C(=O)OR^{21e}$,
- $NR^{10e}C(=O)NR^{10e}R^{11e}$, $NR^{10e}SO_2NR^{10e}R^{11e}$,
- $NR^{10e}SO_2R^{21e}$, $S(O)_pR^{11e}$, $SO_2NR^{10e}R^{11e}$,

35

aryl substituted with 0-3 groups selected from
halogen, C₁-C₆ alkoxy, C₁-C₆ alkyl, CF₃,
S(O)_m^eMe, and -NMe₂,
aryl(C₁-C₄ alkyl)-, said aryl being substituted with
5 0-3 groups selected from halogen, C₁-C₆ alkoxy,
C₁-C₆ alkyl, CF₃, S(O)_p^eMe, and -NMe₂, and
a 5-10 membered heterocyclic ring containing 1-3 N, O, or
S heteroatoms, wherein said heterocyclic ring may be
saturated, partially saturated, or fully unsaturated,
10 said heterocyclic ring being substituted with 0-2 R^{7e};

R^{10e} is selected from:
H, CF₃, C₃-C₆ alkenyl, C₃-C₁₁ cycloalkyl, aryl,
(C₃-C₁₁ cycloalkyl)methyl, aryl(C₁-C₄ alkyl), and C₁-
15 C₁₀ alkyl substituted with 0-2 R^{6e};

R^{11e} is selected from:
H, hydroxy, C₁-C₈ alkyl, C₃-C₆ alkenyl, C₃-C₁₁
cycloalkyl, (C₃-C₁₁ cycloalkyl)methyl, C₁-C₆ alkoxy,
20 benzyloxy, aryl, heteroaryl, heteroaryl(C₁-C₄ alkyl)-,
aryl(C₁-C₄ alkyl), adamantylmethyl, and
C₁-C₁₀ alkyl substituted with 0-2 R^{4e};

R^{4e} is selected from:
25 H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇
cycloalkyl(C₁-C₄ alkyl)-, (C₁-C₁₀ alkyl)carbonyl,
aryl, heteroaryl, aryl(C₁-C₆ alkyl)-, and
heteroaryl(C₁-C₆ alkyl)-, wherein said aryl or
heteroaryl groups are substituted with 0-2
30 substituents independently selected from the group
consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, F, Cl, Br,
CF₃, and NO₂,

alternatively, when R^{10e} and R^{11e} are both substituents on
35 the same nitrogen atom (as in -NR^{10e}R^{11e}) they may be
taken together with the nitrogen atom to which they
are attached to form a heterocycle selected from:
3-azabicyclononyl, 1,2,3,4-tetrahydro-1-quinolinyl,

1,2,3,4-tetrahydro-2-isoquinolinyl, 1-piperidinyl,
 1-morpholinyl, 1-pyrrolidinyl, thiamorpholinyl,
 thiazolidinyl, and 1-piperazinyl;
 said heterocycle being substituted with 0-3 groups
 selected from: C₁-C₆ alkyl, aryl, heteroaryl,
 aryl(C₁-C₄ alkyl)-, (C₁-C₆ alkyl)carbonyl, (C₃-C₇
 cycloalkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl, aryl(C₁-
 C₄ alkoxy)carbonyl, C₁-C₆ alkylsulfonyl, and
 arylsulfonyl;

10

R^{12e} is selected from:

H, C₁-C₆ alkyl, triphenylmethyl, methoxymethyl,
 methoxyphenyldiphenylmethyl,
 trimethylsilylethoxymethyl, (C₁-C₆ alkyl)carbonyl,
 (C₁-C₆ alkoxy)carbonyl, (C₁-C₆ alkyl)aminocarbonyl,
 C₃-C₆ alkenyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-
 C₄ alkyl)-, aryl, heteroaryl(C₁-C₆ alkyl)carbonyl,
 heteroarylcarbonyl, aryl(C₁-C₆ alkyl)-,
 (C₁-C₆ alkyl)carbonyl, arylcarbonyl, C₁-C₆
 alkylsulfonyl, arylsulfonyl, aryl(C₁-C₆
 alkyl)sulfonyl, heteroarylsulfonyl, heteroaryl(C₁-C₆
 alkyl)sulfonyl, aryloxy carbonyl, and aryl(C₁-C₆
 alkoxy)carbonyl, wherein said aryl groups are
 substituted with 0-2 substituents selected from the
 group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, halo,
 CF₃, and nitro;

25

R^{16e} is selected from:

-C(=O)OR^{18ae}, -C(=O)R^{18be}, -C(=O)N(R^{18be})₂,
 -C(=O)NHSO₂R^{18ae}, -C(=O)NHC(=O)R^{18be},
 -C(=O)NHC(=O)OR^{18ae}, -C(=O)NHSO₂NHR^{18be}, -SO₂R^{18ae},
 -SO₂N(R^{18be})₂, and -SO₂NHC(=O)OR^{18be};

30

R^{17e} is selected from:

H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇
 cycloalkyl(C₁-C₄ alkyl)-, aryl, aryl(C₁-C₆ alkyl)-,
 and heteroaryl(C₁-C₆ alkyl);

35

R^{18ae} is selected from:

- 5 C₁-C₈ alkyl optionally substituted with a bond to L_n,
 C₃-C₁₁ cycloalkyl optionally substituted with a
 bond to L_n, aryl(C₁-C₆ alkyl)- optionally
 10 substituted with a bond to L_n, heteroaryl(C₁-C₆
 alkyl)- optionally substituted with a bond to
 L_n, (C₁-C₆ alkyl)heteroaryl optionally
 substituted with a bond to L_n, biaryl(C₁-C₆
 alkyl) optionally substituted with a bond to
 15 L_n, heteroaryl optionally substituted with a
 bond to L_n, phenyl substituted with 3-4 R^{19e} and
 optionally substituted with a bond to L_n,
 naphthyl substituted with 0-4 R^{19e} and
 optionally substituted with a bond to L_n, and a
 bond to L_n, wherein said aryl or heteroaryl
 groups are optionally substituted with 0-4 R^{19e};

R^{18be} is H or R^{18ae};

20 R^{19e} is selected from:

- H, halogen, CF₃, CO₂H, CN, NO₂, -NR^{11e}R^{12e}, OCF₃,
 C₁-C₈ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl,
 C₃-C₁₁ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-,
 aryl(C₁-C₆ alkyl)-, C₁-C₆ alkoxy, C₁-C₄
 25 alkoxy carbonyl, aryl, aryl-O-, aryl-SO₂-,
 heteroaryl, and heteroaryl-SO₂-, wherein said aryl
 and heteroaryl groups are substituted with 0-4
 groups selected from hydrogen, halogen, CF₃, C₁-C₃
 alkyl, and C₁-C₃ alkoxy;

30

R^{20e} is selected from:


- hydroxy, C₁-C₁₀ alkyloxy, C₃-C₁₁ cycloalkyloxy,
 aryloxy, aryl(C₁-C₄ alkyl)oxy,
 C₂-C₁₀ alkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 35 C₂-C₁₀ alkoxy carbonyloxy(C₁-C₂ alkyl)oxy-,
 C₂-C₁₀ alkoxy carbonyl(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkoxy carbonyloxy(C₁-C₂ alkyl)oxy-,

- C₃-C₁₀ cycloalkoxycarbonyl(C₁-C₂ alkyl)oxy-,
 aryloxy carbonyl(C₁-C₂ alkyl)oxy-,
 aryloxy carbonyloxy(C₁-C₂ alkyl)oxy-,
 arylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 5 C₁-C₅ alkoxy(C₁-C₅ alkyl)carbonyloxy(C₁-C₂ alkyl)oxy,
 (5-(C₁-C₅ alkyl)-1,3-dioxo-cyclopenten-2-one-
 yl)methyloxy,
 (5-aryl-1,3-dioxo-cyclopenten-2-one-yl)methyloxy,
 and
 10 (R^{10e}) (R^{11e})N-(C₁-C₁₀ alkoxy)-;

- R^{21e} is selected from:
 C₁-C₈ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl, (C₃-C₁₁
 cycloalkyl)methyl, aryl, aryl(C₁-C₄ alkyl)-, and C₁-
 15 C₁₀ alkyl substituted with 0-2 R^{7e};

- R^{22e} is selected from:
 -C(=O)-R^{18be}, -C(=O)N(R^{18be})₂, -C(=O)NHSO₂R^{18ae}, -
 C(=O)NHC(=O)R^{18be}, -C(=O)NHC(=O)OR^{18ae}, and -
 20 C(=O)NHSO₂NHR^{18be};

- Y^e is selected from:
 -COR^{20e}, -SO₃H, -PO₃H, -CONHNHSO₂CF₃, -CONHSO₂R^{18ae},
 -CONHSO₂NHR^{18be}, -NHCOCF₃, -NHCONHSO₂R^{18ae},
 25 -NHSO₂R^{18ae}, -OPO₃H₂, -OSO₃H, -PO₃H₂, -SO₂NHCOR^{18ae},
 -SO₂NHCO₂R^{18ae},



- 30 m^e is 0-2;

n^e is 0-4;

p^e is 0-2;

35

r^e is 0-2;

with the following proviso: n^e and m^e are chosen such that the number of atoms connecting R^{1e} and Y^e is in the range of 8-14;

5

d is selected from 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

L_n is a linking group having the formula:

10 $((W)_h - (CR^6R^7)_g)_x - (Z)_k - ((CR^{6a}R^{7a})_{g'} - (W)_{h'})_{x'}$;

W is independently selected at each occurrence from the group: O, S, NH, NHC(=O), C(=O)NH, $NR^8C(=O)$, C(=O)N R^8 , C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂,
 15 SO₂NH, (OCH₂CH₂)₂₀₋₂₀₀, (CH₂CH₂O)₂₀₋₂₀₀, (OCH₂CH₂CH₂)₂₀₋₂₀₀, (CH₂CH₂CH₂O)₂₀₋₂₀₀, and (aa)_t;

aa is independently at each occurrence an amino acid;

20 Z is selected from the group: aryl substituted with 0-3 R^{10} , C₃₋₁₀ cycloalkyl substituted with 0-3 R^{10} , and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R^{10} ;

25

R^6 , R^{6a} , R^7 , R^{7a} , and R^8 are independently selected at each occurrence from the group: H, =O, COOH, SO₃H, PO₃H, C₁₋₅ alkyl substituted with 0-3 R^{10} , aryl substituted with 0-3 R^{10} , benzyl substituted with 0-3 R^{10} , and C₁₋₅ alkoxy substituted with 0-3 R^{10} ,
 30 NHC(=O) R^{11} , C(=O)NHR¹¹, NHC(=O)NHR¹¹, NHR¹¹, R^{11} , and a bond to S_f ;

35 R^{10} is independently selected at each occurrence from the group: a bond to S_f , COOR¹¹, C(=O)NHR¹¹, NHC(=O) R^{11} , OH, NHR¹¹, SO₃H, PO₃H, -OPO₃H₂, -OSO₃H, aryl substituted with 0-3 R^{11} , C₁₋₅ alkyl substituted with 0-1 R^{12} , C₁₋₅ alkoxy substituted with 0-1 R^{12} , and a

5-10 membered heterocyclic ring system containing
1-4 heteroatoms independently selected from N, S,
and O and substituted with 0-3 R¹¹;

- 5 R¹¹ is independently selected at each occurrence from the
group: H, alkyl substituted with 0-1 R¹², aryl
substituted with 0-1 R¹², a 5-10 membered
heterocyclic ring system containing 1-4 heteroatoms
independently selected from N, S, and O and
10 substituted with 0-1 R¹², C₃₋₁₀ cycloalkyl
substituted with 0-1 R¹², polyalkylene glycol
substituted with 0-1 R¹², carbohydrate substituted
with 0-1 R¹², cyclodextrin substituted with 0-1 R¹²,
amino acid substituted with 0-1 R¹², polycarboxyalkyl
15 substituted with 0-1 R¹², polyazaalkyl substituted
with 0-1 R¹², peptide substituted with 0-1 R¹²,
wherein the peptide is comprised of 2-10 amino
acids, 3,6-O-disulfo-B-D-galactopyranosyl,
bis(phosphonomethyl)glycine, and a bond to S_f;

20 R¹² is a bond to S_f;

- k is selected from 0, 1, and 2;
h is selected from 0, 1, and 2;
25 h' is selected from 0, 1, and 2;
g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
x is selected from 0, 1, 2, 3, 4, and 5;
30 x' is selected from 0, 1, 2, 3, 4, and 5;

S_f is a surfactant which is a lipid or a compound of the

formula: $A^9 \overset{E^1}{\text{---}} A^{10}$;

35

A⁹ is selected from the group: OH and OR²⁷;

A¹⁰ is OR²⁷;

R²⁷ is C(=O)C₁₋₂₀ alkyl;

E¹ is C₁₋₁₀ alkylene substituted with 1-3 R²⁸;

5

R²⁸ is independently selected at each occurrence from the group: R³⁰, -PO₃H-R³⁰, =O, -CO₂R²⁹, -C(=O)R²⁹, -C(=O)N(R²⁹)₂, -CH₂OR²⁹, -OR²⁹, -N(R²⁹)₂, C₁₋₅ alkyl, and C₂₋₄ alkenyl;

10

R²⁹ is independently selected at each occurrence from the group: R³⁰, H, C₁₋₆ alkyl, phenyl, benzyl, and trifluoromethyl;

15 R³⁰ is a bond to L_n;


and a pharmaceutically acceptable salt thereof.

[40] In another preferred embodiment, the compound is of the formula:



wherein, Q is a compound of Formula (IV):


25



(IV)

30 including stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, or pharmaceutically acceptable salt or prodrug forms thereof wherein:

R^{1e} is selected from:



R^{2e} and R^{3e} are independently selected from:

15 H, C₁-C₄ alkoxy, NR^{11e}R^{12e}, halogen, NO₂, CN, CF₃, C₁-C₆ alkyl, C₃-C₆ alkenyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl(C₁-C₆ alkyl)-, (C₁-C₆ alkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl, arylcarbonyl, and aryl substituted with 0-4 R^{7e},

20 alternatively, when R^{2e} and R^{3e} are substituents on adjacent atoms, they can be taken together with the carbon atoms to which they are attached to form a 5-7 membered carbocyclic or 5-7 membered heterocyclic aromatic or nonaromatic ring system, said carbocyclic or heterocyclic ring being substituted
25 with 0-2 groups selected from C₁-C₄ alkyl, C₁-C₄ alkoxy, halo, cyano, amino, CF₃ and NO₂;

R^{2ae} is selected from:

30 H, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl,

- aryl(C₁-C₄ alkyl)-, (C₂-C₇ alkyl)carbonyl,
 arylcarbonyl,
 (C₂-C₁₀ alkoxy)carbonyl, C₃-C₇ cycloalkoxycarbonyl,
 C₇-C₁₁ bicycloalkoxycarbonyl, aryloxycarbonyl,
 5 aryl(C₁-C₁₀ alkoxy)carbonyl,
 C₁-C₆ alkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl,
 arylcarbonyloxy(C₁-C₄ alkoxy)carbonyl, and
 C₃-C₇ cycloalkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl;
- 10 R^{7e} is selected from:
 H, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, aryl, aryl(C₁-
 C₄ alkyl)-, (C₁-C₄ alkyl)carbonyl, CO₂R^{18ae}, SO₂R^{11e},
 SO₂NR^{10e}R^{11e}, OR^{10e}, and N(R^{11e})R^{12e};
- 15 U^e is selected from:
 -(CH₂)_n^e-, -(CH₂)_n^eO(CH₂)_m^e-, -NH(CH₂)_n^e-,
 -N(R^{10e})C(=O)-, -NHC(=O)(CH₂)_n^e-, and -C(=O)N(R^{10e})-;
- G^e is N or CR^{19e};
- 20 R^{8e} is selected from:
 H, CO₂R^{18be}, C(=O)R^{18be}, CONR^{17e}R^{18be},
 C₁-C₁₀ alkyl substituted with 0-1 R^{6e},
 C₂-C₁₀ alkenyl substituted with 0-1 R^{6e},
 25 C₂-C₁₀ alkynyl substituted with 0-1 R^{6e},
 C₃-C₈ cycloalkyl substituted with 0-1 R^{6e},
 C₅-C₆ cycloalkenyl substituted with 0-1 R^{6e},
 (C₁-C₁₀ alkyl)carbonyl,
 C₃-C₁₀ cycloalkyl(C₁-C₄ alkyl)-,
 30 phenyl substituted with 0-3 R^{6e},
 naphthyl substituted with 0-3 R^{6e},
 a 5-10 membered heterocyclic ring containing 1-3 N,
 O, or S heteroatoms, wherein said heterocyclic
 ring may be saturated, partially saturated, or
 35 fully unsaturated, said heterocyclic ring being
 substituted with 0-2 R^{7e};

R^{9e} is selected from:

- C₁-C₁₀ alkyl substituted with 0-1 R^{6e},
 C₁-C₁₀ alkoxy substituted with 0-2 R^{7e},
 H, nitro, N(R^{11e})R^{12e}, OC(=O)R^{10e}, OR^{10e},
 OC(=O)NR^{10e}R^{11e}, NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},
 5 NR^{10e}C(=O)NR^{10e}R^{11e}, NR^{10e}SO₂NR^{10e}R^{11e},
 NR^{10e}SO₂R^{21e}, hydroxy, OR^{22e}, -N(R^{10e})R^{11e}, -
 N(R^{16e})R^{17e}, aryl(C₀-C₆ alkyl)carbonyl, aryl(C₁-
 C₆ alkyl), heteroaryl(C₁-C₆ alkyl), CONR^{18ae}R^{20e},
 SO₂R^{18ae}, and SO₂NR^{18ae}R^{20e},
 10 providing that any of the above alkyl, cycloalkyl,
 aryl or heteroaryl groups may be unsubstituted
 or substituted independently with 1-2 R^{7e};
- R^{6e} is selected from:
- 15 H, C₁-C₁₀ alkyl, hydroxy, C₁-C₁₀ alkoxy, nitro, C₁-C₁₀
 alkylcarbonyl, -N(R^{11e})R^{12e}, cyano, halo, CF₃,
 CHO, CO₂R^{18be}, C(=O)R^{18be}, CONR^{17e}R^{18be},
 OC(=O)R^{10e}, OR^{10e}, OC(=O)NR^{10e}R^{11e},
 NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},
 20 NR^{10e}C(=O)NR^{10e}R^{11e}, NR^{10e}SO₂NR^{10e}R^{11e},
 NR^{10e}SO₂R^{21e}, S(O)_pR^{11e}, SO₂NR^{10e}R^{11e},
 aryl substituted with 0-3 groups selected from
 halogen, C₁-C₆ alkoxy, C₁-C₆ alkyl, CF₃,
 S(O)_mMe, and -NMe₂,
 25 aryl(C₁-C₄ alkyl)-, said aryl being substituted with
 0-3 groups selected from halogen, C₁-C₆ alkoxy,
 C₁-C₆ alkyl, CF₃, S(O)_pMe, and -NMe₂, and
 a 5-10 membered heterocyclic ring containing 1-3 N, O, or
 S heteroatoms, wherein said heterocyclic ring may be
 30 saturated, partially saturated, or fully unsaturated,
 said heterocyclic ring being substituted with 0-2 R^{7e};

R^{10e} is selected from:

- 35 H, CF₃, C₃-C₆ alkenyl, C₃-C₁₁ cycloalkyl, aryl,
 (C₃-C₁₁ cycloalkyl)methyl, aryl(C₁-C₄ alkyl), and C₁-
 C₁₀ alkyl substituted with 0-2 R^{6e};

R^{11e} is selected from:

H, hydroxy, C₁-C₈ alkyl, C₃-C₆ alkenyl, C₃-C₁₁ cycloalkyl, (C₃-C₁₁ cycloalkyl)methyl, C₁-C₆ alkoxy, benzyloxy, aryl, heteroaryl, heteroaryl(C₁-C₄ alkyl)-, aryl(C₁-C₄ alkyl), adamantylmethyl, and
5 C₁-C₁₀ alkyl substituted with 0-2 R^{4e};

R^{4e} is selected from:

H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-, aryl, heteroaryl, aryl(C₁-C₆ alkyl)-, and heteroaryl(C₁-C₆ alkyl)-, wherein
10 said aryl or heteroaryl groups are substituted with 0-2 substituents independently selected from the group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, F, Cl, Br, CF₃, and NO₂,

15

R^{12e} is selected from:

H, C₁-C₆ alkyl, triphenylmethyl, methoxymethyl, methoxyphenyldiphenylmethyl, trimethylsilylethoxymethyl, (C₁-C₆ alkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl, (C₁-C₆ alkyl)aminocarbonyl, C₃-C₆ alkenyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-, aryl, heteroaryl(C₁-C₆ alkyl)carbonyl, heteroarylcarbonyl, aryl(C₁-C₆ alkyl)-, (C₁-C₆ alkyl)carbonyl, arylcarbonyl, C₁-C₆ alkylsulfonyl, arylsulfonyl, aryl(C₁-C₆ alkyl)sulfonyl, heteroarylsulfonyl, heteroaryl(C₁-C₆ alkyl)sulfonyl, aryloxycarbonyl, and aryl(C₁-C₆ alkoxy)carbonyl, wherein said aryl groups are
20 substituted with 0-2 substituents selected from the group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, halo, CF₃, and nitro;
25
30

R^{16e} is selected from:

-C(=O)OR^{18ae}, -C(=O)R^{18be}, -C(=O)N(R^{18be})₂, -SO₂R^{18ae},
35 and -SO₂N(R^{18be})₂;

R^{17e} is selected from:

H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-, aryl, aryl(C₁-C₆ alkyl)-, and heteroaryl(C₁-C₆ alkyl);

5 R^{18ae} is selected from:

C₁-C₈ alkyl optionally substituted with a bond to L_n,
 C₃-C₁₁ cycloalkyl optionally substituted with a
 bond to L_n, aryl(C₁-C₆ alkyl)- optionally
 substituted with a bond to L_n, heteroaryl(C₁-C₆
 10 alkyl)- optionally substituted with a bond to
 L_n, (C₁-C₆ alkyl)heteroaryl optionally
 substituted with a bond to L_n, biaryl(C₁-C₆
 alkyl) optionally substituted with a bond to
 L_n, heteroaryl optionally substituted with a
 15 bond to L_n, phenyl substituted with 3-4 R^{19e} and
 optionally substituted with a bond to L_n,
 naphthyl substituted with 0-4 R^{19e} and
 optionally substituted with a bond to L_n, and a
 bond to L_n, wherein said aryl or heteroaryl
 20 groups are optionally substituted with 0-4 R^{19e};

R^{18be} is H or R^{18ae};

R^{19e} is selected from:

25 H, halogen, CF₃, CO₂H, CN, NO₂, -NR^{11e}R^{12e}, OCF₃,
 C₁-C₈ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl,
 C₃-C₁₁ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-,
 aryl(C₁-C₆ alkyl)-, C₁-C₆ alkoxy, C₁-C₄
 alkoxy carbonyl, aryl, aryl-O-, aryl-SO₂-,
 30 heteroaryl, and heteroaryl-SO₂-, wherein said aryl
 and heteroaryl groups are substituted with 0-4
 groups selected from hydrogen, halogen, CF₃, C₁-C₃
 alkyl, and C₁-C₃ alkoxy;

35 R^{20e} is selected from:

hydroxy, C₁-C₁₀ alkyloxy, C₃-C₁₁ cycloalkyloxy,
 aryloxy, aryl(C₁-C₄ alkyl)oxy,
 C₂-C₁₀ alkylcarbonyloxy(C₁-C₂ alkyl)oxy-,

C₂-C₁₀ alkoxy carbonyloxy(C₁-C₂ alkyl)oxy-,
 C₂-C₁₀ alkoxy carbonyl(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkyl carbonyloxy(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkoxy carbonyloxy(C₁-C₂ alkyl)oxy-,
 5 C₃-C₁₀ cycloalkoxy carbonyl(C₁-C₂ alkyl)oxy-,
 aryloxy carbonyl(C₁-C₂ alkyl)oxy-,
 aryloxy carbonyloxy(C₁-C₂ alkyl)oxy-,
 aryl carbonyloxy(C₁-C₂ alkyl)oxy-,
 C₁-C₅ alkoxy(C₁-C₅ alkyl) carbonyloxy(C₁-C₂ alkyl)oxy,
 10 (5-(C₁-C₅ alkyl)-1,3-dioxo-cyclopenten-2-one-
 yl)methyloxy,
 (5-aryl-1,3-dioxo-cyclopenten-2-one-yl)methyloxy,
 and
 (R^{10e})(R^{11e})N-(C₁-C₁₀ alkoxy)-;

15

R^{21e} is selected from:

C₁-C₈ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl, (C₃-C₁₁
 cycloalkyl)methyl, aryl, aryl(C₁-C₄ alkyl)-, and
 C₁-C₁₀ alkyl substituted with 0-2 R^{7e};

20

R^{22e} is selected from:

-C(=O)-R^{18be}, -C(=O)N(R^{18be})₂, -C(=O)NHSO₂R^{18ae},
 -C(=O)NHC(=O)R^{18be}, -C(=O)NHC(=O)OR^{18ae}, and
 -C(=O)NHSO₂NHR^{18be};

25

m^e is 0-2;

n^e is 0-4; and

30 p^e is 0-2;

with the following proviso: n^e and m^e are chosen such
 that the number of atoms connecting R¹ and -COR^{20e} in
 Formula (IV) is in the range of 8-14;

35

W is independently selected at each occurrence from the
 group: O, S, NH, NHC(=O), C(=O)NH, NR⁸C(=O), C(=O)N R⁸,
 C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, SO₂NH,

(OCH₂CH₂)₂₀₋₂₀₀, (CH₂CH₂O)₂₀₋₂₀₀, (OCH₂CH₂CH₂)₂₀₋₂₀₀,
(CH₂CH₂CH₂O)₂₀₋₂₀₀, and (aa)_t;

aa is independently at each occurrence an amino acid;

5

Z is selected from the group: aryl substituted with 0-1
R¹⁰, C₃₋₁₀ cycloalkyl substituted with 0-1 R¹⁰, and a
5-10 membered heterocyclic ring system containing
1-4 heteroatoms independently selected from N, S,
10 and O and substituted with 0-1 R¹⁰;

R⁶, R^{6a}, R⁷, R^{7a}, and R⁸ are independently selected at
each occurrence from the group: H, =O, COOH, SO₃H,
C₁₋₅ alkyl substituted with 0-1 R¹⁰, aryl
15 substituted with 0-1 R¹⁰, benzyl substituted with 0-1
R¹⁰, and C₁₋₅ alkoxy substituted with 0-1 R¹⁰,
NHC(=O)R¹¹, C(=O)NHR¹¹, NHC(=O)NHR¹¹, NHR¹¹, R¹¹, and
a bond to S_f;

20 k is 0 or 1;

S_f is a surfactant which is a lipid or a compound of the

formula: $A^9-E^1-A^{10}$;

25

A⁹ is OR²⁷;

A¹⁰ is OR²⁷;

30 R²⁷ is C(=O)C₁₋₁₅ alkyl;

E¹ is C₁₋₄ alkylene substituted with 1-3 R²⁸;

R²⁸ is independently selected at each occurrence from the
35 group: R³⁰, -PO₃H-R³⁰, =O, -CO₂R²⁹, -C(=O)R²⁹,
-CH₂OR²⁹, -OR²⁹, and C₁₋₅ alkyl;

R²⁹ is independently selected at each occurrence from the group: R³⁰, H, C₁-C₆ alkyl, phenyl, and benzyl;

R³⁰ is a bond to L_n;

5

and a pharmaceutically acceptable salt thereof.

[41] In another more preferred embodiment, the present invention provides a novel ultrasound contrast agent composition, comprising:

- 10 (a) a compound of Claim 39, comprising: an quinolone that binds to the integrin $\alpha_v\beta_3$, a surfactant and a linking group between the quinolone and the surfactant;
- 15 (b) a parenterally acceptable carrier; and,
- (c) an echogenic gas.

[42] In another more preferred embodiment, the present invention provides a novel ultrasound contrast agent composition, further comprising: 1,2-dipalmitoyl-sn-glycero-3-phosphotidic acid, 1,2-dipalmitoyl-sn-glycero-3-phosphatidylcholine, and N-(methoxypolyethylene glycol 5000 carbamoyl)-1,2-dipalmitoyl-sn-glycero-3-phosphatidylethanolamine.

25

[43] In another more preferred embodiment, the echogenic gas is a C₂₋₅ perfluorocarbon.

[44] In another preferred embodiment, the present invention provides a method of imaging cancer in a patient comprising: (1) administering, by injection or infusion, a ultrasound contrast agent composition of Claim 41 to a patient; and (2) imaging the patient using sonography.

35

[45] In another preferred embodiment, the present invention provides a method of imaging formation of new blood vessels in a patient comprising: (1)

administering, by injection or infusion, a ultrasound contrast agent composition of of Claim 41 to a patient; (2) imaging the area of the patient wherein the desired formation of new blood vessels is located.

5

[46] In another preferred embodiment, the present invention provides a method of imaging therapeutic angiogenesis in a patient comprising: (1) administering, by injection or infusion, an ultrasound contrast agent composition of Claim 41 to a patient; (2) imaging the area of the patient wherein the desired formation of new blood vessels is located.

15

[47] In another preferred embodiment, the present invention provides a method of imaging atherosclerosis in a patient comprising: (1) administering, by injection or infusion, an ultrasound contrast agent composition of Claim 41 to a patient; (2) imaging the area of the patient wherein the atherosclerosis is located.

20

[48] In another preferred embodiment, the present invention provides a method of imaging restenosis in a patient comprising: (1) administering, by injection or infusion, an ultrasound contrast agent composition of Claim 41 to a patient; (2) imaging the area of the patient wherein the restenosis is located.

30

[49] In another preferred embodiment, the present invention provides a method of imaging cardiac ischemia in a patient comprising: (1) administering, by injection or infusion, an ultrasound contrast agent composition of Claim 41 to a patient; (2) imaging the area of the myocardium wherein the ischemic region is located.

35

[50] In another preferred embodiment, the present invention provides a method of imaging myocardial reperfusion injury in a patient comprising: (1) administering, by injection or infusion, an ultrasound

contrast agent composition of Claim 41 to a patient; (2) imaging the area of myocardium wherein the reperfusion injury is located.

- 5 [51] In another preferred embodiment, the present invention provides a novel therapeutic radiopharmaceutical composition, comprising:
- (a) a therapeutic radiopharmaceutical of Claim 19;
 - and,
 - 10 (b) a parenterally acceptable carrier.

[52] In another preferred embodiment, the present invention provides a novel diagnostic pharmaceutical composition, comprising:

- 15 (a) a diagnostic radiopharmaceutical, a MRI contrast agent, or a X-ray contrast agent of Claim 11; and,
- (b) a parenterally acceptable carrier.

- [53] In another preferred embodiment, the present invention provides a method of treating restenosis in a patient comprising: administering to a patient, either systemically or locally, a therapeutic radiopharmaceutical of Claim 19 capable of localizing in the restenotic area and delivering an effective dose of radiation.
- 20
- 25

- [54] In another preferred embodiment, the present invention provides a method of imaging atherosclerosis in a patient comprising: (1) administering a diagnostic radiopharmaceutical, a MRI contrast agent, or a X-ray contrast agent of Claim 11 to a patient by injection or infusion; (2) imaging the area of the patient wherein the atherosclerosis is located.
- 30

- 35 [55] In another preferred embodiment, the present invention provides a method of imaging restenosis in a patient comprising: (1) administering a diagnostic radiopharmaceutical, a MRI contrast agent, or a X-ray

contrast agent of Claim 11 to a patient by injection or infusion; (2) imaging the area of the patient wherein the restenosis is located.

- 5 [56] In another preferred embodiment, the present invention provides a method of imaging cardiac ischemia in a patient comprising: (1) administering a diagnostic radiopharmaceutical, a MRI contrast agent, or a X-ray contrast agent of Claim 11 to a patient by injection or
10 infusion; (2) imaging the area of the myocardium wherein the ischemic region is located.

- [57] In another preferred embodiment, the present invention provides a method of imaging myocardial
15 reperfusion injury in a patient comprising: (1) administering a diagnostic radiopharmaceutical, a MRI contrast agent, or a X-ray contrast agent of Claim 11 to a patient by injection or infusion; (2) imaging the area of myocardium wherein the reperfusion injury is located.

- 20 Another aspect of the present invention are diagnostic kits for the preparation of radiopharmaceuticals useful as imaging agents for cancer. Diagnostic kits of the present invention comprise one or
25 more vials containing the sterile, non-pyrogenic, formulation comprised of a predetermined amount of a reagent of the present invention, and optionally other components such as one or two ancillary ligands, reducing agents, transfer ligands, buffers, lyophilization aids,
30 stabilization aids, solubilization aids and bacteriostats. The inclusion of one or more optional components in the formulation will frequently improve the ease of synthesis of the radiopharmaceutical by the practicing end user, the ease of manufacturing the kit,
35 the shelf-life of the kit, or the stability and shelf-life of the radiopharmaceutical. The inclusion of one or two ancillary ligands is required for diagnostic kits comprising reagent comprising a hydrazine or hydrazone bonding moiety. The one or more vials that

contain all or part of the formulation can independently be in the form of a sterile solution or a lyophilized solid.

Another aspect of the present invention contemplates
5 a method of imaging cancer in a patient involving: (1)
synthesizing a diagnostic radiopharmaceutical of the
present invention, using a reagent of the present
invention, capable of localizing in tumors; (2)
administering said radiopharmaceutical to a patient by
10 injection or infusion; (3) imaging the patient using
planar or SPECT gamma scintigraphy, or positron emission
tomography.

Another aspect of the present invention contemplates
a method of imaging cancer in a patient involving: (1)
15 administering a paramagnetic metallopharmaceutical of the
present invention capable of localizing in tumors to a
patient by injection or infusion; and (2) imaging the
patient using magnetic resonance imaging.

Another aspect of the present invention contemplates
20 a method of imaging cancer in a patient involving: (1)
administering a X-ray contrast agent of the present
invention capable of localizing in tumors to a patient by
injection or infusion; and (2) imaging the patient using
X-ray computed tomography.

25 Another aspect of the present invention contemplates
a method of imaging cancer in a patient involving: (1)
administering a ultrasound contrast agent of the present
invention capable of localizing in tumors to a patient by
injection or infusion; and (2) imaging the patient using
30 sonography.

Another aspect of the present invention contemplates
a method of treating cancer in a patient involving: (1)
administering a therapeutic radiopharmaceutical of the
present invention capable of localizing in tumors to a
35 patient by injection or infusion.

DEFINITIONS

The compounds herein described may have asymmetric centers. Unless otherwise indicated, all chiral, diastereomeric and racemic forms are included in the present invention. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. It will be appreciated that compounds of the present invention contain asymmetrically substituted carbon atoms, and may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically active starting materials. Two distinct isomers (cis and trans) of the peptide bond are known to occur; both can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. The D and L-isomers of a particular amino acid are designated herein using the conventional 3-letter abbreviation of the amino acid, as indicated by the following examples: D-Leu, or L-Leu.

When any variable occurs more than one time in any substituent or in any formula, its definition on each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R^{52} , then said group may optionally be substituted with up to two R^{52} , and R^{52} at each occurrence is selected independently from the defined list of possible R^{52} . Also, by way of example, for the group $-N(R^{53})_2$, each of the two R^{53} substituents on N is independently selected from the defined list of possible R^{53} . Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. When a bond to a substituent is shown to cross the bond connecting two atoms in a

ring, then such substituent may be bonded to any atom on the ring.

The term "nonpeptide" means preferably less than three amide bonds in the backbone core of the targeting moiety or preferably less than three amino acids or amino acid mimetics in the targeting moiety.

The term "metallopharmaceutical" means a pharmaceutical comprising a metal. The metal is the cause of the imageable signal in diagnostic applications and the source of the cytotoxic radiation in radiotherapeutic applications. Radiopharmaceuticals are metallopharmaceuticals in which the metal is a radioisotope.

By "reagent" is meant a compound of this invention capable of direct transformation into a metallopharmaceutical of this invention. Reagents may be utilized directly for the preparation of the metallopharmaceuticals of this invention or may be a component in a kit of this invention.

The term "binding agent" means a metallopharmaceutical of this invention having affinity for and capable of binding to the vitronectin receptor. The binding agents of this invention have $K_i < 1000\text{nM}$.

By "stable compound" or "stable structure" is meant herein a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious pharmaceutical agent.

The term "substituted", as used herein, means that one or more hydrogens on the designated atom or group is replaced with a selection from the indicated group, provided that the designated atom's or group's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e., $=O$), then 2 hydrogens on the atom are replaced.

The term "bond", as used herein, means either a single or double bond.

The term "salt", as used herein, is used as defined in the CRC Handbook of Chemistry and Physics, 65th

Edition, CRC Press, Boca Raton, Fla, 1984, as any substance which yields ions, other than hydrogen or hydroxyl ions. As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds modified by making acid or base salts.

Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared
5 by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or
10 acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the disclosure of which is hereby incorporated by reference.

As used herein, "alkyl" is intended to include both
15 branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, examples of which include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, sec-butyl, t-butyl, pentyl, hexyl, heptyl, octyl, nonyl,
20 and decyl; "cycloalkyl" or "carbocycle" is intended to include saturated and partially unsaturated ring groups, including mono-, bi- or poly-cyclic ring systems, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl and adamantyl; "bicycloalkyl" or
25 "bicyclic" is intended to include saturated bicyclic ring groups such as [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, and so forth.

As used herein, the term "alkene" or "alkenyl" is
30 intended to include hydrocarbon chains having the specified number of carbon atoms of either a straight or branched configuration and one or more unsaturated carbon-carbon bonds which may occur in any stable point along the chain, such as ethenyl, propenyl, and the like.

35 As used herein, the term "alkyne" or "alkynyl" is intended to include hydrocarbon chains having the specified number of carbon atoms of either a straight or branched configuration and one or more unsaturated

carbon-carbon triple bonds which may occur in any stable point along the chain, such as propargyl, and the like.

As used herein, "aryl" or "aromatic residue" is intended to mean phenyl or naphthyl, which when
5 substituted, the substitution can be at any position.

As used herein, the term "heterocycle" or "heterocyclic system" is intended to mean a stable 5- to 7- membered monocyclic or bicyclic or 7- to 10-membered bicyclic heterocyclic ring which is saturated partially
10 unsaturated or unsaturated (aromatic), and which consists of carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O and S and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene
15 ring. The nitrogen and sulfur heteroatoms may optionally be oxidized. The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a
20 nitrogen atom if the resulting compound is stable. If specifically noted, a nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one
25 another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. As used herein, the term "aromatic heterocyclic system" is intended to mean a stable 5- to 7- membered monocyclic or bicyclic or 7- to 10-membered bicyclic heterocyclic
30 aromatic ring which consists of carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O and S. It is preferred that the total number of S and O atoms in the aromatic heterocycle is not more than 1.

35 Examples of heterocycles include, but are not limited to, 1H-indazole, 2-pyrrolidonyl, 2H,6H-1,5,2-dithiazinyl, 2H-pyrrolyl, 3H-indolyl, 4-piperidonyl, 4aH-carbazole, 4H-quinoliziny, 6H-1,2,5-thiadiazinyl, acridinyl, azocinyl,

- benzimidazolyl, benzofuranyl, benzothiofuranyl,
benzothiophenyl, benzoxazolyl, benzthiazolyl,
benztriazolyl, benztetrazolyl, benzisoxazolyl,
benzisothiazolyl, benzimidazalonyl, carbazolyl,
5 4aH-carbazolyl, --carbolinyl, chromanyl, chromenyl,
cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl,
dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl,
imidazolidinyl, imidazolyl, imidazolyl, 1H-indazolyl,
indolenyl, indolinyl, indoliziny, indolyl,
10 isobenzofuranyl, isochromanyl, isoindazolyl,
isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl,
isoxazolyl, morpholinyl, naphthyridinyl,
octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl,
1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl,
15 oxazolidinyl., oxazolyl, oxazolidinylperimidinyl,
phenanthridinyl, phenanthrolinyl, phenarsazinyl,
phenazinyl, phenothiazinyl, phenoxathiinyl, phenoxazinyl,
phthalazinyl, piperazinyl, piperidinyl, pteridinyl,
piperidonyl, 4-piperidonyl, pteridinyl, purinyl, pyranyl,
20 pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl,
pyridazinyl, pyridooxazole, pyridoimidazole,
pyridothiazole, pyridinyl, pyridyl, pyrimidinyl,
pyrrolidinyl, pyrrolinyl, pyrrolyl, quinazolinyl,
quinolinyl, 4H-quinoliziny, quinoxaliny, quinuclidinyl,
25 carbolinyl, tetrahydrofuranyl, tetrahydroisoquinolinyl,
tetrahydroquinolinyl, 6H-1,2,5-thiadiazinyl,
1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl,
1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl,
thiazolyl, thienyl, thienothiazolyl, thienooxazolyl,
30 thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl,
1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl,
xanthylenyl. Preferred heterocycles include, but are not
limited to, pyridinyl, furanyl, thienyl, pyrrolyl,
pyrazolyl, imidazolyl, indolyl, benzimidazolyl,
35 1H-indazolyl, oxazolidinyl, benzotriazolyl,
benzisoxazolyl, oxindolyl, benzoxazolinyl, or isatinoyl.
Also included are fused ring and spiro compounds
containing, for example, the above heterocycles.

As used herein, the term "alkaryl" means an aryl group bearing an alkyl group of 1-10 carbon atoms; the term "aralkyl" means an alkyl group of 1-10 carbon atoms bearing an aryl group; the term "arylalkaryl" means an
5 aryl group bearing an alkyl group of 1-10 carbon atoms bearing an aryl group; and the term "heterocycloalkyl" means an alkyl group of 1-10 carbon atoms bearing a heterocycle.

A "polyalkylene glycol" is a polyethylene glycol,
10 polypropylene glycol or polybutylene glycol having a molecular weight of less than about 5000, terminating in either a hydroxy or alkyl ether moiety.

A "carbohydrate" is a polyhydroxy aldehyde, ketone, alcohol or acid, or derivatives thereof, including
15 polymers thereof having polymeric linkages of the acetal type.

A "cyclodextrin" is a cyclic oligosaccharide. Examples of cyclodextrins include, but are not limited to, α -cyclodextrin, hydroxyethyl- α -cyclodextrin,
20 hydroxypropyl- α -cyclodextrin, β -cyclodextrin, hydroxypropyl- β -cyclodextrin, carboxymethyl- β -cyclodextrin, dihydroxypropyl- β -cyclodextrin, hydroxyethyl- β -cyclodextrin, 2,6
25 di-O-methyl- β -cyclodextrin, sulfated- β -cyclodextrin, γ -cyclodextrin, hydroxypropyl- γ -cyclodextrin, dihydroxypropyl- γ -cyclodextrin, hydroxyethyl- γ -cyclodextrin, and sulfated γ -cyclodextrin.

As used herein, the term "polycarboxyalkyl" means an
30 alkyl group having between two and about 100 carbon atoms and a plurality of carboxyl substituents; and the term "polyazaalkyl" means a linear or branched alkyl group having between two and about 100 carbon atoms, interrupted by or substituted with a plurality of amine
35 groups.

A "reducing agent" is a compound that reacts with a radionuclide, which is typically obtained as a relatively unreactive, high oxidation state compound, to lower its oxidation state by transferring electron(s) to the
5 radionuclide, thereby making it more reactive. Reducing agents useful in the preparation of radiopharmaceuticals and in diagnostic kits useful for the preparation of said radiopharmaceuticals include but are not limited to stannous chloride, stannous fluoride, formamidine
10 sulfinic acid, ascorbic acid, cysteine, phosphines, and cuprous or ferrous salts. Other reducing agents are described in Brodack et. al., PCT Application 94/22496, which is incorporated herein by reference.

A "transfer ligand" is a ligand that forms an
15 intermediate complex with a metal ion that is stable enough to prevent unwanted side-reactions but labile enough to be converted to a metallopharmaceutical. The formation of the intermediate complex is kinetically favored while the formation of the metallopharmaceutical
20 is thermodynamically favored. Transfer ligands useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the preparation of diagnostic radiopharmaceuticals include but are not limited to gluconate, glucoheptonate, mannitol, glucarate,
25 N,N,N',N'-ethylenediaminetetraacetic acid, pyrophosphate and methylenediphosphonate. In general, transfer ligands are comprised of oxygen or nitrogen donor atoms.

The term "donor atom" refers to the atom directly attached to a metal by a chemical bond.

30 "Ancillary" or "co-ligands" are ligands that are incorporated into a radiopharmaceutical during its synthesis. They serve to complete the coordination sphere of the radionuclide together with the chelator or radionuclide bonding unit of the reagent. For
35 radiopharmaceuticals comprised of a binary ligand system, the radionuclide coordination sphere is composed of one or more chelators or bonding units from one or more reagents and one or more ancillary or co-ligands, provided that there are a total of two types of ligands,

chelators or bonding units. For example, a radiopharmaceutical comprised of one chelator or bonding unit from one reagent and two of the same ancillary or co-ligands and a radiopharmaceutical comprised of two
5 chelators or bonding units from one or two reagents and one ancillary or co-ligand are both considered to be comprised of binary ligand systems. For radiopharmaceuticals comprised of a ternary ligand system, the radionuclide coordination sphere is composed
10 of one or more chelators or bonding units from one or more reagents and one or more of two different types of ancillary or co-ligands, provided that there are a total of three types of ligands, chelators or bonding units. For example, a radiopharmaceutical comprised of one
15 chelator or bonding unit from one reagent and two different ancillary or co-ligands is considered to be comprised of a ternary ligand system.

Ancillary or co-ligands useful in the preparation of radiopharmaceuticals and in diagnostic kits useful for
20 the preparation of said radiopharmaceuticals are comprised of one or more oxygen, nitrogen, carbon, sulfur, phosphorus, arsenic, selenium, and tellurium donor atoms. A ligand can be a transfer ligand in the synthesis of a radiopharmaceutical and also serve as an
25 ancillary or co-ligand in another radiopharmaceutical. Whether a ligand is termed a transfer or ancillary or co-ligand depends on whether the ligand remains in the radionuclide coordination sphere in the radiopharmaceutical, which is determined by the
30 coordination chemistry of the radionuclide and the chelator or bonding unit of the reagent or reagents.

A "chelator" or "bonding unit" is the moiety or group on a reagent that binds to a metal ion through the formation of chemical bonds with one or more donor atoms.
35 The term "binding site" means the site in vivo or in vitro that binds a biologically active molecule.

A "diagnostic kit" or "kit" comprises a collection of components, termed the formulation, in one or more vials which are used by the practicing end user in a

clinical or pharmacy setting to synthesize diagnostic radiopharmaceuticals. The kit provides all the requisite components to synthesize and use the diagnostic radiopharmaceutical except those that are commonly
5 available to the practicing end user, such as water or saline for injection, a solution of the radionuclide, equipment for heating the kit during the synthesis of the radiopharmaceutical, if required, equipment necessary for administering the radiopharmaceutical to the patient such
10 as syringes and shielding, and imaging equipment.

Therapeutic radiopharmaceuticals, X-ray contrast agent pharmaceuticals, ultrasound contrast agent pharmaceuticals and metallopharmaceuticals for magnetic resonance imaging contrast are provided to the end user
15 in their final form in a formulation contained typically in one vial, as either a lyophilized solid or an aqueous solution. The end user reconstitutes the lyophilized with water or saline and withdraws the patient dose or just withdraws the dose from the aqueous solution
20 formulation as provided.

A "lyophilization aid" is a component that has favorable physical properties for lyophilization, such as the glass transition temperature, and is added to the formulation to improve the physical properties of the
25 combination of all the components of the formulation for lyophilization.

A "stabilization aid" is a component that is added to the metallopharmaceutical or to the diagnostic kit either to stabilize the metallopharmaceutical or to
30 prolong the shelf-life of the kit before it must be used. Stabilization aids can be antioxidants, reducing agents or radical scavengers and can provide improved stability by reacting preferentially with species that degrade other components or the metallopharmaceutical.

35 A "solubilization aid" is a component that improves the solubility of one or more other components in the medium required for the formulation.

A "bacteriostat" is a component that inhibits the growth of bacteria in a formulation either during its

storage before use of after a diagnostic kit is used to synthesize a radiopharmaceutical.

The following abbreviations are used herein:

5	AcM	acetamidomethyl
	b-Ala, beta-Ala	
	or bAla	3-aminopropionic acid
	ATA	2-aminothiazole-5-acetic acid or 2-aminothiazole-5-acetyl group
10	Boc	t-butyloxycarbonyl
	CBZ, Cbz or Z	Carbobenzyloxy
	Cit	citrulline
	Dap	2,3-diaminopropionic acid
	DCC	dicyclohexylcarbodiimide
15	DIEA	diisopropylethylamine
	DMAP	4-dimethylaminopyridine
	EOE	ethoxyethyl
	HBTU	2-(1H-Benzotriazol-1-yl)-1,1,3,3-tetramethyluronium
20	hexafluorophosphate	
	hynic	boc-hydrazinonicotinyl group or 2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid,
25	NMeArg or MeArga-N-methyl arginine	
	NMeAsp	a-N-methyl aspartic acid
	NMM	N-methylmorpholine
	OcHex	O-cyclohexyl
	OBzl	O-benzyl
30	oSu	O-succinimidyl
	TBTU	2-(1H-Benzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate
	THF	tetrahydrofuran
	THP	tetrahydropyranyl
35	Tos	tosyl
	Tr	trityl

The following conventional three-letter amino acid abbreviations are used herein; the conventional one-letter amino acid abbreviations are NOT used herein:

5	Ala	=	alanine
	Arg	=	arginine
	Asn	=	asparagine
	Asp	=	aspartic acid
	Cys	=	cysteine
10	Gln	=	glutamine
	Glu	=	glutamic acid
	Gly	=	glycine
	His	=	histidine
	Ile	=	isoleucine
15	Leu	=	leucine
	Lys	=	lysine
	Met	=	methionine
	Nle	=	norleucine
	Orn	=	ornithine
20	Phe	=	phenylalanine
	Phg	=	phenylglycine
	Pro	=	proline
	Sar	=	sarcosine
	Ser	=	serine
25	Thr	=	threonine
	Trp	=	tryptophan
	Tyr	=	tyrosine
	Val	=	valine

30 As used herein, the term "bubbles", as used herein, refers to vesicles which are generally characterized by the presence of one or more membranes or walls surrounding an internal void that is filled with a gas or precursor thereto. Exemplary bubbles include, for
35 example, liposomes, micelles and the like.

As used herein, the term "lipid" refers to a synthetic or naturally-occurring amphipathic compound which comprises a hydrophilic component and a hydrophobic component. Lipids include, for example, fatty acids,

neutral fats, phosphatides, glycolipids, aliphatic alcohols and waxes, terpenes and steroids.

As used herein, the term "lipid composition" refers to a composition which comprises a lipid compound.

- 5 Exemplary lipid compositions include suspensions, emulsions and vesicular compositions.

As used herein, the term "lipid formulation" refers to a composition which comprises a lipid compound and a bioactive agent.

- 10 As used herein, the term "vesicle" refers to a spherical entity which is characterized by the presence of an internal void. Preferred vesicles are formulated from lipids, including the various lipids described herein. In any given vesicle, the lipids may be in the
15 form of a monolayer or bilayer, and the mono- or bilayer lipids may be used to form one of more mono- or bilayers. In the case of more than one mono- or bilayer, the mono- or bilayers are generally concentric. The lipid vesicles described herein include such entities commonly referred
20 to as liposomes, micelles, bubbles, microbubbles, microspheres and the like. Thus, the lipids may be used to form a unilamellar vesicle (comprised of one monolayer or bilayer), an oligolamellar vesicle (comprised of about two or about three monolayers or bilayers) or a
25 multilamellar vesicle (comprised of more than about three monolayers or bilayers). The internal void of the vesicles may be filled with a liquid, including, for example, an aqueous liquid, a gas, a gaseous precursor, and/or a solid or solute material, including, for
30 example, a bioactive agent, as desired.

As used herein, the term "vesicular composition" refers to a composition which is formulate from lipids and which comprises vesicles.

- 35 As used herein, the term "vesicle formulation" refers to a composition which comprises vesicles and a bioactive agent.

As used herein, the term "lipsomes" refers to a generally spherical cluster or aggregate of amphipathic compounds, including lipid compounds, typically in the

form of one or more concentric layers, for example, bilayers. They may also be referred to herein as lipid vesicles.

Angiogenesis is the process of formation of new
5 capillary blood vessels from existing vasculature. It is an important component of a variety of physiological processes including ovulation, embryonic development, wound repair, and collateral vascular generation in the myocardium. It is also central to a number of
10 pathological conditions such as tumor growth and metastasis, diabetic retinopathy, and macular degeneration. The process begins with the activation of existing vascular endothelial cells in response to a variety of cytokines and growth factors. The activated
15 endothelial cells secrete enzymes that degrade the basement membrane of the vessels. The endothelial cells then proliferate and migrate into the extracellular matrix first forming tubules and subsequently new blood vessels.

20 Under normal conditions, endothelial cell proliferation is a very slow process, but it increases for a short period of time during embryogenesis, ovulation and wound healing. This temporary increase in cell turnover is governed by a combination of a number of
25 growth stimulatory factors and growth suppressing factors. In pathological angiogenesis, this normal balance is disrupted resulting in continued increased endothelial cell proliferation. Some of the pro-angiogenic factors that have been identified include
30 basic fibroblast growth factor (bFGF), angiogenin, TGF-alpha, TGF-beta, and vascular endothelium growth factor (VEGF), while interferon-alpha, interferon-beta and thrombospondin are examples of angiogenesis suppressors.

Angiogenic factors interact with endothelial cell
35 surface receptors such as the receptor tyrosine kinases EGFR, FGFR, PDGFR, Flk-1/KDR, Flt-1, Tek, Tie, neuropilin-1, endoglin, endosialin, and Axl. The receptors Flk-1/KDR, neuropilin-1, and Flt-1 recognize VEGF and these interactions play key roles in VEGF-

induced angiogenesis. The Tie subfamily of receptor tyrosine kinases are also expressed prominently during blood vessel formation.

The proliferation and migration of endothelial cells
5 in the extracellular matrix is mediated by interaction
with a variety of cell adhesion molecules. Integrins are
a diverse family of heterodimeric cell surface receptors
by which endothelial cells attach to the extracellular
matrix, each other and other cells. Angiogenesis induced
10 by bFGF or TNF-alpha depend on the agency of the integrin
avb3, while angiogenesis induced by VEGF depends on the
integrin avb5 (Cheresh et. al., Science, 1995, 270, 1500-
2). Induction of expression of the integrins albl and
a2b1 on the endothelial cell surface is another important
15 mechanism by which VEGF promotes angiogenesis (Senger,
et. al., Proc. Natl. Acad. Sci USA, 1997, 94, 13612-7).

The pharmaceuticals of the present invention are
comprised of a non-peptide targeting moiety for the
vitronectin receptor that is expressed or upregulated in
20 angiogenic tumor vasculature.

The ultrasound contrast agents of the present
invention comprise a plurality of vitronectin receptor
targeting moieties attached to or incorporated into a
25 microbubble of a biocompatible gas, a liquid carrier, and
a surfactant microsphere, further comprising an optional
linking moiety, L_n , between the targeting moieties and
the microbubble. In this context, the term liquid
carrier means aqueous solution and the term surfactant
30 means any amphiphilic material which produces a reduction
in interfacial tension in a solution. A list of suitable
surfactants for forming surfactant microspheres is
disclosed in EP0727225A2, herein incorporated by
reference. The term surfactant microsphere includes
35 nanospheres, liposomes, vesicles and the like. The
biocompatible gas can be air, or a fluorocarbon, such as
a C₃-C₅ perfluoroalkane, which provides the difference in
echogenicity and thus the contrast in ultrasound imaging.

The gas is encapsulated or contained in the microsphere to which is attached the biodirecting group, optionally via a linking group. The attachment can be covalent, ionic or by van der Waals forces. Specific examples of such contrast agents include lipid encapsulated perfluorocarbons with a plurality of tumor neovasculature receptor binding peptides, polypeptides or peptidomimetics.

X-ray contrast agents of the present invention are comprised of one or more vitronectin receptor targeting moieties attached to one or more X-ray absorbing or "heavy" atoms of atomic number 20 or greater, further comprising an optional linking moiety, L_n , between the targeting moieties and the X-ray absorbing atoms. The frequently used heavy atom in X-ray contrast agents is iodine. Recently, X-ray contrast agents comprised of metal chelates (Wallace, R., U.S. 5,417,959) and polychelates comprised of a plurality of metal ions (Love, D., U.S. 5,679,810) have been disclosed. More recently, multinuclear cluster complexes have been disclosed as X-ray contrast agents (U.S. 5,804,161, PCT WO91/14460, and PCT WO 92/17215).

MRI contrast agents of the present invention are comprised of one or more vitronectin receptor targeting moieties attached to one or more paramagnetic metal ions, further comprising an optional linking moiety, L_n , between the targeting moieties and the paramagnetic metal ions. The paramagnetic metal ions are present in the form of metal complexes or metal oxide particles. U.S. 5,412,148, and 5,760,191, describe examples of chelators for paramagnetic metal ions for use in MRI contrast agents. U.S. 5,801,228, U.S. 5,567,411, and U.S. 5,281,704, describe examples of polychelants useful for complexing more than one paramagnetic metal ion for use in MRI contrast agents. U.S. 5,520,904, describes particulate compositions comprised of paramagnetic metal ions for use as MRI contrast agents.

The pharmaceuticals of the present invention have the formulae, $(Q)_d-L_n-(C_h-X)$, $(Q)_d-L_n-(C_h-X^1)_d$, $(Q)_d-L_n-(X^2)_d$, and $(Q)_d-L_n-(X^3)$, wherein Q represents a non-peptide that binds to a receptor expressed in angiogenic tumor vasculature, d is 1-10, L_n represents an optional linking group, C_h represents a metal chelator or bonding moiety, X represents a radioisotope, X^1 represents paramagnetic metal ion, X^2 represents a paramagnetic metal ion or heavy atom containing insoluble solid particle, d^* is 1-100, and X^3 represents a surfactant microsphere of an echogenic gas. The interaction of the non-peptide recognition sequences of the vitronectin receptor binding portion of the pharmaceuticals with the $\alpha v \beta 3$ receptor results in localization of the pharmaceuticals in angiogenic tumor vasculature, which express the $\alpha v \beta 3$ receptor.

The pharmaceuticals of the present invention can be synthesized by several approaches. One approach involves the synthesis of the targeting non-peptide moiety, Q, and direct attachment of one or more moieties, Q, to one or more metal chelators or bonding moieties, C_h , or to a paramagnetic metal ion or heavy atom containing solid particle, or to an echogenic gas microbubble. Another approach involves the attachment of one or more moieties, Q, to the linking group, L_n , which is then attached to one or more metal chelators or bonding moieties, C_h , or to a paramagnetic metal ion or heavy atom containing solid particle, or to an echogenic gas microbubble. Another approach involves the synthesis of a non-peptide, Q, bearing a fragment of the linking group, L_n , one or more of which are then attached to the remainder of the linking group and then to one or more metal chelators or bonding moieties, C_h , or to a paramagnetic metal ion or heavy atom containing solid particle, or to an echogenic gas microbubble.

The non-peptide vitronectin binding moieties, Q, optionally bearing a linking group, L_n , or a fragment of


the linking group, can be synthesized using standard synthetic methods known to those skilled in the art. Preferred methods include but are not limited to those methods described below.

- 5 The attachment of linking groups, L_n , to the non-peptides, Q; chelators or bonding units, C_h , to the non-peptides, Q, or to the linking groups, L_n ; and non-peptides, bearing a fragment of the linking group to the remainder of the linking group, in combination forming
10 the moiety, $(Q)_d-L_n$, and then to the moiety C_h ; can all be performed by standard techniques. These include, but are not limited to, amidation, esterification, alkylation, and the formation of ureas or thioureas. Procedures for performing these attachments can be found in Brinkley,
15 M., *Bioconjugate Chemistry* 1992, 3(1), which is incorporated herein by reference.

- A number of methods can be used to attach the non-peptides, Q, to paramagnetic metal ion or heavy atom containing solid particles, X^2 , by one of skill in the
20 art of the surface modification of solid particles. In general, the targeting moiety Q or the combination $(Q)_d-L_n$ is attached to a coupling group that react with a constituent of the surface of the solid particle. The coupling groups can be any of a number of silanes which
25 react with surface hydroxyl groups on the solid particle surface, as described in co-pending United States Patent Application Serial No. 09/356,178, and can also include polyphosphonates, polycarboxylates, polyphosphates or mixtures thereof which couple with the surface of the
30 solid particles, as described in U.S. 5,520,904.

- A number of reaction schemes can be used to attach the non-peptides, Q, to the surfactant microsphere, X^3 . These are illustrated in following reaction schemes where S_f represents a surfactant moiety that forms the
35 surfactant microsphere.

Acylation Reaction:




Y is a leaving group or active ester

5 Disulfide Coupling:



Sulfonamide Coupling:

10



Reductive Amidation:

15



20 In these reaction schemes, the substituents S_f and Q can be reversed as well.

The linking group L_n can serve several roles. First it provides a spacing group between the metal chelator or bonding moiety, C_h , the paramagnetic metal ion or heavy atom containing solid particle, X^2 , and the surfactant
 25 microsphere, X^3 , and the one or more of the non-peptides, Q , so as to minimize the possibility that the moieties C_h-X , C_h-X^1 , X^2 , and X^3 , will interfere with the interaction of the recognition sequences of Q with angiogenic tumor vasculature receptors. The necessity of
 30 incorporating a linking group in a reagent is dependent on the identity of Q , C_h-X , C_h-X^1 , X^2 , and X^3 . If C_h-X , C_h-X^1 , X^2 , and X^3 , cannot be attached to Q without substantially diminishing its affinity for the receptors, then a linking group is used. A linking group also
 35 provides a means of independently attaching multiple non-peptides, Q , to one group that is attached to C_h-X , C_h-X^1 , X^2 , or X^3 .

The linking group also provides a means of incorporating a pharmacokinetic modifier into the pharmaceuticals of the present invention. The pharmacokinetic modifier serves to direct the

5 biodistribution of the injected pharmaceutical other than by the interaction of the targeting moieties, Q, with the vitronectin receptors expressed in the tumor neovasculature. A wide variety of functional groups can serve as pharmacokinetic modifiers, including, but not

10 limited to, carbohydrates, polyalkylene glycols, peptides or other polyamino acids, and cyclodextrins. The modifiers can be used to enhance or decrease hydrophilicity and to enhance or decrease the rate of blood clearance. The modifiers can also be used to

15 direct the route of elimination of the pharmaceuticals. Preferred pharmacokinetic modifiers are those that result in moderate to fast blood clearance and enhanced renal excretion.

The metal chelator or bonding moiety, Ch, is

20 selected to form stable complexes with the metal ion chosen for the particular application. Chelators or bonding moieties for diagnostic radiopharmaceuticals are selected to form stable complexes with the radioisotopes that have imageable gamma ray or positron emissions, such

25 as ^{99m}Tc , ^{95}Tc , ^{111}In , ^{62}Cu , ^{60}Cu , ^{64}Cu , ^{67}Ga , ^{68}Ga , ^{86}Y .

Chelators for technetium, copper and gallium isotopes are selected from diaminedithiols, monoamine-monoamidedithiols, triamide-monothiol, monoamine-diamide-monothiol, diaminedioximes, and

30 hydrazines. The chelators are generally tetradentate with donor atoms selected from nitrogen, oxygen and sulfur. Preferred reagents are comprised of chelators having amine nitrogen and thiol sulfur donor atoms and hydrazine bonding units. The thiol sulfur atoms and the

35 hydrazines may bear a protecting group which can be displaced either prior to using the reagent to synthesize a radiopharmaceutical or preferably in situ during the synthesis of the radiopharmaceutical.

Exemplary thiol protecting groups include those listed in Greene and Wuts, "Protective Groups in Organic Synthesis" John Wiley & Sons, New York (1991), the disclosure of which is hereby incorporated by reference.

- 5 Any thiol protecting group known in the art can be used. Examples of thiol protecting groups include, but are not limited to, the following: acetamidomethyl, benzamidomethyl, 1-ethoxyethyl, benzoyl, and triphenylmethyl.

- 10 Exemplary protecting groups for hydrazine bonding units are hydrazones which can be aldehyde or ketone hydrazones having substituents selected from hydrogen, alkyl, aryl and heterocycle. Particularly preferred hydrazones are described in co-pending U.S.S.N.
15 08/476,296 the disclosure of which is herein incorporated by reference in its entirety.

- The hydrazine bonding unit when bound to a metal radionuclide is termed a hydrazido, or diazenido group and serves as the point of attachment of the radionuclide
20 to the remainder of the radiopharmaceutical. A diazenido group can be either terminal (only one atom of the group is bound to the radionuclide) or chelating. In order to have a chelating diazenido group at least one other atom of the group must also be bound to the radionuclide. The
25 atoms bound to the metal are termed donor atoms.

- Chelators for ^{111}In and ^{86}Y are selected from cyclic and acyclic polyaminocarboxylates such as DTPA, DOTA, DO3A, 2-benzyl-DOTA, alpha-(2-phenethyl)1,4,7,10-tetraazacyclododecane-1-acetic-4,7,10-
30 tris(methylacetic)acid, 2-benzyl-cyclohexyldiethylenetriaminepentaacetic acid, 2-benzyl-6-methyl-DTPA, and 6,6"-bis[N,N,N",N"-tetra(carboxymethyl)aminomethyl)-4'-(3-amino-4-methoxyphenyl)-2,2':6',2"-terpyridine. Procedures for
35 synthesizing these chelators that are not commercially available can be found in Brechbiel, M. and Gansow, O., *J. Chem. Soc. Perkin Trans. 1992, 1, 1175*; Brechbiel, M. and Gansow, O., *Bioconjugate Chem. 1991, 2, 187*; Deshpande, S., et. al., *J. Nucl. Med. 1990, 31, 473*;

Kruper, J., U.S. Patent 5,064,956, and Toner, J., U.S. Patent 4,859,777, the disclosures of which are hereby incorporated by reference in their entirety.

5 The coordination sphere of metal ion includes all
the ligands or groups bound to the metal. For a
transition metal radionuclide to be stable it typically
has a coordination number (number of donor atoms)
comprised of an integer greater than or equal to 4 and
less than or equal to 8; that is there are 4 to 8 atoms
10 bound to the metal and it is said to have a complete
coordination sphere. The requisite coordination number
for a stable radionuclide complex is determined by the
identity of the radionuclide, its oxidation state, and
the type of donor atoms. If the chelator or bonding unit
15 does not provide all of the atoms necessary to stabilize
the metal radionuclide by completing its coordination
sphere, the coordination sphere is completed by donor
atoms from other ligands, termed ancillary or co-ligands,
which can also be either terminal or chelating.

20 A large number of ligands can serve as ancillary or
co-ligands, the choice of which is determined by a
variety of considerations such as the ease of synthesis
of the radiopharmaceutical, the chemical and physical
properties of the ancillary ligand, the rate of
25 formation, the yield, and the number of isomeric forms of
the resulting radiopharmaceuticals, the ability to
administer said ancillary or co-ligand to a patient
without adverse physiological consequences to said
patient, and the compatibility of the ligand in a
30 lyophilized kit formulation. The charge and
lipophilicity of the ancillary ligand will effect the
charge and lipophilicity of the radiopharmaceuticals.
For example, the use of 4,5-dihydroxy-1,3-benzene
disulfonate results in radiopharmaceuticals with an
35 additional two anionic groups because the sulfonate
groups will be anionic under physiological conditions.
The use of N-alkyl substituted 3,4-hydroxypyridinones
results in radiopharmaceuticals with varying degrees of

lipophilicity depending on the size of the alkyl substituents.

Preferred technetium radiopharmaceuticals of the present invention are comprised of a hydrazido or diazenido bonding unit and an ancillary ligand, A_{L1} , or a bonding unit and two types of ancillary A_{L1} and A_{L2} , or a tetradentate chelator comprised of two nitrogen and two sulfur atoms. Ancillary ligands A_{L1} are comprised of two or more hard donor atoms such as oxygen and amine nitrogen (sp^3 hybridized). The donor atoms occupy at least two of the sites in the coordination sphere of the radionuclide metal; the ancillary ligand A_{L1} serves as one of the three ligands in the ternary ligand system. Examples of ancillary ligands A_{L1} include but are not limited to dioxygen ligands and functionalized aminocarboxylates. A large number of such ligands are available from commercial sources.

Ancillary dioxygen ligands include ligands that coordinate to the metal ion through at least two oxygen donor atoms. Examples include but are not limited to: glucoheptonate, gluconate, 2-hydroxyisobutyrate, lactate, tartrate, mannitol, glucarate, maltol, Kojic acid, 2,2-bis(hydroxymethyl)propionic acid, 4,5-dihydroxy-1,3-benzene disulfonate, or substituted or unsubstituted 1,2 or 3,4 hydroxypyridinones. (The names for the ligands in these examples refer to either the protonated or non-protonated forms of the ligands.)

Functionalized aminocarboxylates include ligands that have a combination of amine nitrogen and oxygen donor atoms. Examples include but are not limited to: iminodiacetic acid, 2,3-diaminopropionic acid, nitrilotriacetic acid, N,N'-ethylenediamine diacetic acid, N,N,N'-ethylenediamine triacetic acid, hydroxyethylethylenediamine triacetic acid, and N,N'-ethylenediamine bis-hydroxyphenylglycine. (The names for the ligands in these examples refer to either the protonated or non-protonated forms of the ligands.)

A series of functionalized aminocarboxylates are disclosed by Bridger et. al. in U.S. Patent 5,350,837, herein incorporated by reference, that result in improved rates of formation of technetium labeled hydrazino
5 modified proteins. We have determined that certain of these aminocarboxylates result in improved yields of the radiopharmaceuticals of the present invention. The preferred ancillary ligands A_{L1} functionalized aminocarboxylates that are derivatives of glycine; the
10 most preferred is tricine (tris(hydroxymethyl)methylglycine).

The most preferred technetium radiopharmaceuticals of the present invention are comprised of a hydrazido or diazenido bonding unit and two types of ancillary
15 designated A_{L1} and A_{L2} , or a diaminedithiol chelator. The second type of ancillary ligands A_{L2} are comprised of one or more soft donor atoms selected from the group: phosphine phosphorus, arsine arsenic, imine nitrogen (sp^2 hybridized), sulfur (sp^2 hybridized) and carbon (sp
20 hybridized); atoms which have p-acid character. Ligands A_{L2} can be monodentate, bidentate or tridentate, the denticity is defined by the number of donor atoms in the ligand. One of the two donor atoms in a bidentate ligand and one of the three donor atoms in a tridentate ligand
25 must be a soft donor atom. We have disclosed in co-pending U.S.S.N. 08/415,908, and U.S.S.N. 60/013360 and 08/646,886, the disclosures of which are herein incorporated by reference in their entirety, that radiopharmaceuticals comprised of one or more ancillary
30 or co-ligands A_{L2} are more stable compared to radiopharmaceuticals that are not comprised of one or more ancillary ligands, A_{L2} ; that is, they have a minimal number of isomeric forms, the relative ratios of which do not change significantly with time, and that remain
35 substantially intact upon dilution.

The ligands A_{L2} that are comprised of phosphine or arsine donor atoms are trisubstituted phosphines, trisubstituted arsines, tetrasubstituted diphosphines and

- tetrasubstituted diarsines. The ligands A_{L2} that are comprised of imine nitrogen are unsaturated or aromatic nitrogen-containing, 5 or 6-membered heterocycles. The ligands that are comprised of sulfur (sp^2 hybridized) donor atoms are thiocarbonyls, comprised of the moiety $C=S$. The ligands comprised of carbon (sp hybridized) donor atoms are isonitriles, comprised of the moiety CNR , where R is an organic radical. A large number of such ligands are available from commercial sources.
- 5 Isonitriles can be synthesized as described in European Patent 0107734 and in U.S. Patent 4,988,827, herein incorporated by reference.

Preferred ancillary ligands A_{L2} are trisubstituted phosphines and unsaturated or aromatic 5 or 6 membered heterocycles. The most preferred ancillary ligands A_{L2} are trisubstituted phosphines and unsaturated 5 membered heterocycles.

The ancillary ligands A_{L2} may be substituted with alkyl, aryl, alkoxy, heterocycle, aralkyl, alkaryl and arylalkaryl groups and may or may not bear functional groups comprised of heteroatoms such as oxygen, nitrogen, phosphorus or sulfur. Examples of such functional groups include but are not limited to: hydroxyl, carboxyl, carboxamide, nitro, ether, ketone, amino, ammonium, sulfonate, sulfonamide, phosphonate, and phosphonamide.

20 The functional groups may be chosen to alter the lipophilicity and water solubility of the ligands which may affect the biological properties of the radiopharmaceuticals, such as altering the distribution into non-target tissues, cells or fluids, and the mechanism and rate of elimination from the body.

Chelators or bonding moieties for therapeutic radiopharmaceuticals are selected to form stable complexes with the radioisotopes that have alpha particle, beta particle, Auger or Coster-Kronig electron emissions, such as ^{186}Re , ^{188}Re , ^{153}Sm , ^{166}Ho , ^{177}Lu , ^{149}Pm , ^{90}Y , ^{212}Bi , ^{103}Pd , ^{109}Pd , ^{159}Gd , ^{140}La , ^{198}Au , ^{199}Au , ^{169}Yb , ^{175}Yb , ^{165}Dy , ^{166}Dy , ^{67}Cu , ^{105}Rh , ^{111}Ag , and ^{192}Ir .

- Chelators for rhenium, copper, palladium, platinum, iridium, rhodium, silver and gold isotopes are selected from diaminedithiols, monoamine-monoamidedithiols, triamide-monothiol, monoamine-diamide-monothiol, 5 diaminedioximes, and hydrazines. Chelators for yttrium, bismuth, and the lanthanide isotopes are selected from cyclic and acyclic polyaminocarboxylates such as DTPA, DOTA, DO3A, 2-benzyl-DOTA, alpha-(2-phenethyl)1,4,7,10-tetraazacyclododecane-1-acetic-4,7,10- 10 tris(methylacetic)acid, 2-benzyl-cyclohexyldiethylenetriaminepentaacetic acid, 2-benzyl-6-methyl-DTPA, and 6,6"-bis[N,N,N",N"-tetra(carboxymethyl)aminomethyl)-4'-(3-amino-4-methoxyphenyl)-2,2':6',2"-terpyridine.
- 15 Chelators for magnetic resonance imaging contrast agents are selected to form stable complexes with paramagnetic metal ions, such as Gd(III), Dy(III), Fe(III), and Mn(II), are selected from cyclic and acyclic polyaminocarboxylates such as DTPA, DOTA, DO3A, 20 2-benzyl-DOTA, alpha-(2-phenethyl)1,4,7,10-tetraazacyclododecane-1-acetic-4,7,10-tris(methylacetic)acid, 2-benzyl-cyclohexyldiethylenetriaminepentaacetic acid, 2-benzyl-6-methyl-DTPA, and 6,6"-bis[N,N,N",N"- 25 tetra(carboxymethyl)aminomethyl)-4'-(3-amino-4-methoxyphenyl)-2,2':6',2"-terpyridine.

The technetium and rhenium radiopharmaceuticals of the present invention comprised of a hydrazido or diazenido bonding unit can be easily prepared by admixing 30 a salt of a radionuclide, a reagent of the present invention, an ancillary ligand A_{L1} , an ancillary ligand A_{L2} , and a reducing agent, in an aqueous solution at temperatures from 0 to 100 °C. The technetium and rhenium radiopharmaceuticals of the present invention 35 comprised of a tetradentate chelator having two nitrogen and two sulfur atoms can be easily prepared by admixing a salt of a radionuclide, a reagent of the present

invention, and a reducing agent, in an aqueous solution at temperatures from 0 to 100 °C.

When the bonding unit in the reagent of the present invention is present as a hydrazone group, then it must first be converted to a hydrazine, which may or may not be protonated, prior to complexation with the metal radionuclide. The conversion of the hydrazone group to the hydrazine can occur either prior to reaction with the radionuclide, in which case the radionuclide and the ancillary or co-ligand or ligands are combined not with the reagent but with a hydrolyzed form of the reagent bearing the chelator or bonding unit, or in the presence of the radionuclide in which case the reagent itself is combined with the radionuclide and the ancillary or co-ligand or ligands. In the latter case, the pH of the reaction mixture must be neutral or acidic.

Alternatively, the radiopharmaceuticals of the present invention comprised of a hydrazido or diazenido bonding unit can be prepared by first admixing a salt of a radionuclide, an ancillary ligand A_{L1} , and a reducing agent in an aqueous solution at temperatures from 0 to 100 °C to form an intermediate radionuclide complex with the ancillary ligand A_{L1} then adding a reagent of the present invention and an ancillary ligand A_{L2} and reacting further at temperatures from 0 to 100 °C.

Alternatively, the radiopharmaceuticals of the present invention comprised of a hydrazido or diazenido bonding unit can be prepared by first admixing a salt of a radionuclide, an ancillary ligand A_{L1} , a reagent of the present invention, and a reducing agent in an aqueous solution at temperatures from 0 to 100 °C to form an intermediate radionuclide complex, and then adding an ancillary ligand A_{L2} and reacting further at temperatures from 0 to 100 °C.

The technetium and rhenium radionuclides are preferably in the chemical form of pertechnetate or perrhenate and a pharmaceutically acceptable cation. The pertechnetate salt form is preferably sodium

pertechnetate such as obtained from commercial Tc-99m generators. The amount of pertechnetate used to prepare the radiopharmaceuticals of the present invention can range from 0.1 mCi to 1 Ci, or more preferably from 1 to 5 200 mCi.

The amount of the reagent of the present invention used to prepare the technetium and rhenium radiopharmaceuticals of the present invention can range from 0.01 μ g to 10 mg, or more preferably from 0.5 μ g to 10 200 μ g. The amount used will be dictated by the amounts of the other reactants and the identity of the radiopharmaceuticals of the present invention to be prepared.

The amounts of the ancillary ligands A_{L1} used can 15 range from 0.1 mg to 1 g, or more preferably from 1 mg to 100 mg. The exact amount for a particular radiopharmaceutical is a function of identity of the radiopharmaceuticals of the present invention to be prepared, the procedure used and the amounts and 20 identities of the other reactants. Too large an amount of A_{L1} will result in the formation of by-products comprised of technetium labeled A_{L1} without a biologically active molecule or by-products comprised of technetium labeled biologically active molecules with the ancillary 25 ligand A_{L1} but without the ancillary ligand A_{L2} . Too small an amount of A_{L1} will result in other by-products such as technetium labeled biologically active molecules with the ancillary ligand A_{L2} but without the ancillary ligand A_{L1} , or reduced hydrolyzed technetium, or 30 technetium colloid.

The amounts of the ancillary ligands A_{L2} used can range from 0.001 mg to 1 g, or more preferably from 0.01 mg to 10 mg. The exact amount for a particular radiopharmaceutical is a function of the identity of the 35 radiopharmaceuticals of the present invention to be prepared, the procedure used and the amounts and identities of the other reactants. Too large an amount of A_{L2} will result in the formation of by-products

comprised of technetium labeled A_{L2} without a biologically active molecule or by-products comprised of technetium labeled biologically active molecules with the ancillary ligand A_{L2} but without the ancillary ligand A_{L1} . If the
5 reagent bears one or more substituents that are comprised of a soft donor atom, as defined above, at least a ten-fold molar excess of the ancillary ligand A_{L2} to the reagent of formula 2 is required to prevent the substituent from interfering with the coordination of the
10 ancillary ligand A_{L2} to the metal radionuclide.

Suitable reducing agents for the synthesis of the radiopharmaceuticals of the present invention include stannous salts, dithionite or bisulfite salts, borohydride salts, and formamidinesulfinic acid, wherein
15 the salts are of any pharmaceutically acceptable form. The preferred reducing agent is a stannous salt. The amount of a reducing agent used can range from 0.001 mg to 10 mg, or more preferably from 0.005 mg to 1 mg.

The specific structure of a radiopharmaceutical of
20 the present invention comprised of a hydrazido or diazenido bonding unit will depend on the identity of the reagent of the present invention used, the identity of any ancillary ligand A_{L1} , the identity of any ancillary ligand A_{L2} , and the identity of the radionuclide.

25 Radiopharmaceuticals comprised of a hydrazido or diazenido bonding unit synthesized using concentrations of reagents of $<100 \mu\text{g/mL}$, will be comprised of one hydrazido or diazenido group. Those synthesized using $>1 \text{ mg/mL}$ concentrations will be comprised of two hydrazido
30 or diazenido groups from two reagent molecules. For most applications, only a limited amount of the biologically active molecule can be injected and not result in undesired side-effects, such as chemical toxicity, interference with a biological process or an altered
35 biodistribution of the radiopharmaceutical. Therefore, the radiopharmaceuticals which require higher concentrations of the reagents comprised in part of the

biologically active molecule, will have to be diluted or purified after synthesis to avoid such side-effects.

The identities and amounts used of the ancillary ligands A_{L1} and A_{L2} will determine the values of the variables y and z . The values of y and z can independently be an integer from 1 to 2. In combination, the values of y and z will result in a technetium coordination sphere that is made up of at least five and no more than seven donor atoms. For monodentate ancillary ligands A_{L2} , z can be an integer from 1 to 2; for bidentate or tridentate ancillary ligands A_{L2} , z is 1. The preferred combination for monodentate ligands is y equal to 1 or 2 and z equal to 1. The preferred combination for bidentate or tridentate ligands is y equal to 1 and z equal to 1.

The indium, copper, gallium, silver, palladium, rhodium, gold, platinum, bismuth, yttrium and lanthanide radiopharmaceuticals of the present invention can be easily prepared by admixing a salt of a radionuclide and a reagent of the present invention, in an aqueous solution at temperatures from 0 to 100 °C. These radionuclides are typically obtained as a dilute aqueous solution in a mineral acid, such as hydrochloric, nitric or sulfuric acid. The radionuclides are combined with from one to about one thousand equivalents of the reagents of the present invention dissolved in aqueous solution. A buffer is typically used to maintain the pH of the reaction mixture between 3 and 10.

The gadolinium, dysprosium, iron and manganese metallopharmaceuticals of the present invention can be easily prepared by admixing a salt of the paramagnetic metal ion and a reagent of the present invention, in an aqueous solution at temperatures from 0 to 100 °C. These paramagnetic metal ions are typically obtained as a dilute aqueous solution in a mineral acid, such as hydrochloric, nitric or sulfuric acid. The paramagnetic metal ions are combined with from one to about one thousand equivalents of the reagents of the present

invention dissolved in aqueous solution. A buffer is typically used to maintain the pH of the reaction mixture between 3 and 10.

5 The total time of preparation will vary depending on the identity of the metal ion, the identities and amounts of the reactants and the procedure used for the preparation. The preparations may be complete, resulting in > 80% yield of the radiopharmaceutical, in 1 minute or may require more time. If higher purity
10 metallopharmaceuticals are needed or desired, the products can be purified by any of a number of techniques well known to those skilled in the art such as liquid chromatography, solid phase extraction, solvent extraction, dialysis or ultrafiltration.

15 Buffers useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the preparation of said radiopharmaceuticals include but are not limited to phosphate, citrate, sulfosalicylate, and acetate. A more complete list can be found in the
20 United States Pharmacopeia.

Lyophilization aids useful in the preparation of diagnostic kits useful for the preparation of radiopharmaceuticals include but are not limited to mannitol, lactose, sorbitol, dextran, Ficoll, and
25 polyvinylpyrrolidone (PVP).

Stabilization aids useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the preparation of radiopharmaceuticals include but are not limited to ascorbic acid, cysteine, monothioglycerol,
30 sodium bisulfite, sodium metabisulfite, gentisic acid, and inositol.

Solubilization aids useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the preparation of radiopharmaceuticals include but are
35 not limited to ethanol, glycerin, polyethylene glycol, propylene glycol, polyoxyethylene sorbitan monooleate, sorbitan monooleate, polysorbates, poly(oxyethylene)poly(oxypropylene)poly(oxyethylene)

block copolymers (Pluronic) and lecithin. Preferred solubilizing aids are polyethylene glycol, and Pluronic.

Bacteriostats useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the preparation of radiopharmaceuticals include but are not limited to benzyl alcohol, benzalkonium chloride, chlorbutanol, and methyl, propyl or butyl paraben.

A component in a diagnostic kit can also serve more than one function. A reducing agent can also serve as a stabilization aid, a buffer can also serve as a transfer ligand, a lyophilization aid can also serve as a transfer, ancillary or co-ligand and so forth.

The diagnostic radiopharmaceuticals are administered by intravenous injection, usually in saline solution, at a dose of 1 to 100 mCi per 70 kg body weight, or preferably at a dose of 5 to 50 mCi. Imaging is performed using known procedures.

The therapeutic radiopharmaceuticals are administered by intravenous injection, usually in saline solution, at a dose of 0.1 to 100 mCi per 70 kg body weight, or preferably at a dose of 0.5 to 5 mCi per 70 kg body weight.

The magnetic resonance imaging contrast agents of the present invention may be used in a similar manner as other MRI agents as described in U.S. Patent 5,155,215; U.S. Patent 5,087,440; Margerstadt et al., Magn. Reson. Med., 1986, 3, 808; Runge et al., Radiology, 1988, 166, 835; and Bousquet et al., Radiology, 1988, 166, 693. Generally, sterile aqueous solutions of the contrast agents are administered to a patient intravenously in dosages ranging from 0.01 to 1.0 mmoles per kg body weight.

For use as X-ray contrast agents, the compositions of the present invention should generally have a heavy atom concentration of 1 mM to 5 M, preferably 0.1 M to 2 M. Dosages, administered by intravenous injection, will typically range from 0.5 mmol/kg to 1.5 mmol/kg, preferably 0.8 mmol/kg to 1.2 mmol/kg. Imaging is

performed using known techniques, preferably X-ray computed tomography.

The ultrasound contrast agents of the present invention are administered by intravenous injection in an amount of 10 to 30 μ L of the echogenic gas per kg body weight or by infusion at a rate of approximately 3 μ L/kg/min. Imaging is performed using known techniques of sonography.

Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

EXAMPLES


Representative materials and methods that may be used in preparing the compounds of the invention are described further below.

1-methyl-4-oxo-7-(((1-(triphenylmethyl)imidazol-2-yl)amino)methyl)hydroquinoline-3-carboxylic acid, ethyl 7-bromo-4-oxohydroquinoline-3-carboxylate, 1-(triphenylmethyl)imidazole-2-ylamine, and methyl 3-amino-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)propanoate hydrochloride were prepared as described in PCT WO 98/23608. Boc-L-cysteic acid, Boc-L-cysteic acid N-hydroxyphenyl ester, and Boc-L-cysteic acid p-nitrophenyl ester were prepared as described in *Liebigs Ann. Chem.* 1979, 776-783. Benzotriazole-1-yloxy-tris-pyrrolidinophosphonium hexafluorophosphate (PyBOP) was purchased from Novabiochem.

(tert-butoxy)-N-(3-bromopropyl)formamide and 2-(2-aza-2-((5-((2,5-dioxopyrrolidinyl)carbonyl)(2-pyridyl))-amino)vinyl)benzenesulfonic acid were prepared as described in PCT WO 96/40637. All other chemicals and solvents (reagent grade) were used as supplied from the vendors cited without further purification. t-Butyloxycarbonyl (Boc) amino acids and other starting amino acids may be obtained commercially from Bachem Inc., Bachem Biosciences Inc. (Philadelphia, PA), Advanced ChemTech (Louisville, KY), Peninsula

Laboratories (Belmont, CA), or Sigma (St. Louis, MO). 2-(1H-Benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) and TBTU were purchased from Advanced ChemTech. N-methylmorpholine (NMM), *m*-cresol, 5 D-2-aminobutyric acid (Abu), trimethylacetylchloride, diisopropylethylamine (DIEA), 1,2,4-triazole, stannous chloride dihydrate, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC), triethylsilane (Et₃SiH) and tris(3-sulfonatophenyl)phosphine trisodium 10 salt (TPPTS) were purchased from Aldrich Chemical Company. Bis(3-sulfonatophenyl)phenylphosphine disodium salt (TPPDS) was prepared by the published procedure (Kuntz, E., U.S. Patent 4,248,802). (3-Sulfonatophenyl)diphenylphosphine monosodium salt 15 (TPPMS) was purchased from TCI America, Inc. Tricine was obtained from Research Organics, Inc. Technetium-99m-pertechnetate (^{99m}TcO₄⁻) was obtained from a DuPont Pharma ⁹⁹Mo/^{99m}Tc Technelite® generator. In-111-chloride (Indichlor®) was obtained from Amersham Medi-Physics, 20 Inc. Sm-153-chloride and Lutetium-177-chloride were obtained from the University of Missouri Research Reactor (MURR). Yttrium-90 chloride was obtained from the Pacific Northwest Research Laboratories. Dimethylformamide (DMF), ethyl acetate, chloroform (CHCl₃), methanol 25 (MeOH), pyridine and hydrochloric acid (HCl) were obtained from Baker. Acetonitrile, dichloromethane (DCM), acetic acid (HOAc), trifluoroacetic acid (TFA), ethyl ether, triethylamine, acetone, and magnesium sulfate were commercially obtained. Absolute ethanol was 30 obtained from Quantum Chemical Corporation.

Synthesis of Boc-Glu-(OTFP)-OTFP



To a solution of Boc-Glu-OH (28.9 g, 117 mmol) in DMF (500 mL) at room temperature, and under nitrogen, was added a solution of 2,3,5,6-tetrafluorophenol (48.2 g, 290 mmol) in DMF (50 mL). After stirring for 10 min. EDC (55.6 g, 290 mmol) was added and the reaction mixture was stirred for about 96 h. The volatiles were removed in vacuo and the residue was triturated in 0.1 N HCl (750 mL). To this mixture was added ethyl acetate (600 mL), the layers separated. The aqueous layer was extracted with ethyl acetate (3 x ~500 mL), and all the ethyl acetate fractions were combined, washed with water (300 mL) and brine (300 mL), dried (MgSO₄), and concentrated to give a tan solid (62 g). The tan solid was washed with acetonitrile to give the title compound (45.5 g, 73%) in purified form.

ESMS: Calculated for C₂₂H₁₇F₈NO₆, 543.09; found, 566.0 [M+Na]⁺¹.

20

Example 1


2-(((4-(4-(((3-(2-(2-(3-((6-((1-Aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)-propoxy)ethoxy)-ethoxy)propyl)amino)sulfonyl)phenyl)phenyl)sulfonyl)-amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic Acid
Trifluoroacetate Salt

30 Part A - N-(3-(2-(2-(3-aminopropoxy)ethoxy)ethoxy)-propyl)(phenylmethoxy)formamide

A solution of 4,7,10-trioxa-1,13-tridecanediamine (158 mL, 0.72 mol), TEA (16.7 mL, 0.12 mol), and MeOH (300 mL) in peroxide-free THF (1,000 mL) was placed in a 3 liter 3-neck flask fitted with a mechanical stirrer, a thermometer, and an addition funnel with nitrogen line. The addition funnel was charged with a solution of benzyl chloroformate (17.1 mL, 0.12 mol) in peroxide-free THF

(1,000 mL). The contents of the flask were cooled below 5 °C. The contents of the addition funnel were added to the flask with rapid stirring over 4 h while keeping the temperature below 5 °C. The solution was stirred an
 5 additional 30 min and concentrated to give a thick syrup. This syrup was taken up in saturated NaCl (1800 mL) and 10% Na₂CO₃ (200 mL) and extracted with ether (3 x 1,000 mL). The combined ether extracts were washed with saturated NaCl (500 mL), dried (MgSO₄), and concentrated
 10 to give a pale yellow oil (36.74 g). Flash chromatography on a 7 x 29 cm silica gel column (DCM/MeOH/TEA, 20/15/0.5) gave the title compound as a colorless syrup (19.14 g, 45%). ¹H NMR (CDCl₃): 7.33-7.25 (m, 5H), 5.59 (s, 1H), 5.06 (s, 2H), 3.62-3.45 (m,
 15 12H), 3.32-3.25 (m, 2H), 2.74 (t, J = 6.7 Hz, 2H), 1.75 (pentet, J = 6.0 Hz, 2H), 1.67 (pentet, J = 6.4 Hz, 2H), 1.33 (s, 2H); MS: m/e 355.4 [M+H]; High Resolution MS: Calcd for C₁₈H₃₁N₂O₅ [M+H]: 355.2233, Found: 355.2222.


20



Part B - Methyl 3-((tert-Butoxy)carbonylamino)-2-(((4-(4-
 (((3-(2-(2-(3-
 ((phenylmethoxy)carbonylamino)propoxy)ethoxy)-
 25 ethoxy)propyl)amino)sulfonyl)phenyl)phenyl)sulfonyl)-
 amino)propanoate

Biphenyl-4,4'-disulfonyl chloride (2.64 g, 7.5 mmol, freshly recrystallized from CHCl₃) and DCM (200 mL) were placed in a 500 mL 3-neck flask fitted with a
 30 thermometer, an addition funnel, and a nitrogen line. The addition funnel was charged with a solution of N-(3-(2-(2-(3-aminopropoxy)ethoxy)ethoxy)propyl)-
 (phenylmethoxy)formamide (1.77 g, 5.0 mmol) and DIEA (0.87 mL, 5.0 mmol) in DCM (40 mL). The contents of the
 35 flask were cooled below 5 °C. The contents of the addition funnel were added to the flask with rapid

stirring over 3 h while keeping the temperature of the flask below 5 °C. The addition funnel was charged with a solution of N-β-Boc-L-α,β,-diaminopropionic acid methyl ester hydrochloride (2.55 g, 10 mmol) and DIEA (3.8 mL, 22 mmol) in DCM (25 mL). This solution was added to the flask with stirring at 5 °C over 15 min, and stirred at ambient temperatures for an additional 20 h. The reaction solution was washed consecutively with 0.1 N HCl (100 mL) and water (2 x 100 mL), dried (MgSO₄), and concentrated to give a viscous oil (5.79 g). Flash chromatography on a 5 x 21 cm silica gel column (85/15 EtOAc/hexanes, followed by 100% EtOAc) gave a colorless amorphous solid. Recrystallization from toluene (85 mL) gave the title compound as a colorless solid (2.52 g, 59%). MP: 104.5-106.5 °C; ¹H NMR (CDCl₃): 8.00-7.90 (m, 4H), 7.72-7.64 (m, 4H), 7.46-7.24 (m, 5H), 5.96-5.88 (m, 1H), 5.86-5.73 (m, 1H), 5.41 (s, 1H), 5.16-5.00 (m, 3H), 4.15-4.02 (m, 1H), 3.68-3.39 (m, 17H), 3.34-3.22 (m, 2H), 3.13-3.03 (m, 2H), 1.80-1.62 (m, 4H), 1.39 (s, 9H); ¹³C NMR (CDCl₃): 170.2, 156.5, 156.1, 143.9, 143.0, 140.4, 139.4, 136.7, 128.4, 128.1, 128.0, 127.9, 127.9, 127.8, 127.3, 80.1, 70.6, 70.5, 70.2, 70.1, 70.0, 69.6, 66.5, 56.1, 52.9, 43.2, 42.4, 39.3, 29.4, 28.5, 28.2; MS: m/e 868.3 [M+NH₄]; High Resolution MS: Calcd for C₃₉H₅₅N₄O₁₃S₂ [M+H]: 851.3207, Found: 851.3226.




Part C - Methyl 3-((1-Methyl-4-oxo-7-(((1-(triphenylmethyl)imidazol-2-yl)amino)methyl)(3-hydroquinolyl))carbonylamino)-2-(((4-(4-(((3-(2-(2-(3-(phenylmethoxy)carbonylamino)propoxy)ethoxy)ethoxy)propyl)amino)sulfonyl)phenyl)phenyl)sulfonyl)amino)propanoate

The product of Part B, above (748 mg, 0.88 mmol) was dissolved in 25/75 TFA/DCM (15 mL) and allowed to stand

at ambient temperatures under nitrogen for 15 min. The TFA was removed under vacuum and the resulting amber oil was taken up in 50/50 ACN/water (50 mL), and treated portion wise with Bio-Rad AG-3-X4A resin, hydroxide form, to raise the pH from 2 to 6. The resin was removed by filtration and the filtrate was lyophilized to give a sticky pale yellow foam.

In a separate flask, 1-methyl-4-oxo-7-(((1-(triphenylmethyl)imidazol-2-yl)amino)methyl)hydroquinoline-3-carboxylic acid (432 mg, 0.80 mmol), TEA (0.33 mL), and HBTU (364 mg, 0.96 mmol) were dissolved in anhydrous DMF (25 mL). The resulting solution was stirred at ambient temperatures under a nitrogen atmosphere for 10 min and combined with a solution of the yellow foam in anhydrous DMF (15 mL). The DMF was removed under vacuum after 18 h to give a viscous yellow oil. This oil was taken up in EtOAc (175 mL), washed consecutively with water (25 mL), saturated NaHCO₃ (50 mL), and saturated NaCl (25 mL), dried (MgSO₄), and concentrated to give a viscous yellow oil. Purification by flash chromatography on a 7 x 25 cm silica gel column using a CHCl₃/EtOAc/MeOH step gradient (47/47/6, 46/46/8, 60/30/10) gave the title compound as a pale yellow solid (510 mg, 50%). MP: 136-140 °C; MS: m/e 1273.4 [M+H]; High Resolution MS: Calcd for C₆₈H₇₃N₈O₁₃S₂ [M+H]: 1273.4738, Found: 1273.4730.




30

Part D - 3-(((1-Methyl-4-oxo-7-(((1-(triphenylmethyl)-imidazol-2-yl)amino)methyl) (3-hydroquinolyl)) carbonylamino)-2-(((4-(4-((3-(2-(2-(3-(phenylmethoxy) carbonylamino) propoxy) ethoxy)-

ethoxy)propyl)amino)sulfonyl)phenyl)phenyl)sulfonyl)-
amino)propanoic Acid


- The product from Part C, above (295 mg, 0.232 mmol) was dissolved in a mixture of peroxide-free THF (12 mL),
5 water (1.8 mL), and 3 N LiOH (1.2 mL), and stirred at ambient temperatures under a nitrogen atmosphere for 30 min. The THF was removed under vacuum and the resulting mixture was dissolved in CHCl₃ (75 mL) and water (50 mL). The aqueous layer was adjusted to pH 3 with 0.5 N HCl and
10 the layers were thoroughly mixed. The aqueous layer was extracted with additional CHCl₃ (2 x 25 mL). The combined CHCl₃ extracts were washed with saturated NaCl (50 mL), dried (MgSO₄), and concentrated to give the title compound as a pale yellow solid (291 mg, 100%).
15 MS: m/e 1259.3 [M+H]; High Resolution MS: Calcd for C₆₇H₇₁N₈O₁₃S₂ [M+H]: 1259.4582, Found: 1259.4610.



- 20 Part E - 2-(((4-(4-(((3-(2-(2-(3-Aminopropoxy)ethoxy)ethoxy)propyl)amino)sulfonyl)phenyl)phenyl)sulfonyl)-amino)-3-((7-((imidazol-2-yl)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic Acid

- The product from Part D, above (279 mg, 0.222 mmol) was dissolved in degassed TFA (30 mL) and treated with
25 Et₃SiH (0.424 mL, 2.66 mmol). The solution was heated at 70 °C under a nitrogen atmosphere for 1 h and concentrated to a viscous oil. This oil was dissolved in water (20 mL) and washed with ether (2 x 20 mL). The
30 combined ether washings were back-extracted with water (10 mL). The combined water extracts were diluted with an equal volume of ACN and treated with Bio-Rad AG-3-X4A resin, hydroxide form to raise the pH from 4 to 6. The resin was removed by filtration and the filtrate was
35 lyophilized to give the title compound as a colorless solid (220 mg). MS: m/e 883.4 [M+H], 442.5 [M+2H]; High


Resolution MS: Calcd for $C_{40}H_{51}N_8O_{11}S_2$ $[M+H]^+$: 833.3118,
Found: 833.3118.



5

Part F - 2-(((4-(4-(((3-(2-(2-(3-((6-((1-Aza-2-(2-sulphophenyl)vinyl)amino)(3-pyridyl))carbonylamino)-propoxy)ethoxy)-ethoxy)propyl)amino)sulfonyl)phenyl)phenyl)sulfonyl)-amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic Acid
Trifluoroacetate Salt

A solution of the product from Part F, above (15 mg, 0.0135 mmol), TEA (0.007 mL), and 2-(2-aza-2-((5-((2,5-dioxopyrrolidinyl)carbonyl)(2-pyridyl))amino)vinyl)-benzenesulfonic acid (9.0 mg, 0.0204 mmol) in anhydrous DMF (2.5 mL) was allowed to stand at ambient temperatures under a nitrogen atmosphere for 22 h. The DMF was removed under vacuum and the glassy solid was dissolved in 20% ACN and purified by preparative HPLC on a Vydac C-18 column (22 x 250 mm) using 0.1% TFA in water for 5 min followed by a 2.52%/min gradient of 0 to 63% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 21.2 min was collected and lyophilized to give the title compound as a colorless powder (3.5 mg, 20%). MS: m/e 1186.7 $[M+H]^+$; High Resolution MS: Calcd for $C_{53}H_{60}N_{11}O_{15}S_3$ $[M+H]^+$: 1186.3432, Found: 1186.3410.




•TFA

30


Example 2

3-((7-((Imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((4-(4-((3-(2-(2-(3-(2-(1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)cyclododecyl)acetyl amino)-propoxy)ethoxy)ethoxy)propyl)amino)sulfonyl)-phenyl)phenyl)sulfonyl)amino)propanoic Acid
Bis(trifluoroacetate) Salt

Part A - Phenylmethyl 2-(1,4,7,10-Tetraaza-4,7,10-tris(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)acetate
A solution of tert-butyl (1,4,7,10-tetraaza-4,7-bis(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)acetate (0.922 g, 1.79 mmol), TEA (1.8 mL) and benzyl bromoacetate (0.86 mL, 5.37 mmol) in anhydrous DMF (24 mL) was stirred at ambient temperatures under a nitrogen atmosphere for 24 h. The DMF was removed under vacuum and the resulting oil was dissolved in EtOAc (300 mL). This solution was washed consecutively with water (2 x 50 mL) and saturated NaCl (50 mL), dried (MgSO₄), and concentrated to give the title compound as an amorphous solid (1.26 g). MS: m/e 663.5 [M+H].




Part B - 2-(1,4,7,10-tetraaza-4,7,10-tris(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)acetic acid
The product from Part A, above (165 mg, 0.25 mmol) was hydrogenolyzed over 10% Pd on carbon (50 mg) in EtOH (15 mL) at 60 psi for 24 h. The catalyst was removed by filtration through filter aid and washed with EtOH. The filtrates were concentrated to give the title compound as an amorphous solid (134 mg, 94%). MS: m/e 573.5 [M+H].



Part C - Methyl 3-((7-((Imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((4-(4-
 5 (((3-(2-(2-(3-(2-(1,4,7,10-tetraaza-4,7,10-tris(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)acetyl-amino)-propoxy)ethoxy)-ethoxy)propyl)amino)sulfonyl)phenyl)phenyl)sulfonyl)-amino)propanoate Pentakis(trifluoroacetate) Salt

10 A solution of the product of Example 1, Part C (68 mg, 0.0534 mmol) and Et₃SiH (0.051 mL, 0.32 mmol) in degassed TFA (5.0 mL) was stirred at 70 °C under a nitrogen atmosphere for 1 h and concentrated to dryness. The resulting amber oil was dissolved in anhydrous DMF (2
 15 mL) and treated with TEA until basic to pH paper. A solution of the product of Part B, above (46 mg, 0.080 mmol) in anhydrous DMF (1.0 mL) was added, followed by HBTU (24 mg, 0.064 mmol), and the solution was stirred at ambient temperatures under a nitrogen atmosphere for 3 h.
 20 The DMF was removed under vacuum and the residue was dissolved in 50% ACN and purified by preparative HPLC on a Vydac C-18 column (22 x 250 mm) using a 2.1%/min gradient of 0 to 63% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 23.8
 25 min was collected and lyophilized to give the title compound as a colorless powder (16 mg, 15%). MS: m/e 1451.7 [M+H]; High Resolution MS: Calcd for C₆₉H₁₀₃N₁₂O₁₈S₂ [M+H]: 1451.6954, Found: 1451.698.



• 5 TFA


30

Part D - 3-((7-((Imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((4-(4-(((3-(2-(2-

(3-(2-(1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)cyclododecyl)acetylamino)-propoxy)ethoxy)ethoxy)propyl)amino)sulfonyl)-phenyl)phenyl)sulfonyl)amino)propanoic Acid

5 Bis(trifluoroacetate) Salt

The product of Part C, above (16 mg, 0.0102 mmol) was dissolved in a mixture of peroxide-free THF (1 mL), water (0.115 mL), and 3 N LiOH (0.075 mL), and stirred at ambient temperatures under a nitrogen atmosphere for 24
 10 h. The reaction was concentrated to give an oily solid. This solid was dissolved in 50% ACN and purified by preparative HPLC on a Vydac C-18 column (22 x 250 mm) using a 2.52%/min gradient of 0 to 63% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product
 15 peak eluting at 24.0 min was collected and lyophilized to give a colorless powder (6.0 mg). This solid was dissolved in degassed TFA (2.0 mL) and Et₃SiH (0.050 mL), stirred at 70 °C under a nitrogen atmosphere for 4.5 h, and concentrated to dryness. The resulting oil was
 20 dissolved in 25% ACN and purified by preparative HPLC on a Vydac C-18 column (22 x 250 mm) using a 1.5%/min gradient of 0 to 45% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 19.0 min was collected and lyophilized to give the title
 25 compound as a colorless powder (2.0 mg, 17%). MS: m/e 1269.5 [M+H], 635.5 [M+2H], 424.3 [M+3H]; High Resolution MS: Calcd for C₅₆H₇₇N₁₂O₁₈S₂ [M+H]: 1269.4920, Found: 1269.4950.



• 2 TFA

30

Example 3


2-(((4-(3-(N-(3-(2-(3-((6-((1-Aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)-

propoxy)ethoxy)ethoxy)propyl)carbamoyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)propanoic Acid Trifluoroacetate Salt

5

Part A - Ethyl 4-(3,5-Dimethylphenoxy)butanoate

Sodium metal (17.12 g, 0.744 mol) was added to anhydrous EtOH (350 mL) and stirred until dissolved. 3,5-Dimethylphenol was added and the solution was stirred
10 15 min at ambient temperatures. Ethyl 4-bromoacetate (58.7 mL, 0.41 mol) was added and the solution was stirred at ambient temperatures under a nitrogen atmosphere for 28 h. The EtOH was removed under vacuum and the oily solid was partitioned between water (1 L)
15 and EtOAc (500 mL). The aqueous layer was extracted with additional EtOAc (500 mL). The combined EtOAc extracts were washed consecutively with saturated NaHCO₃ (300 mL) and saturated NaCl (300 mL), dried (MgSO₄), and concentrated to give an amber liquid. This liquid was
20 vacuum fractional distilled through a 15 cm Vigreux column. The main fraction was collected from 91-117 °C/6 mm Hg to give the title compound as a colorless liquid (77.77 g, 89%). ¹H NMR (CDCl₃): 6.59 (s, 1H), 6.52 (s, 2H), 4.16 (q, J = 7.16 Hz, 2H), 3.98 (t, J = 6.14 Hz, 2H), 2.49 (t, J = 7.34 Hz, 2H), 2.28 (s, 6H), 2.11-2.07
25 (m, 2H), 1.26 (t, J = 7.16 Hz, 3H); Anal. calcd for C₁₄H₂₀O₃: C, 71.16; H, 8.53, Found: C, 71.35; H, 8.59.




30

Part B - 4-(3,5-Dimethylphenoxy)butanoic Acid

The product of part A, above (75.52 g, 0.320 mol) and KOH pellets (38.5 g, 0.584 mol) were dissolved in absolute EtOH (1.50 L) and heated at reflux for 3 h. The
35 solution was concentrated to a colorless solid, which was taken up in water (2.0 L) and washed with ether (2 x 750


mL). The aqueous layer was adjusted to pH 1 with concd HCl (55 mL) and the resulting oily ppt was extracted into EtOAc (2 x 500 mL). The combined EtOAc extracts were washed consecutively with water (300 mL) and saturated NaCl, dried (MgSO₄), and concentrated to give a colorless solid (64.13 g). Recrystallization from hexanes (500 mL) gave the title compound as a colorless solid (59.51 g, 89%). MP: 66-68.5 °C; ¹H NMR (CDCl₃): 11.70 (bs, 1H), 6.59 (s, 1H), 6.52 (s, 2H), 3.99 (t, J = 6.06 Hz, 2H), 2.57 (t, J = 7.29 Hz, 2H), 2.28 (s, 6H), 2.12-2.08 (m, 2H); Anal. calcd for C₁₂H₁₆O₃: C, 69.21; H, 7.74, Found: C, 69.23; H, 7.40.



15


Part C - 4-(4-(Chlorosulfonyl)-3,5-dimethylphenoxy)butanoic Acid

A solution of the product of Part B, above (20.8 g, 0.100 mol) in CHCl₃ (100 mL) was cooled to 0 °C and treated with chlorosulfonic acid (36 mL, 0.54 mol) dropwise and with rapid stirring while keeping the temperature of the reaction at 0 °C. The resulting gelatinous mixture was stirred an additional 10 min and poured onto an ice/water mixture (600 mL). The resulting solid ppt was collected by filtration, washed with water (3 x 75 mL), and dried under vacuum to give a colorless solid (12.52 g). MP: 114-115 °C (with decomp); ¹H NMR (CDCl₃): 13.84 (bs, 1H), 6.50 (s, 2H), 3.91 (t, J = 6.48 Hz, 2H), 2.48 (s, 6H), 2.32 (t, J = 7.32 Hz, 2H), 1.89-1.84 (m, 2H); IR (KBr cm⁻¹): 1705 (s), 1370 (s), 1175 (s); MS: m/e 305.1 [M-H].



Part D - 4-(4-(((2-((tert-Butoxy)carbonylamino)-1-(methoxycarbonyl)ethyl)amino)sulfonyl)-3,5-dimethylphenoxy)butanoic Acid

A solution of N- β -Boc-L- α,β -diaminopropionic acid methyl ester hydrochloride (568 mg, 2.10 mmol) and DIEA (0.73 mL, 4.2 mmol) in DCM (5 mL) was cooled to 0 °C and treated with a suspension of the product of Part C, above (656 mg, 2.10 mmol) in DCM (20 mL) in small portions over a 15 min period. The reaction was stirred at ambient temperatures under a nitrogen atmosphere for 18 h. The reaction was diluted with DCM (100 mL) and washed with water (3 x 75 mL). The organic phase was dried (MgSO₄), and concentrated to give crude product (698 mg), which was purified by preparative HPLC on a Vydac C-18 column (50 x 250 mm) using a 0.96%/min gradient of 18 to 58.5% ACN containing 0.1% TFA at a flow rate of 80 mL/min. The main product fraction eluting at 23.8 min was collected adjusted to pH 3, partially concentrated to remove ACN, and extracted with DCM (2 x 100 mL). The DCM extracts were dried (MgSO₄) and concentrated to give the title compound as a colorless solid (297 mg, 29%). ¹H NMR (CDCl₃): δ 6.61 (s, 2H), 5.66 (d, J = 7.2 Hz, 1H), 4.90 (s, 1H), 4.03 (bs, 2H), 3.86 (bs, 1H), 3.59 (s, 3H), 3.49 (bs, 2H), 2.62 (s, 6H), 2.58-2.51 (m, 2H), 2.18-2.07 (m, 2H), 1.41 (s, 9H); MS: m/e 489.4 [M+H]; High Resolution MS: Calcd for C₂₁H₃₃N₂O₉S [M+Na]: 511.1726, Found: 511.1747; Anal. calcd for C₂₁H₃₂N₂O₉S: C, 51.62; H, 6.61; N, 5.74, Found: C, 51.47; H, 6.27; N, 5.48.




30

Part E - Methyl 3-((tert-Butoxy)carbonylamino)-2-(((2,6-dimethyl-4-(3-(N-(3-(2-(2-(3-((phenylmethoxy)carbonylamino)propoxy)ethoxy)-

ethoxy)propyl)carbamoyl)propoxy)phenyl)-
sulfonyl)amino)propanoate

A solution of the product from Part D, above (233 mg, 0.477 mmol), the product of Example 1, Part A (190 mg, 0.536 mmol), TEA (0.2 mL, 1.43 mmol), and HBTU (226 mg, 0.701 mmol) in anhydrous DMF (8 mL) was stirred at ambient temperatures under a nitrogen atmosphere for 1 h. The DMF was removed under vacuum and the oily residue was taken up in EtOAc (50 mL) and washed consecutively with 0.1 N HCl (35 mL), water (35 mL), and saturated NaCl (35 mL), dried (MgSO₄), and concentrated to give crude product as a yellow viscous oil. Flash chromatography on a 3 x 18 cm silica gel column (EtOAc/MeOH, 95/5) gave the title compound as a colorless viscous oil (393 mg, 100%).

¹H NMR (CDCl₃): δ 7.34-7.28 (m, 5H), 6.60 (s, 2H), 6.26 (bs, 1H), 5.67 (bs, 1H), 5.29 (bs, 1H), 5.08 (s, 2H), 4.88 (bs, 1H), 3.99 (t, J = 6.1 Hz, 2H), 3.88-3.84 (m, 1H), 3.62-3.40 (m, 17H), 3.37-3.26 (m, 4H), 2.62 (s, 6H), 2.32 (t, J = 7.2 Hz, 2H), 2.08 (t, J = 6.3 Hz, 2H), 1.79-1.70 (m, 4H), 1.41 (s, 9H); MS: m/e 825.5 [M+H]; High Resolution MS: Calcd for C₃₉H₆₁N₄O₁₃S [M+H]: 825.3955, Found: 825.3940.




25

Part F - Methyl 3-Amino-2-(((2,6-dimethyl-4-(3-(N-(3-(2-(2-(3-((phenylmethoxy)carbonylamino)propoxy)ethoxy)-ethoxy)propyl)carbamoyl)propoxy)phenyl)-sulfonyl)amino)propanoate


30 The product of Part E, above (750 mg, 0.91 mmol) was dissolved in 4 M HCl/dioxane (25 mL) and stirred at ambient temperatures for 1 h. The solution was diluted with ether (500 mL) and the resulting gummy ppt was

trituated with fresh ether (2 x 250 mL). The gummy solid was dissolved in water (100 mL) and adjusted to pH 9 with NaHCO₃, causing an oily ppt to form. This ppt was extracted into DCM (2 x 75 mL). The DCM extracts were
 5 dried (MgSO₄) and concentrated to give the title compound as a colorless oil (386 mg, 56%). MS: m/e 725.5 [M+H].




- 10 Part G - Methyl 2-(((2,6-Dimethyl-4-(3-(N-(3-(2-(2-(3-
 ((phenylmethoxy)carbonylamino)propoxy)ethoxy)ethoxy)-
 propyl)carbamoyl)propoxy)phenyl)sulfonyl)amino)-3-((1-
 methyl-4-oxo-7-(((1-(triphenylmethyl)imidazol-2-
 yl)amino)methyl)(3-
 15 hydroquinolyl))carbonylamino)propanoate.

A solution of 1-methyl-4-oxo-7-(((1-(triphenylmethyl)imidazol-2-yl)amino)methyl)hydroquinoline-3-carboxylic acid (274 mg, 0.51 mmol), TEA (0.22 mL, 1.52 mmol), and HBTU (192 mg, 0.51 mmol) in anhydrous DMF (3 mL) was stirred at ambient
 20 temperatures for 5 min. A solution of the product of Part F, above (367 mg, (0.51 mmol) in anhydrous DMF (7 mL) was added and the resulting solution was stirred at ambient temperatures under a nitrogen atmosphere for 2 h.
 25 The DMF was removed under vacuum and the resulting oily solid was dissolved in EtOAc (150 mL). The EtOAc solution was washed consecutively with water (50 mL), saturated NaHCO₃ (25 mL), and saturated NaCl (25 mL), dried (MgSO₄), and concentrated to give a yellow solid.
 30 Purification by flash chromatography on a silica gel column using a EtOAc/MeOH step gradient (95/5, 92.5/7.5) gave the title compound as a pale yellow solid (254 mg, 43%). MS: m/e 1247.7 [M+H], 624.6 [M+2H].



Part H - 2-(((2,6-Dimethyl-4-(3-(N-(3-(2-(2-(3-
 5 ((phenylmethoxy)carbonylamino)propoxy)ethoxy)ethoxy)-
 propyl)carbamoyl)propoxy)phenyl)sulfonyl)amino)-3-((1-
 methyl-4-oxo-7-(((1-(triphenylmethyl)imidazol-2-
 yl)amino)methyl)(3-hydroquinolyl))carbonylamino)propanoic
 Acid

10 The product of Part G, above (60.0 mg, 0.048 mmol)
 was dissolved in a mixture of peroxide-free THF (2.5 mL),
 water (0.37 mL), and 3 N LiOH (0.244 mL), and stirred at
 ambient temperatures under a nitrogen atmosphere for 30
 min. The THF was removed under vacuum and the resulting
 15 mixture was dissolved in CHCl₃ (25 mL) and water (20 mL).
 The aqueous layer was adjusted to pH 3 with 0.1 N HCl and
 the layers were thoroughly mixed. The aqueous layer was
 extracted with additional CHCl₃ (2 x 20 mL). The
 combined CHCl₃ extracts were washed with saturated NaCl
 20 (30 mL), dried (MgSO₄), and concentrated to give the
 title compound as a pale yellow solid (44.0 mg, 74%).
 MS: m/e 1233.7 [M+H]; High Resolution MS: Calcd for
 C₆₇H₇₇N₈O₁₃S [M+H]: 1233.5330, Found: 1233.5330.




25

Part I - 2-(((4-(3-(N-(3-(2-(2-(3-Aminopropoxy)ethoxy)-ethoxy)propyl)carbamoyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic

5 Acid

The product of Part H, above (42.1 mg, 0.0341 mmol) and Et₃SiH (0.033 mL, 0.205 mmol) were dissolved in degassed TFA (3.5 mL), heated at 70 °C under a nitrogen atmosphere for 1 h, and concentrated to give a viscous
 10 amber oil. This oil was dissolved in water (20 mL) and washed with ether (2 x 20 mL). The combined ether washings were back-extracted with water (10 mL). The combined water extracts were diluted with an equal volume of ACN and treated with Bio-Rad AG-3-X4A resin, hydroxide
 15 form to raise the pH from 4 to 6. The resin was removed by filtration and the filtrate was lyophilized to give the title compound as a colorless solid (34 mg). MS: m/e 857.5 [M+H], 429.4 [M+2H].




20

Part J - 2-(((4-(3-(N-(3-(2-(2-(3-((6-((1-Aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)-propoxy)ethoxy)ethoxy)propyl)carbamoyl)propoxy)-2,6-
 25 dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)propanoic Acid Trifluoroacetate Salt

A solution of the product from Part I, above (30 mg, 0.035 mmol), DIEA (0.018 mL, 0.105 mmol) and 2-(2-aza-2-
 30 ((5-((2,5-dioxopyrrolidinyl)carbonyl)(2-pyridyl))-amino)vinyl)benzenesulfonic acid (18.5 mg, 0.042 mmol) in anhydrous DMF (1.5 mL) was allowed to stand at ambient temperatures under a nitrogen atmosphere for 20 h. The

DMF was removed under vacuum and the amber oil was dissolved in 50% ACN and purified by preparative HPLC on a Zorbax C-18 RX column (21.2 x 250 mm) using a 1.5%/min gradient of 0 to 45% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 21.0 min was collected and lyophilized to give the title compound as a colorless powder (8.9 mg, 20%). MS: m/e 1160.6 [M+H], 581.0 [M+2H].



10

Example 4


3-((1-(3-((6-((1-Aza-2-(2-sulfohenyl)vinyl)amino)(3-pyridyl))carbonylamino)propyl)-7-((imidazole-2-ylamino)-methoxy)-4-oxo(3-hydroquinolyl))carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic Acid
Trifluoroacetate Salt

20 Part A - Ethyl 1-(3-((tert-Butoxy)carbonylamino)propyl)-7-bromo-4-oxohydroquinoline-3-carboxylate

A mixture of ethyl 7-bromo-4-oxohydroquinoline-3-carboxylate (6.28 g, 0.0212 mol), (tert-butoxy)-N-(3-bromopropyl)formamide (30.3 g, 0.127 mol), and anhydrous K₂CO₃ (12.5 g, 0.904 mol) in anhydrous DMF (200 mL) was stirred at 60 °C under a nitrogen atmosphere for 4 h, and then at ambient temperatures for 72 h. The DMF was removed under vacuum and the resulting oily solid was dissolved in EtOAc (500 mL). The EtOAc solution was washed consecutively with water (500 mL), saturated NaHCO₃ (500 mL), and saturated NaCl (500 mL), dried (MgSO₄), and concentrated to give a red oil. This oil

30


- was taken up in EtOAc (250 mL) and cooled, causing a solid ppt to form. This ppt was collected by filtration, washed with cold EtOAc, and dried to give the title compound as a colorless solid (6.25 g, 65%). MP: 140-142 °C; ¹H NMR (CDCl₃): 8.49 (s, 1H), 8.39 (d, J = 8.6 Hz, 1H), 7.58 (s, 1H), 7.53 (d, J = 8.6 Hz, 1H), 4.72 (bs, 1H), 4.39 (q, J = 7.1 Hz, 2H), 4.20 (t, J = 7.6 Hz, 2H), 3.28-3.24 (m, 2H), 2.10-2.06 (m, 2H), 1.46 (s, 9H), 1.40 (t, J = 7.1 Hz, 3H); MS: m/e 455.2. [M+H]⁺; High Resolution MS: Calcd for C₂₀H₂₆BrN₂O₅ [M+H]⁺: 453.1025, Found: 453.1028.



- Part B - Ethyl 1-(3-((tert-butoxy)carbonylamino)propyl)-4-oxo-7-vinylhydroquinoline-3-carboxylate

- The product from Part A, above (2.98 g, 6.60 mmol) was dissolved in toluene (50 mL) at a temperature of 100 °C and treated with tetrakis(triphenylphosphine)palladium(0) (152 mg, 0.132 mmol). After 5 min the mixture was treated with tributyl(vinyl)tin (1.93 mL, 6.60 mmol) and stirred 4.5 h at 100 °C under a nitrogen atmosphere, and 18 h at ambient temperatures. Additional tributyl(vinyl)tin (0.386 mL) and tetrakis(triphenylphosphine)palladium(0) (152 mg) were added and the mixture was heated at 100 °C for an additional 17 h. The toluene was removed under vacuum and the solid residue was triturated with ether to give the title compound as a pale green solid (1.67 g, 63%). MP: 133-135 °C; ¹H NMR (CDCl₃): 8.52 (d, J = 8.4 Hz, 1H), 8.51 (s, 1H), 7.55 (d, J = 8.4 Hz, 1H), 7.38 (s, 1H), 6.88-6.82 (m, 1H), 5.97 (d, J = 17.4 Hz, 1H), 5.51 (d, J = 10.8 Hz, 1H), 4.75 (bs, 1H), 4.42 (q, J = 7.2 Hz, 2H), 4.27 (t, J = 7.8 Hz, 2H), 3.6-3.25 (m, 2H), 2.16-2.11 (m, 2H), 1.49 (s, 9H), 1.45 (t, J = 7.2 Hz, 3H); MS:


m/e 401.3 [M+H]; High Resolution MS: Calcd for C₂₂H₂₉N₂O₅
[M+H]: 401.2076, Found: 401.2075.



5


Part C - Ethyl 1-(3-((tert-butoxy)carbonylamino)propyl)-
7-formyl-4-oxohydroquinoline-3-carboxylate

A solution of the product of Part B, above (1.50 g,
3.75 mmol) in dioxane (119 mL) and water (39 mL) was
10 treated with a solution of osmium tetroxide (19.6 mg,
0.077 mmol) in dioxane (0.600 mL) and stirred at ambient
temperatures under a nitrogen atmosphere for 5 min.
Sodium periodate (2.40 g, 11.2 mmol) was added and the
stirred at ambient temperatures for 2 h. The dioxane was
15 removed under vacuum and the residue was taken up in DCM
(500 mL). The DCM solution was washed consecutively with
water (500 mL) and saturated NaCl (500 mL), dried
(MgSO₄), and concentrated to give the title compound as
an orange oily solid (1.52 g, 100%). ¹H NMR (CDCl₃):
20 10.17 (s, 1H), 8.68 (d, J = 8.2 Hz, 1H), 8.64 (s, 1H),
8.01 (s, 1H), 7.88 (d, J = 8.2 Hz, 1H), 4.82 (bs, 1H),
4.41-4.35 (m, 4H), 3.28 (s, 2H), 2.15-2.07 (m, 2H), 1.45
(s, 9H), 1.41 (t, J = 7.1 Hz, 3H); MS: m/e 403.3 [M+H];
High Resolution MS: Calcd for C₂₁H₂₇N₂O₆ [M+H]: 403.1870,
25 Found: 403.1875.



Part D - Ethyl 1-(3-((tert-butoxy)carbonylamino)propyl)-
4-oxo-7-(((1-(triphenylmethyl)imidazole-2-
30 yl)amino)methyl)hydroquinoline-3-carboxylate

A solution of the product of Part C, above (544 mg, 1.35 mmol) and 1-(triphenylmethyl)imidazole-2-ylamine (456 mg, 1.35 mmol) in toluene (60 mL) was heated at reflux under a nitrogen atmosphere with removal of water
5 for 5 h. The solution was cooled, treated with $\text{Na}(\text{OAc})_3\text{BH}$ (1.14 g, 5.38 mmol) and stirred at ambient temperatures for 18 h. The mixture was diluted with EtOAc (400 mL), washed consecutively with water (500 mL) and saturated NaCl (500 mL), dried (MgSO_4), and
10 concentrated to give an orange solid. This solid was dissolved in 50% ACN and purified by preparative HPLC on a Vydac C-18 column (50 x 250 mm) using a 0.60%/min gradient of 18 to 52% ACN containing 0.1% TFA at a flow rate of 49 mL/min. The main product peak eluting at 30.8
15 min was collected and lyophilized to give the title compound as a pale yellow solid (407 mg, 60%). MS: m/e 712.4 [M+H]; High Resolution MS: Calcd for $\text{C}_{43}\text{H}_{46}\text{N}_5\text{O}_5$ [M+H]: 712.3499, Found: 712.3485.




20

Part E - 1-(3-((tert-Butoxy)carbonylamino)propyl)-4-oxo-7-(((1-(triphenylmethyl)imidazole-2-yl)amino)methyl)-hydroquinoline-3-carboxylic Acid

25 A mixture of the product of Part D, above (997 mg, 1.40 mmol), water (7.3 mL), 3 N LiOH (3.5 mL), and THF (50 mL) was stirred at ambient temperatures under a nitrogen atmosphere for 3 h. The THF was removed under vacuum and the resulting mixture was dissolved in CHCl_3
30 (500 mL) and water (100 mL). The aqueous layer was adjusted to pH 3 with 1.0 N HCl and the layers were thoroughly mixed. The organic layer was washed consecutively with water (500 mL) and saturated NaCl (500 mL), dried (MgSO_4), and concentrated to give the title


compound as a pale yellow solid (998 mg). MP: 153-160 °C; ¹H NMR (CDCl₃): δ 14.83 (s, 1H), 8.76 (s, 1H), 8.68 (s, 1H), 8.24 (d, J = 6 Hz, 1H), 7.49-7.35 (m, 9H), 7.12-7.10 (m, 6H), 6.82 (s, 1H), 6.52 (s, 1H), 6.24 (d, J = 6 Hz, 1H), 5.75 (bs, 1H), 4.87-4.83 (m, 2H), 4.77 (bs, 1H), 4.51 (t, J = 9 Hz, 2H), 3.38 (s, 2H), 2.23 (s, 2H), 1.42 (s, 9H); MS: m/e 684.3 [M+H]; High Resolution MS: Calcd for C₄₁H₄₂N₅O₅ [M+H]: 684.3186, Found: 684.3181.



10


Part F - Methyl 3-((1-(3-((tert-Butoxy)carbonylamino)propyl)-4-oxo-7-((1-(triphenylmethyl)imidazole-2-yl)amino)methyl)(3-hydroquinolyl))carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoate

A solution of the product of Part E, above (300 mg, 0.437 mmol), TEA (0.243 mL, 1.75 mmol), and HBTU (230 mg, 0.606 mmol) in anhydrous DMF (4 mL) was stirred at ambient temperatures for 5 min. A solution of methyl 3-amino-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoate hydrochloride (184 mg, 0.637 mmol) in anhydrous DMF (3 mL) was added and the solution was stirred at ambient temperatures under a nitrogen atmosphere for 2 h. The solution was diluted with EtOAc (200 mL) and washed consecutively with water (2 x 50 mL), saturated NaHCO₃ (50 mL), and saturated NaCl (50 mL), dried (MgSO₄), and concentrated to give a viscous amber oil. Purification by flash chromatography on a 2.5 x 24 cm silica gel column using a EtOAc/MeOH step gradient (98/2, 95/5, 75/25) gave the title compound as a pale yellow oil (330 mg, 78%). MS: m/e 966.6 [M+H]; High Resolution MS: Calcd for C₅₄H₆₀N₇O₈S [M+H]: 966.4224, Found: 966.4224.



Part G - 3-((1-(3-((tert-Butoxy)carbonylamino)propyl)-4-oxo-7-(((1-(triphenylmethyl)imidazole-2-yl)amino)methyl)(3-hydroquinolyl))carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic Acid

A solution of the product of Part F, above (51 mg, 0.052 mmol), water (0.27 mL), and 3 N LiOH (0.13 mL) in MeOH (2 mL) was allowed to stand at ambient temperatures for 3.5 h and concentrated under vacuum. The resulting solid was dissolved in water (10 mL) and adjusted to pH 3 with 1.0 N HCl. The aqueous mixture was extracted with DCM (2 x 30 mL). The combined DCM extracts were washed with saturated NaCl (30 mL), dried (MgSO₄), and concentrated to give the title compound as a colorless solid (72 mg). MS: m/e 952.5 [M+H]; High Resolution MS: Calcd for C₅₃H₅₈N₇O₈S [M+H]: 952.4067, Found: 952.4056.



20


Part H - 3-((1-(3-Aminopropyl)-7-((imidazole-2-ylamino)-methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic Acid
Bis(trifluoroacetate) Salt

25 The product of Part I, above (0.052 mmol) and Et₃SiH (0.042 mL, 0.26 mmol) were dissolved in degassed TFA (2 mL), heated at 70 °C for 2.5 h, and concentrated to give an amber oil. This oil was dissolved in water (25 mL) and washed with ether (2 x 15 mL). The combined ether washings were back-extracted with water (15 mL). The

30

combined water extracts were lyophilized to give the title compound as a colorless powder (34 mg, 78%). MS: m/e 610.4 [M+H]; High Resolution MS: Calcd for C₂₉H₃₆N₇O₆S [M+H]: 610.2448, Found: 610.2462.


5



Part I - 3-((1-(3-((6-((1-Aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)propyl)-7-((imidazole-2-ylamino)-methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic Acid
Trifluoroacetate Salt

A solution of the product of Part H, above (13.7 mg, 0.0163 mmol), TEA (0.015 mL, 0.108 mmol), and 2-(2-aza-2-((5-((2,5-dioxopyrrolidinyl)carbonyl)(2-pyridyl))-amino)vinyl)benzenesulfonic acid (8.2 mg, 0.0186 mmol) in anhydrous DMF (2.0 mL) was allowed to stand at ambient temperatures under a nitrogen atmosphere for 24 h. The DMF was removed under reduced pressure and the amber oil was dissolved in 50% ACN and purified by preparative HPLC on a Vydac C-18 column (22 x 250 mm) using 0.1% TFA in water for 5 min followed by a 2.52%/min gradient of 0 to 63% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 21.4 min was collected and lyophilized to give the title compound as a colorless powder (12.5 mg, 75%). MS: m/e 913.3 [M+H]; High Resolution MS: Calcd for C₄₂H₄₅N₁₀O₁₀S₂ [M+H]: 913.2761, Found: 913.2751.


30

Example 5

5 3-((1-(3-((6-((1-Aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)propyl)-7-(((1-hydroxyimidazole-2-yl)amino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-
 10 ((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic Acid
 Trifluoroacetate Salt

Part A - Methyl 3-((1-(3-Aminopropyl)-7-((imidazole-2-ylamino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-
 ((2,4,6-trimethylphenyl)sulfonyl)amino)propanoate
 Bis(trifluoroacetate) Salt


15 A solution of the product of Example 4, Part F (120 mg, 0.124 mmol) and Et₃SiH (0.99 mL, 6.20 mmol) in TFA (10 mL) was heated at 70 °C for 1 h, and concentrated to give an amber oil. This oil was dissolved in water (50 mL) and washed with ether (2 x 30 mL). The combined
 20 ether washings were back-extracted with water (20 mL). The combined water extracts were lyophilized to give the title compound as a colorless powder (105 mg, 100%). MS: m/e 624.4 [M+H]; High Resolution MS: Calcd for C₃₀H₃₈N₇O₆S [M+H]: 624.2604, Found: 624.2608.



Part B - 3-((1-(3-Aminopropyl)-7-(((1-hydroxyimidazol-2-yl)amino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-

((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic Acid
Trifluoroacetate Salt

- A mixture of the product of Part A, above (105 mg, 0.126 mmol), water (3.0 mL), and 3 N LiOH (1.82 mL) in
5 peroxide-containing THF (4 mL) was allowed to stand at ambient temperatures for 1 h and concentrated under vacuum. The resulting solid was dissolved in water (10 mL) and adjusted to pH 5 with 1.0 N HCl. Insoluble impurities were removed by filtration and the filtrate
10 was lyophilized to give a colorless solid. This solid was dissolved in water and purified by preparative HPLC on a Vydac C-18 column (22 x 250 mm) using 0.1% TFA in water for 5 min followed by a 2.52%/min gradient of 0 to 63% ACN containing 0.1% TFA at a flow rate of 20 mL/min.
15 The main product peak eluting at 19.5 min was collected and lyophilized to give the title compound as a colorless powder (10.0 mg, 11%). MS: m/e 314.0 [M+2H]




20

- Part C - 3-((1-(3-((6-((1-Aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)propyl)-7-((1-hydroxyimidazole-2-yl)amino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-
25 (((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic Acid
Trifluoroacetate Salt

- A solution of the product of Part B, above (10.0 mg, 0.0135 mmol), TEA (0.018 mL, 0.129 mmol), and 2-(2-aza-2-((5-((2,5-dioxopyrrolidinyl)carbonyl)(2-pyridyl))-
30 amino)vinyl)benzenesulfonic acid (7.2 mg, 0.0163 mmol) in anhydrous DMF (4 mL) was allowed to stand at ambient temperatures under a nitrogen atmosphere for 20 h. The DMF was removed under vacuum and the amber oil was dissolved in 30% ACN and purified by preparative HPLC on

a Vydac C-18 column (22 x 250 mm) using 0.1% TFA in water for 5 min followed by a 2.52%/min gradient of 0 to 63% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 21.5 min was collected and lyophilized to give the title compound as a colorless powder (3.5 mg, 25%). MS: m/e 929.4 [M+H]; High Resolution MS: Calcd for C₄₂H₄₅N₁₀O₁₁S₂ [M+H]: 929.2710, Found: 929.2698.



10.

Example 6


3-((1-(3-(3-(N-(3-(2-(2-(3-((6-((1-Aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)propoxy)ethoxy)-ethoxy)propyl)carbamoyl)propanoylamino)propyl)-7-((imidazole-2-ylamino)methyl)-4-oxo(3-hydroquinolyl))-carbonylamino)-2-(((2,4,6-trimethylphenyl)sulfonyl)-amino)propanoic Acid Trifluoroacetate Salt

20

Part A - 3-(N-(3-(2-(2-(3-((tert-Butoxy)carbonylamino)-propoxy)ethoxy)ethoxy)propyl)carbamoyl)propanoic Acid


A solution of N-(3-(2-(2-(3-aminopropoxy)ethoxy)ethoxy)propyl) (tert-butoxy)formamide (as described by D. S. Wilbur et al. in *Bioconjugate Chem.* 1998, 9, 322-330) (2.00 g, 6.24 mmol), TEA (1.0 mL, 7.49 mmol), and succinic anhydride (624 mg, 6.24 mmol) in anhydrous DMF (5 mL) was stirred at ambient temperatures under a nitrogen atmosphere for 4 h. The DMF was removed under reduced pressure to give the title compound as a pale yellow oil (2.80 g). MS: m/e 839.5 [2M-H], 419.4 [M-H].

30



Part B - Methyl 3-((1-(3-(3-(N-(3-(2-(2-(3-((tert-
 Butoxy) carbonylamino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)propanoylamino)propyl-4-oxo-7-(((1-(triphenylmethyl)-
 5 imidazole-2-yl)amino)methyl)(3-hydroquinolyl))carbonylamino)-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)propanoate


The product of Example 4, Part F (46.1 mg, 0.477
 10 mmol) was dissolved in 50% TFA/DCM (2.0 mL) for 15 min at ambient temperatures and concentrated to give a yellow oil. This oil was dissolved in anhydrous DMF (1.0 mL) and made basic to pH paper with TEA. In a separate flask, the product of Part A, above (26.1 mg, 0.062
 15 mmol), TEA (0.014 mL, 0.099 mmol), and HBTU (27.7 mg, 0.074 mmol) were dissolved in anhydrous DMF (1.0 mL). The resulting solution was allowed to react for 5 min and combined with the DMF solution from the TFA deprotection reaction. The combined solutions were allowed to stand
 20 at ambient temperatures under a nitrogen atmosphere for 20 min and concentrated under vacuum. The resulting oil was dissolved in 50% ACN and purified by preparative HPLC on a Vydac C-18 column (22 x 250 mm) using a 1.8%/min gradient of 18 to 72% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 26.8
 25 min was collected and lyophilized to give the title compound as a colorless powder (44.5 mg, 68%). MS: m/e 1268.6 [M+H]; High Resolution MS: Calcd for C₆₈H₈₆N₉O₁₃S [M+H]: 1268.6065, Found: 1268.6070.



Part C - 3-((1-(3-(3-(N-(3-(2-(2-(3-((tert-Butoxy)carbonylamino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)propanoylamino)propyl)-4-oxo-7-(((1-(triphenylmethyl)-imidazole-2-yl)amino)methyl)(3-

5 hydroquinolyl))carbonylamino)-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic Acid


A solution of the product of Part B, above (31.1 mg, 0.0227 mmol), 3 n LiOH (0.091 mL), and water (0.117 mL) in MeOH (1.30 mL) was stirred at ambient temperatures for 10 8.5 h. The MeOH was removed under vacuum and the aqueous mixture was diluted with water (30 mL) and adjusted to pH 4 with 1.0 N HCl. The resulting aqueous mixture was extracted with DCM (2 x 50 mL). The combined DCM extracts were washed with saturated NaCl (50 mL), dried 15 (MgSO₄), and concentrated to give the title compound as a colorless solid (24.6 mg, 86%).



20 Part D - 3-((1-(3-(3-(N-(3-(2-(2-(3-aminopropoxy)ethoxy)ethoxy)propyl)carbamoyl)propanoylamino)propyl)-7-((imidazole-2-ylamino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic Acid
25 Bis(trifluoroacetate) Salt

A solution of the product of Part C, above (24.6 mg, 0.0194 mmol) and Et₃SiH (0.016 mL, 0.097 mmol) in TFA (2.0 mL) was heated at 70 °C under a nitrogen atmosphere for 3 h, and concentrated to give a yellow solid. This 30 solid was dissolved in water (50 mL) and washed with ether (2 x 25 mL). The aqueous layer was lyophilized to


give the title compound as a pale yellow solid (20.7 mg, 93%). MS: m/e 912.5 [M+H].



5

Part E - 3-((1-(3-(3-(N-(3-(2-(2-(3-((6-((1-Aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)propoxy)ethoxy)-ethoxy)propyl)carbamoyl)propanoylamino)propyl)-7-((imidazole-2-ylamino)methyl)-4-oxo(3-hydroquinolyl))-carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)-amino)propanoic Acid Trifluoroacetate Salt

A solution of the product of Part D, above (15.5 mg, 0.0136 mmol), TEA (0.010 mL, 0.0746 mmol), and 2-(2-aza-2-((5-((2,5-dioxopyrrolidinyl)carbonyl)(2-pyridyl))-amino)vinyl)benzenesulfonic acid (8.0 mg, 0.0182 mmol) in anhydrous DMF (2.0 mL) was allowed to stand at ambient temperatures under a nitrogen atmosphere for 24 h. The DMF was removed under vacuum and the resulting yellow oil was dissolved in 50% ACN and purified by preparative HPLC on a Vydac C-18 column (22 x 250 mm) using 0.1% TFA in water for 5 min followed by a 2.52%/min gradient of 0 to 63% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 21.7 min was collected and lyophilized to give the title compound as a colorless powder (7.2 mg, 40%). MS: m/e 1215.5 [M+H]; High Resolution MS: Calcd for C₅₆H₇₁N₁₂O₁₅S₂ [M+H]: 1215.4603, Found: 1215.4580.



Example 7


- 5 2-(2-Aza-2-(5-(N-(1,3-bis(3-(2-(2-(3-(3-(N-(3-(3-(N-(3-carboxy-2-((2,4,6-trimethylphenyl)sulfonyl)amino)ethyl)-carbamoyl)-7-((imidazole-2-ylamino)methyl)4-oxohydroquinolyl)propyl)carbamoyl)propanoylamino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)(2-pyridyl))amino)vinyl)-
 10 benzenesulfonic Acid Bis(trifluoroacetate) Salt

- Part A - N,N'-Bis(3-(2-(2-(3-(3-(N-(3-(3-(N-(3-carbomethoxy-2-((2,4,6-trimethylphenyl)sulfonyl)amino)ethyl)carbamoyl)-4-oxo-7-((1-
 15 (triphenylmethyl)imidazole-2-yl)amino)methyl)-hydroquinolyl)propyl)carbamoyl)propanoylamino)propoxy)ethoxy)ethoxy)propyl-2-((tert-butoxy)carbonylamino)pentane-1,5-diamide

- A solution of the product of Example 6, Part B (50.5
 20 mg, 0.0398 mmol) in 50/50 TFA/DCM (2 mL) was allowed to react for 20 min at ambient temperatures and concentrated to a viscous oil. This oil was taken up in anhydrous DMF and made basic to pH paper with TEA. This solution was treated with Boc-L-Glu-OH (4.5 mg, 0.0181 mmol) and HBTU
 25 (16.6 mg, 0.0438 mmol), and allowed to stand at ambient temperatures for 2 h. The DMF was removed under vacuum and the resulting oil was dissolved in 60% ACN and purified by preparative HPLC on a Vydac C-18 column (22 x 250 mm) using a 1.8%/min gradient of 18 to 72% ACN
 30 containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 21.5 min was collected and lyophilized to give the title compound as a colorless

powder (38.8 mg, 84%). MS: m/e 2306.5 [M+H-Tr], 2064.4 [M+H-2Tr], 1275.0 [M+2H]; High Resolution MS: Calcd for C₁₁₇H₁₅₄N₁₉OS₂ [M+H-Tr]: 2305.0753, Found: 2305.0770.

5




Part B - 2-Amino-N,N'-bis(3-(2-(2-(3-(3-(N-(3-(3-(N-(3-carboxy-2-((2,4,6-trimethylphenyl)sulfonyl)amino)ethyl)-carbamoyl)-7-((imidazole-2-ylamino)methyl)4-oxohydroquinolyl)propyl)carbamoyl)propanoylamino)propoxy)ethoxy)ethoxy)propyl)pentane-1,5-diamide
 10 Tris(trifluoroacetate) Salt

A solution of the product from Part A, above (38.8 mg, 0.0152 mmol), 3 N LiOH (0.075 mL), and water (0.156 mL) in MeOH (2.0 mL) was stirred at ambient temperatures for 18 h. The MeOH was removed under vacuum and the aqueous mixture was diluted with water (50 mL) and adjusted to pH 3 using 0.5 N HCl. The mixture was extracted with DCM (2 x 50 mL). The combined DCM
 15 extracts were washed with saturated NaCl (50 mL), dried (MgSO₄), and concentrated to give a colorless solid. This solid was dissolved in TFA (3.0 mL) along with Et₃SiH (0.031 mL, 0.178 mol), heated at 70 °C under a nitrogen atmosphere for 11 h, and concentrated to give a
 20 yellow oil. This oil was dissolved in water (25 mL) and washed with ether (2 x 25 mL). The aqueous solution was lyophilized to give a pale yellow solid. This solid was dissolved in water and purified by preparative HPLC on a Vydac C-18 column (22 x 250 mm) using 0.1% TFA in water
 25 for 5 min followed by a 2.52%/min gradient of 0 to 63% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The


main product peak eluting at 22.4 min was collected and lyophilized to give the title compound as a colorless powder (5.1 mg, 18%). MS: m/e 968.2 [M+2H], 646.0 [M+3H].

5



Part C - 2-(2-Aza-2-(5-(N-(1,3-bis(3-(2-(2-(3-(3-(N-(3-(3-(N-(3-carboxy-2-((2,4,6-trimethylphenyl)sulfonyl)amino)ethyl)carbamoyl)-7-((imidazole-2-ylamino)methyl)4-oxohydroquinolyl)propyl)carbamoyl)propanoylamino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)(2-pyridyl)amino)vinyl)benzenesulfonic Acid Bis(trifluoroacetate) Salt

15 A solution of the product of Part B, above (5.1 mg, 0.00224 mmol), TEA (0.002 mL, 0.0115 mmol), and 2-(2-aza-2-((5-((2,5-dioxopyrrolidinyl)carbonyl)(2-pyridyl))-amino)vinyl)benzenesulfonic acid (1.2 mg, 0.00272 mmol) in anhydrous DMF (2.0 mL) was allowed to stand at ambient
20 temperatures under nitrogen for 72 h. The DMF was removed under vacuum and the resulting oil was dissolved in 50% ACN and purified by preparative HPLC on a Vydac C-18 column (22 x 250 mm) using 0.1% TFA in water for 5 min followed by a 2.52%/min gradient of 0 to 63% ACN
25 containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 23.5 min was collected and lyophilized to give the title compound as a colorless powder (0.5 mg, 9.0%). MS: m/e 1120.0 [M+2H]; High Resolution MS: Calcd for C₁₀₄H₁₃₇N₂₂O₂₈S₃ [M+]: 2237.9055,
30 Found: 2237.9120.



Example 8

- 5 DOTA Conjugate of 3-((1-(3-(3-(N-(3-(2-(2-(N-(L-Asp-L-
Asp)3-aminopropoxy)ethoxy)ethoxy)propyl)carbamoyl)-
propanoylamino)propyl-7-((imidazole-2-ylamino)methyl)-4-
10 oxo(3-hydroquinolyl))carbonylamino)-2-((2,4,6-
trimethylphenyl)sulfonyl)amino)propanoic Acid
Bis(trifluoroacetate Salt)

Part A - Carbobenzyloxy-L-Asp(O-t-Bu)-L-Asp(O-t-Bu)-OMe

- A solution of Cbz-Asp(O-t-Bu)-OH (1.54 g, 4.76
mmol), H-Asp(O-t-Bu)-OMe•HCl (1.14 g, 4.76 mmol), DIEA
15 (1.85 mL, 10.5 mmol), and HBTU (1.99 g, 5.24 mmol) in DMF
(20 mL) was stirred at ambient temperatures for 18 h.
Water (100 mL) and EtOAc (50 mL) were added and the
layers were separated. The water layer was extracted
with EtOAc (2 x 50 mL). The combined EtOAc extracts were
20 washed consecutively with water (50 mL), 10% KHSO₄ (2 x
50 mL), and 10% NaHCO₃ (50 mL). The organic phase was
dried (MgSO₄), and concentrated to give an oily solid.
This material was triturated with ether to give the title
compound as a colorless solid (2.14g, 89%). MS: m/e
25 1017.6 [2M+H], 509.4 [M+H].


Part B - Carbobenzyloxy-L-Asp(O-t-Bu)-L-Asp(O-t-Bu)-OH

A mixture of the product of Part A, above (200 mg, 0.393 mmol), LiOH (38 mg, 0.865 mmol), water (40 mL), and THF (200 mL) was stirred at ambient temperatures for 28 h, and concentrated to remove THF. The aqueous mixture
5 was diluted with additional water (20 mL) and washed with EtOAc (20 mL). The aqueous phase was adjusted to pH 4 with 1.0 N HCl and extracted with EtOAc (20 mL). The EtOAc extract was washed with saturated NaCl (15 mL), dried (MgSO₄), and concentrated to give a colorless
10 solid. This solid was dissolved in 60% ACN and purified by preparative HPLC on a Vydac C-18 column (22 x 250 mm) using a 2.4%/min gradient of 18 to 90% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 19.0 min was collected and lyophilized to
15 give the title compound as a colorless powder (95 mg, 49%).

Part C - Methyl 3-((1-(3-(3-(N-(3-(2-(2-(N-(benzyloxycarbonyl-L-Asp(O-t-Bu)-L-Asp(O-t-Bu))3-
20 aminopropoxy)ethoxy)ethoxy)propyl)carbamoyl)-propanoylamino)propyl-4-oxo-7-((1-(triphenylmethyl)-imidazole-2-yl)amino)methyl)(3-hydroquinolyl))carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoate


25 The product of Example 6, Part B (44.0 mg, 0.0894 mmol) in TFA (1.5 mL) was allowed to stand at ambient temperatures for 45 min and concentrated to a yellow oil. This oil was dissolved in anhydrous DMF (2.0 mL) and made basic to pH paper with TEA. In a separate flask, the
30 product of Part B, above (69.3 mg, 0.0547 mmol) was dissolved in anhydrous DMF (2.0 mL) and pre-activated by treatment with TEA (0.015 mL, 0.104 mmol) and HBTU (32.6 mg, 0.0859 mmol). After 10 min this solution was added to the DMF solution from the TFA deprotection reaction,
35 and the combined solutions were stirred at ambient temperatures for 30 min. The DMF was removed under vacuum and the resulting oil was dissolved in 60% ACN and purified by preparative HPLC on a Vydac C-18 column (22 x 250 mm) using a 1.54%/min gradient of 18 to 72% ACN

containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 29.9 min was adjusted to pH 8 with saturated NaHCO₃ and concentrated to remove the ACN. The remaining aqueous mixture was extracted with EtOAc (2 x 40 mL). The combined EtOAc extracts were washed with saturated NaCl (40 mL), dried (MgSO₄), and concentrated to give the title compound as a colorless solid (56.4 mg, 63%). MS: m/e 1644.8 [M+H]⁺; High Resolution MS: Calcd for C₈₇H₁₁₀N₁₁O₁₉S [M+H]⁺: 1644.7700, Found: 1644.771.



Part D - Methyl 3-((1-(3-(3-(N-(3-(2-(2-(N-(L-Asp(O-t-Bu)-L-Asp(O-t-Bu))3-aminopropoxy)-ethoxy)ethoxy)propyl)carbamoyl)propanoylamino)propyl-4-oxo-7-((1-(triphenylmethyl)imidazole-2-yl)amino)methyl)(3-hydroquinolyl))carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoate


The product of Part E, above (55.0 mg, 0.0335 mmol) was hydrogenolyzed over 10% Pd/C (25 mg) in MeOH (15 mL) at 40 psi for 3.5 h. The catalyst was removed by filtration through filter aid and the filtrate was concentrated to give the title compound as a pale yellow oil (41.8 mg, 83%). MS: m/e 1510.8 [M+H]⁺.



Part E - DOTA-tri-t-butyl Ester Conjugate of Methyl 3-
 ((1-(3-(3-(N-(3-(2-(2-(N-(L-Asp(O-t-Bu)-L-Asp(O-t-Bu))3-
 5 aminopropoxy)ethoxy)ethoxy)propyl)carbamoyl)-
 propanoylamino)propyl-4-oxo-7-(((1-(triphenylmethyl)-
 imidazole-2-yl)amino)methyl)(3-
 hydroquinolyl))carbonylamino)-2-(((2,4,6-
 trimethylphenyl)sulfonyl)amino)propanoate


10 A solution of the product of Part D, above (41.8 mg,
 0.0277 mmol), the product of Example 2, Part B, 39.9 mg,
 0.0436 mmol), TEA (0.023 mL, 0.166 mmol), and HBTU (15.6
 mg, 0.0411 mmol) in anhydrous DMF (3.0 mL) was allowed to
 stand at ambient temperatures under a nitrogen atmosphere
 15 for 20 h. The DMF was removed under vacuum and the
 resulting oil was dissolved in 60% ACN and purified by
 preparative HPLC on a Vydac C-18 column (22 x 250 mm)
 using a 2.4%/min gradient of 18 to 90% ACN containing
 0.1% TFA at a flow rate of 20 mL/min. The main product
 20 peak eluting at 21.2 min was collected and lyophilized to
 give the title compound as a colorless powder (24.8 mg,
 43%). MS: m/e 2066.3 [M+H], 1033.6 [M+2H]; High
 Resolution MS: Calcd for C₁₀₇H₁₅₄N₁₅O₂₄S [M+H]: 2065.1011,
 Found: 2065.1030.

25



Part F - DOTA Conjugate of 3-((1-(3-(3-(N-(3-(2-(2-(N-(L-
 Asp-L-Asp)3-aminopropoxy)ethoxy)ethoxy)propyl)carbamoyl)-
 5 propanoylamino)propyl-7-((imidazole-2-ylamino)methyl)-4-
 oxo(3-hydroquinolyl))carbonylamino)-2-((2,4,6-
 trimethylphenyl)sulfonyl)amino)propanoic Acid
 Bis(trifluoroacetate Salt

A mixture of the product of Part G, above (18.8 mg.
 10 0.0091 mmol), water (0.150 mL), 3 N LiOH (0.015 mL), and
 peroxide-free THF (1.5 mL) was stirred at ambient
 temperatures for 3 h. The THF was removed under vacuum
 and the aqueous mixture was diluted with water (40 mL)
 and adjusted to pH 7 with 0.1 N HCl. The mixture was
 15 extracted with DCM (2 x 30 mL) and the combined extracts
 were concentrated to give a yellow oil. This oil was
 dissolved in TFA (1.0 mL) along with Et₃SiH (0.030 mL,
 0.184 mmol) and heated at 40 °C under a nitrogen
 atmosphere for 48 h. The solution was concentrated and
 20 the resulting oil was dissolved in water and purified by
 preparative HPLC on a Vydac C-18 column (22 x 250 mm)
 using 0.1% TFA in water for 5 min followed by a 2.52%/min
 gradient of 0 to 63% ACN containing 0.1% TFA at a flow
 rate of 20 mL/min. The main product peak eluting at 19.9
 25 min was collected and lyophilized to give the title
 compound as a colorless powder (1.5 mg, 9.4%). MS: m/e
 1528.9 [M+2H], 765.1 [M+2H], 510.7 [M+3H].



Example 9

- 5 DOTA/2-Amino-N,N'-bis(3-(2-(2-(3-(3-(N-(3-(3-(N-(3-carboxy-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)ethyl)-carbamoyl)-7-((imidazole-2-ylamino)methyl)4-oxohydroquinolyl)-propyl)carbamoyl)propanoylamino)propoxy)ethoxy)ethoxy)-propyl)pentane-1,5-diamide Tris(trifluoroacetate) Salt Conjugate


- Part A - DOTA-tri-t-butyl Ester/2-Amino-N,N'-bis(3-(2-(2-(3-(3-(N-(3-(3-(N-(3-carboxy-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)ethyl)carbamoyl)-7-((imidazole-2-ylamino)methyl)4-oxohydroquinolyl)propyl)carbamoyl)-propanoylamino)propoxy)ethoxy)ethoxy)propyl)pentane-1,5-diamide Hexakis(trifluoroacetate) Salt Conjugate

- A solution of the product of Example 2, Part B, HBTU, and DIEA in anhydrous DMF is stirred at ambient temperatures under nitrogen for 15 min and treated with the product of Example 7, Part B. The resulting solution is stirred an additional 18 h and the DMF is removed under vacuum. The resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient. The product fraction is lyophilized to give the title compound.

- Part B - DOTA/2-Amino-N,N'-bis(3-(2-(2-(3-(3-(N-(3-(3-(N-(3-carboxy-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)ethyl)carbamoyl)-7-((imidazole-2-ylamino)methyl)4-oxohydroquinolyl)propyl)carbamoyl)-

propanoylamino)propoxy)ethoxy)ethoxy)propyl)pentane-1,5-diamide Tris(trifluoroacetate) Salt Conjugate

The product of Part B, above, is dissolved in degassed TFA, treated with triethylsilane, and heated at 50 °C under nitrogen for 1 h. The solution is concentrated under vacuum and the resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient. The product fraction is lyophilized to give the title compound.



Example 10

DOTA/2-(((4-(3-(N-(3-(2-(2-(3-(2-Amino-3-sulfopropyl)-propoxy)ethoxy)ethoxy)propyl)carbamoyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)propanoic Acid Trifluoroacetate Salt Conjugate

Part A - 2-(((4-(3-(N-(3-(2-(2-(3-(2-((tert-Butoxy)-carbonylamino)-3-sulfopropyl)propoxy)ethoxy)ethoxy)-propyl)carbamoyl)propoxy)-2,6-

dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)propanoic Acid

The product of Example 3, Part I is dissolved in anhydrous DMF and treated with the N-hydroxysuccinimide ester of Boc-cysteic acid (as described in *Liebigs Ann.*

Chem. 1979, 776-783) and DIEA. The solution is stirred at ambient temperatures under nitrogen for 18 h, and the DMF is removed under vacuum. The resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient. The product fraction is lyophilized to give the title compound.

Part B - DOTA-tri-t-butyl Ester/2-(((4-(3-(N-(3-(2-(2-(3-(2-Amino-3-sulfopropyl)propoxy)ethoxy)ethoxy)propyl)-carbamoyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic Acid
Tetrakis(trifluoroacetate) Salt Conjugate

The product of Part A, above, is dissolved in degassed TFA and stirred at ambient temperatures for 15 min. The solution is concentrated under vacuum, and the resulting residue is dissolved in 50% ACN and lyophilized to remove the last traces of TFA.


In a separate flask, a solution of the product of Example 2, Part B and DIEA in anhydrous DMF are treated with HBTU and allowed to react 15 min at ambient temperatures under nitrogen. The deprotected product from above is added to this solution and stirring is continued at ambient temperatures under nitrogen for 18 h. The DMF is removed under vacuum and the resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient. The product fraction is lyophilized to give the title compound.

Part C - DOTA/2-(((4-(3-(N-(3-(2-(2-(3-(2-Amino-3-sulfopropyl)-propoxy)ethoxy)ethoxy)propyl)carbamoyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)propanoic Acid Trifluoroacetate Salt Conjugate

The product of Part B, above, and Et₃SiH are dissolved in degassed TFA and heated at 50 °C under nitrogen for 1 h. The solution is concentrated and the

resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient. The product fraction is lyophilized to give the title compound.

5




Example 11

DOTA/2-(((4-(3-(N-(3-(2-(2-(3-(2-Amino-3-(4-(
(phosphonooxy)-
phenyl)propanoylamino)propoxy)ethoxy)ethoxy)propyl)carbam
oyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-
((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-
hydroquinolyl))carbonylamino)propanoic Acid
Trifluoroacetate Salt Conjugate

The title compound is prepared by the same procedure described for Example 10 by substituting Boc-Tyr(PO₃H₂)-OSu for Boc-Cys(O₃H)-OSu.

20




25

Example 12

DOTA/2-(((4-(3-(N-(3-(2-(2-(3-(2-Amino-3-(4-(sulfooxy)-
 phenyl)propanoylamino)propoxy)ethoxy)ethoxy)propyl)carbam
 oyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-
 ((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-
 5 hydroquinolyl))carbonylamino)propanoic Acid
 Trifluoroacetate Salt Conjugate

The title compound is prepared by the same procedure
 described for Example 10 by substituting Boc-Tyr(SO₃H)-
 10 OSu for Boc-Cys(O₃H)-OSu.



Example 13

DOTA/2-(((4-(3-(N-(3-(2-(2-(3-(2-Amino-4-(N-(ethyl-3,6-O-
 disulfo- β -D-galactopyranosyl) carbamoyl) butanoylamino)-
 propoxy)ethoxy)ethoxy)propyl) carbamoyl)propoxy)-2,6-
 5 dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-
 ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-
 carbonylamino)propanoic Acid Conjugate

Part A - Preparation of Boc-Glu(aminoethyl-3,6-O-disulfo-
 10 β -D-galactopyranosyl)-OSu


A solution of Boc-Glu-OMe, aminoethyl-3,6-O-disulfo-
 β -D-galactopyranoside (as described in *Tet. Lett.* 1997,
 53, 11937-11952), DIEA, and HBTU in anhydrous DMF is
 stirred at ambient temperatures under nitrogen for 18 h.
 15 The DMF is removed under vacuum and the resulting residue
 is hydrolyzed using aqueous NaOH. The reaction solution
 is adjusted to pH 7 and purified by preparative anion
 exchange chromatography using a resin such as DEAE
 Cellulose and a Et₃NH₂CO₃ gradient. The product fraction
 20 is treated with a cation exchange resin, sodium form, to
 give the intermediate carboxylic acid as the sodium salt.

The above compound, N-hydroxysuccinimide, and DCC
 are dissolved in anhydrous DMF and stirred at ambient
 temperatures under nitrogen for 18 h. The DMF is removed
 25 under vacuum and the resulting residue is purified by
 preparative anion exchange chromatography as above to
 give the title compound as the triethylammonium salt.

Part B - DOTA/2-(((4-(3-(N-(3-(2-(2-(3-(2-Amino-4-(N-
 30 (ethyl-3,6-O-disulfo- β -D-
 galactopyranosyl) carbamoyl) butanoylamino)-
 propoxy)ethoxy)ethoxy)propyl) carbamoyl)propoxy)-2,6-
 dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-
 ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-
 35 carbonylamino)propanoic Acid Conjugate

The title compound is prepared by the same procedure
 described for Example 10 by substituting Boc-

Glu(aminoethyl-3,6-O-disulfo- β -D-galactopyranosyl)-OSu
for Boc-Cys(O₃H)-OSu.



5

Example 14

DOTA/2-(((4-(3-(N-(3-(2-(2-(3-(2-Amino-4-(N-(6-deoxy- β -
cyclodextryl) carbamoyl) butanoylamino)-
propoxy) ethoxy) ethoxy) propyl) carbamoyl) propoxy)-2,6-
10 dimethylphenyl) sulfonyl) amino)-3-((7-((imidazol-2-
ylamino) methyl)-1-methyl-4-oxo(3-hydroquinolyl))-
carbonylamino) propanoic Acid Bis(trifluoroacetate) Salt
Conjugate

15 Part A - Preparation of Boc-Glu(6-amino-6-deoxy- β -
cyclodextryl)-OMe

A solution of Boc-Glu-OMe, 6-amino-6-deoxy- β -
cyclodextrin (as described in *J. Org. Chem.* 1996, 61,
903-908), DIEA, and HBTU in anhydrous DMF is stirred at
20 ambient temperatures under nitrogen for 18 h. The DMF is
removed under vacuum and the resulting residue is
purified by preparative HPLC on a C18 column using a
water:ACN:0.1% TFA gradient. The product fraction is
lyophilized to give the title compound.

25


Part B - Preparation of Boc-Glu(6-amino-6-deoxy- β -
cyclodextryl)-OSu

The product of Part A, above, is hydrolyzed by
stirring in a mixture of LiOH, THF, and water at ambient
30 temperatures under nitrogen for 4 h. The THF is removed

under vacuum and the resulting mixture is diluted with water and adjusted to pH 3 using 0.1 N HCl. The mixture is extracted with EtOAc, and the combined extracts are dried (MgSO₄) and concentrated. The resulting material
 5 is dissolved in anhydrous DMF along with N-hydroxysuccinimide, and DCC, and stirred at ambient temperatures under nitrogen for 18 h. The DMF is removed under vacuum and the resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1%
 10 TFA gradient. The product fraction is lyophilized to give the title compound.

Part C - DOTA/2-(((4-(3-(N-(3-(2-(2-(3-(2-Amino-4-(N-(6-deoxy-β-cyclodextryl) carbamoyl) butanoylamino)-
 15 propoxy)ethoxy)ethoxy)propyl) carbamoyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)propanoic Acid Bis(trifluoroacetate) Salt Conjugate

20 The title compound is prepared by the same procedure described for Example 10 by substituting Boc-Glu(6-amino-6-deoxy-β-cyclodextryl)-OSu for Boc-Cys(O₃H)-OSu.



25

Example 15

DOTA/2-(((4-(3-(N-(3-(2-(2-(3-(2-Amino-4-(N-(ω-methoxypolyethylene(5,000)glycoxyethyl) carbamoyl)-
 30 butanoylamino)propoxy)ethoxy)ethoxy)propyl) carbamoyl)-propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-

((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic Acid
Bis(trifluoroacetate) Salt Conjugate

5 Part A - Preparation of Boc-Glu(amino- ω -methoxypolyethylene glycol)-OMe

A solution of Boc-Glu-OMe, amino- ω -methoxypolyethylene glycol, (MW = 5,000), DIEA, and HBTU in anhydrous DMF is stirred at ambient temperatures under
10 nitrogen for 18 h. The DMF is removed under vacuum and the resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient. The product fraction is lyophilized to give the title compound.


15

Part B - Preparation of Boc-Glu(amino- ω -methoxypolyethylene glycol)-OSu

The product of Part A, above, is hydrolyzed by stirring in a mixture of LiOH, THF, and water at ambient
20 temperatures under nitrogen for 4 h. The THF is removed under vacuum and the resulting solution is adjusted to pH 7 using 0.1 N HCl. The solution is desalted using a Sephadex PD-10 desalting column and the product eluant is lyophilized. The resulting material is dissolved in
25 anhydrous DMF along with N-hydroxysuccinimide, and DCC, and stirred at ambient temperatures under nitrogen for 18 h. The DMF is removed under vacuum and the resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient. The product
30 fraction is lyophilized to give the title compound.

Part C - DOTA/2-(((4-(3-(N-(3-(2-(2-(3-(2-Amino-4-(N-(ω -methoxypolyethylene(5,000)glycoxyethyl)carbamoyl)-butanoylamino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)-propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic Acid
35 Bis(trifluoroacetate) Salt Conjugate

The title compound is prepared by the same procedure described for Example 10 by substituting Boc-Glu(amino- ω -methoxypolyethylene glycol)-OSu for Boc-Cys(O₃H)-OSu.



5


Example 16

2-(((4-(3-(N-(3-(2-(2-(3-(2-(1,4,7,10-Tetraaza-4,7,10-tris(carboxymethyl)cyclododecylacetyl amino)-6-aminohexanoylamino)propoxy)ethoxy)ethoxy)-propyl)carbamoyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)propanoic Acid Tris(trifluoroacetate) Salt

15

The title compound is prepared by the same procedure described for Example 10 by substituting Boc-Lys(Cbz)-OSu for Boc-Cys(O₃H)-OSu.

20




Example 17

2-(((4-(3-(N-(3-(2-(2-(3-(2-(1,4,7,10-Tetraaza-4,7,10-
 tris(carboxymethyl)cyclododecylacetyl amino)-6-(2-
 (bis(phosphonomethyl) amino) acetyl amino) hexanoyl amino)-
 propoxy) ethoxy) ethoxy) propyl) carbamoyl) propoxy)-2,6-
 5 dimethylphenyl) sulfonyl) amino)-3-((7-((imidazol-2-
 yl amino) methyl)-1-methyl-4-oxo(3-hydroquinolyl))-
 carbonyl amino) propanoic Acid Conjugate

A solution of bis(phosphonomethyl)glycine, DIEA, and
 10 HBTU in anhydrous DMF is stirred at ambient temperatures
 under nitrogen for 15 min, and treated with the product
 of Example 16. Stirring is continued for 18 h and the
 DMF is removed under vacuum. The resulting residue is
 purified by ion exchange chromatography.


15



Example 18

20 2-(((4-(3-(N-(3-(2-(2-(3-(2-(2-(2-((2-(2-
 (bis(carboxymethyl)-
 amino) ethyl) (carboxymethyl) amino) ethyl) (carboxymethyl) ami
 no) acetyl amino)-3-sulfopropyl) propoxy) ethoxy)-
 ethoxy) propyl) carbamoyl) propoxy)-2,6-dimethylphenyl)-
 25 sulfonyl) amino)-3-((7-((imidazol-2-yl amino) methyl)-1-
 methyl-4-oxo(3-hydroquinolyl)) carbonyl amino) propanoic
 Acid

The product of Example 10, Part A is dissolved in degassed TFA and stirred at ambient temperatures for 15 min. The solution is concentrated under vacuum, and the resulting residue is dissolved in 50% ACN and lyophilized to remove the last traces of TFA. The material is dissolved in anhydrous DMF along with DIEA and diethylenetriaminepentaacetic dianhydride. The resulting solution is stirred at ambient temperatures under nitrogen for 18 h. The DMF is removed under vacuum and the resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient. The product fraction is lyophilized to give the title compound.



15

The following procedure describe the synthesis of radiopharmaceuticals of the present invention of the formula $^{99m}\text{Tc}(\text{VnA})(\text{tricine})(\text{phosphine})$, in which (VnA) represents a vitronectin receptor antagonist compound of the present invention bonded to the Tc through a diazenido ($-\text{N}=\text{N}-$) or hydrazido ($=\text{N}-\text{NH}-$) moiety. The diazenido or hydrazido moiety results from the reaction of the hydrazinonicotinamido group, present either as the free hydrazine or protected as a hydrazone, with the Tc-99m. The other two ligands in the Tc coordination sphere are tricine and a phosphine.

30

Examples 19 - 23

Synthesis of Complexes [$^{99m}\text{Tc}(\text{HYNIC}-\text{VnA})(\text{tricine})(\text{TPPTS})$].

To a lyophilized vial containing 4.84 mg TPPTS, 6.3 mg tricine, 40 mg mannitol, succinic acid buffer, pH 4.8,

and 0.1% Pluronic F-64 surfactant, was added 1.1 mL sterile water for injection, 0.2 mL (20 µg) of the appropriate HYNIC-conjugated vitronectin antagonist (VnA) in deionized water or 50% aqueous ethanol, and 0.2 mL of ^{99m}TcO₄⁻ (50±5 mCi) in saline. The reconstituted kit was heated in a 100°C water bath for 15 minutes, and was allowed to cool 10 minutes at room temperature. A sample of the reaction mixture was analyzed by HPLC. The RCP results are listed in the Table 1.

HPLC Method

Column: Zorbax C18 , 25 cm x 4.6 mm

Flow rate : 1.0 mL/min

15 Solvent A: 10 mM sodium phosphate buffer, pH 6.0

Solvent B : 100 % CH₃CN

Gradient A (Exs. 19, 20, 21)

t (min)	0	20	21	30	31	40
% Solvent B	0	25	75	75	0	0

20

Gradient B (Ex. 22)

t (min)	0	20	30	31	40
% Solvent B	0	50	50	0	0

25 Gradient C (Ex. 23)

t (min)	0	20	21	30	31	40
% Solvent B	10	30	75	75	0	0

Table 1. Analytical and Yield Data for

30 ^{99m}Tc(VnA) (tricine) (TPPTS) Complexes

Example No.	Reagent No.	Ret. Time (min)	% Yield
19	1	8.8	73
20	3	17.2	81
21	4	17.6	68
22	6	11.7	79
23	7	16.4	52

Example 24

Synthesis of the In-111 Complex of 3-((7-((Imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((4-(4-((3-(2-(2-(3-(2-(1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)cyclododecyl)acetylamino)-propoxy)ethoxy)ethoxy)propyl)amino)sulfonyl)-phenyl)phenyl)sulfonyl)amino)propanoic Acid

To a lead shielded and crimped autosampler vial was added 35 µg of the conjugate of Example 2 and 1.0 mg gentisic acid, sodium salt dissolved in 70 µL ammonium acetate buffer (0.4 M, pH 4.7) followed by the addition of 2 mCi, 20 µL In-111 in 0.05 N HCl (specific activity: 17 µg/mCi). The reaction mixture was heated at 70 - 80 °C for 60 min and analyzed by HPLC and ITLC. The complex was formed in 93% yield and had a retention time of 19.6 min.

20 HPLC Method

Column: Zorbax Rx C18, 25 cm x 4.6 mm
 Column Temperature: Ambient
 Flow: 1.0 mL/min
 Solvent A: 10 % Acetonitrile/0.1%TFA/H₂O

25 Solvent B: Acetonitrile

Detector: Sodium iodide (NaI) radiometric probe
 Gradient

t (min)	0	25	26	35	36	45
%B	10	20	60	60	10	10

30

Examples 25 - 26

Synthesis of ¹⁷⁷Lu and ⁹⁰Y Complexes of 3-((7-((Imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((4-(4-((3-(2-(2-(3-(2-(1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)cyclododecyl)acetylamino)-propoxy)ethoxy)ethoxy)propyl)amino)sulfonyl)-phenyl)phenyl)sulfonyl)amino)propanoic Acid.

To a clean sealed 5 mL vial was added 0.3 mL of a solution of the conjugate of Example 2 (200 µg/mL in 0.5 M ammonium acetate buffer, pH 6.9), followed by 0.05 mL of gentisic acid (sodium salt, 10 mg/mL in 0.5 M ammonium acetate buffer, pH 6.9) solution, 0.3 mL of 0.5 M ammonium acetate buffer (pH 6.9), and 0.010 mL of $^{177}\text{LuCl}_3$ or $^{90}\text{YCl}_3$ solution (1000 mCi/mL for $^{177}\text{LuCl}_3$ and 500 mCi/mL for $^{90}\text{YCl}_3$) in 0.05 N HCl. The resulting mixture was heated at 100 °C for 30 min. After cooling to room temperature, a sample of the resulting solution was analyzed by radio-HPLC and ITLC. The radiolabeling yields were = 90% (after correction for small amount of colloid) for both complex, and the retention time was 19.2 min.

15

HPLC Method

Column: Zorbax C18 , 25 cm x 4.6 mm

Flow rate : 1.0 mL/min

Solvent A: 0.1% TFA aqueous solution

20 Solvent B : 100 % CH_3CN

t (min)	0	20	25	30	31	40
% Solvent B	10	25	60	60	10	10

The instant thin layer chromatography (ITLC) method used Gelman Sciences silica-gel strips and a 1:1 mixture of acetone and saline as eluant.

25

Example 27

Synthesis of ^{177}Lu Complex of the DOTA Conjugate of 3-((1-(3-(3-(N-(3-(2-(2-(N-(L-Asp-L-Asp)3-aminopropoxy)-ethoxy)ethoxy)propyl)carbamoyl)propanoylamino)propyl-7-((imidazole-2-ylamino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic Acid.

To a clean sealed 5 mL vial was added 0.5 mL of a solution of the conjugate of Example 8 (200 µg/mL in 0.5 M ammonium acetate buffer, pH 6.9), followed by 0.05 mL of gentisic acid (sodium salt, 10 mg/mL in 0.5 M ammonium

35

acetate buffer, pH 6.9) solution, 0.25 mL of 0.5 M ammonium acetate buffer (pH 6.9), and 0.05 mL of $^{177}\text{LuCl}_3$ solution (200 mCi/mL) in 0.05 N HCl. The resulting mixture was heated at 100 °C for 30 min. After cooling to room temperature, a sample of the resulting solution was analyzed by radio-HPLC and ITLC. The radiolabeling yield was 75% (after correction for colloid), and the retention time was 20 min.

10 HPLC Method

Column: Zorbax C18 , 25 cm x 4.6 mm

Flow rate : 1.0 mL/min

Solvent A: 10 mM phosphate buffer, pH = 6

Solvent B : 100 % CH₃CN

15	t (min)	0	20	25	30	31	40
	% Solvent B	0	20	50	50	0	0

Example 28

Synthesis of the Gadolinium Complex of 2-(((4-(3-(N-(3-(2-(2-(3-(2-(2-((2-((2-(bis(carboxymethyl)-amino)ethyl)(carboxymethyl)amino)ethyl)(carboxymethyl)amino)acetylamino)-3-sulfopropyl)propoxy)ethoxy)-ethoxy)propyl)carbamoyl)propoxy)-2,6-dimethylphenyl)-sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic Acid


The gadolinium complex of the conjugate of Example 18 is prepared according to the following procedure. 3-3.5 mg of the conjugate is dissolved in 2 mL 1 M ammonium acetate buffer at pH 7.0 , and one equivalent Gd(NO₃)₃ solution (0.02 M in water) is added to it. The reaction mixture is allowed to stay at room temperature for 3-5 hours and the product is isolated by HPLC. The fraction containing the complex is lyophilized and dissolved in 1 mL H₂O. The identity of the complex is confirmed by mass spectroscopy.

Example 29


Synthesis of (2S)-2-[(2,6-Dimethyl-4-[3-(N-{2-[3-sulfo-2-(3-sulfo-2-{2-[1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)

5 cyclododecyl]acetyl amino}propyl)propyl]ethyl]carbamoyl)-
propoxy]phenyl)sulfonyl]amino]-3-({7-[(imidazol-2-
ylamino)methyl]-1-methyl-4-oxo(3-hydroquinolyl)}-
carbonylamino)propanoic Acid Trifluoroacetate Salt

10



Part A - Preparation of Methyl (2S)-3-[(tert-Butoxy)-
15 carbonylamino]-2-[(2,6-dimethyl-4-[3-(N-{2-
[(phenylmethoxy)carbonylamino]ethyl)carbonyl]propoxy]-
phenyl)sulfonyl]amino]propanoate




20

A solution of the product of Example 3, Part D (369 mg, 0.756 mmol), DIEA (0.52 mL, 3.0 mmol), and HBTU (315 mg, 0.832 mmol) in anhydrous DMF (14 mL) was stirred at ambient temperatures under nitrogen for 5 min, and
25 treated with benzyl N-(2-aminoethyl)carbamate hydrochloride (192 mg, 0.832 mmol), and stirred an additional 1 h. The DMF was removed under vacuum, and


the oily residue was taken up in EtOAc (150 mL), washed consecutively with 0.1 N HCl (40 mL), water (40 mL), and saturated NaCl (40 mL), dried (MgSO₄), and concentrated to give a colorless viscous oil. Flash chromatography on a 3 x 16 cm silica gel column (EtOAc) gave the title compound as a colorless viscous oil (450 mg, 89.6%). ¹H NMR (CDCl₃): δ 7.34-7.27 (m, 5H), 6.58 (s, 2H), 6.31 (bs, 1H), 5.86 (bs, 1H), 5.36 (bs, 1H), 5.14-5.03 (m, 3H), 3.96 (t, J = 6.0 Hz, 2H), 3.88-3.83 (m, 1H), 3.56 (s, 3H), 3.47-3.25 (m, 6H), 2.59 (s, 6H), 2.31 (t, J = 6.9 Hz, 2H), 2.05 (p, J = 6.6 Hz, 2H), 1.39 (s, 9H); ¹³C NMR (CDCl₃): δ 172.9, 170.5, 160.6, 157.3, 155.9, 141.8, 136.3, 128.5, 128.2, 128.0, 116.6, 79.9, 66.9, 55.5, 52.8, 43.1, 40.9, 40.3, 32.4, 28.2, 24.9, 23.3; MS: m/e 665.4 [M+H]; 687.3 [M+Na]; High Resolution MS: Calcd for C₃₁H₄₅N₄O₁₀S [M+H]: 665.2856, Found: 665.2883.

Part B - Preparation of Methyl (2S)-3-Amino-2-[(2,6-dimethyl-4-[3-(N-{2-[(phenylmethoxy)carbonylamino]ethyl} carbamoyl)propoxy]phenyl)sulfonyl]amino]propanoate Trifluoroacetate Salt




The product of Part A, above (420 mg, 0.632 mmol) was dissolved in 25/75 DCM/TFA (20 mL) and allowed to stand at ambient temperatures under nitrogen for 10 min. The solution was concentrated, and the resulting viscous oil was dissolved in 50% ACN and lyophilized to give the title compound as a colorless solid (437 mg, 102%). MS: m/e 565.3 [M+H].

Part C - Preparation of Methyl (2S)-2-[(2,6-Dimethyl-4-
 [3-(N-{2-[(phenylmethoxy)carbonylamino]ethyl}carbamoyl)-
 propoxy]phenyl)sulfonyl]amino]-3-[[1-methyl-4-oxo-7-((1-
 (triphenylmethyl)imidazol-2-yl)amino)methyl](3-
 5 hydroquinolyl)]carbonylamino]propanoate




A solution of 1-methyl-4-oxo-7-(((1-
 10 (triphenylmethyl)imidazol-2-
 yl)amino)methyl)hydroquinoline-3-carboxylic acid (702 mg,
 1.30 mmol), DIEA (0.678 mL, 3.90 mmol), and HBTU (542 mg,
 1.43 mmol) in anhydrous DMF (60 mL) was stirred at
 ambient temperatures under nitrogen for 10 min, and
 15 treated with the product of Step B, above (881 mg, 1.30
 mmol). After 75 min the DMF was removed under vacuum and
 the resulting oil was purified by HPLC on a Vydac C-18
 column (50 x 250 mm) using a 1.24%/min gradient of 18 to
 67.5% ACN containing 0.1% TFA at a flow rate of 80
 20 mL/min. A peak eluting at 18.9 min was lyophilized to
 give unreacted 1-methyl-4-oxo-7-(((1-(triphenylmethyl)-
 imidazol-2-yl)amino)methyl)hydroquinoline-3-carboxylic
 acid (308 mg). The main product peak eluting at 23.7 min
 was lyophilized to give the title compound as a colorless
 25 solid (890 mg, 63.0%). ¹H NMR (CDCl₃/D₂O): δ 8.50 (s, 1H),
 8.18 (d, J = 8.3 Hz, 1H), 7.70 (s, 1H), 7.51-7.25 (m,
 15H), 7.25-7.12 (m, 5H), 6.97 (s, 1H), 6.58 (d, J = 2.3
 Hz, 1H), 6.34 (s, 2H), 6.32 (d, J = 8.5 Hz, 1H), 5.09 (s,
 2H), 4.65 (s, 2H), 4.29-4.23 (m, 1H), 3.88 (s, 3H), 3.80-
 30 3.50 (m, 7H), 3.41-3.28 (m, 4H), 2.61 (s, 6H), 2.26-2.11
 (m, 2H), 1.92-1.76 (m, 2H); MS: m/e 1087.4 [M+H]⁺; 845.3
 [M+H-Tr]⁺; High Resolution MS: Calcd for C₆₀H₆₃N₉O₁₀S [M+H]⁺:
 1087.4388; found: 1087.440.

- Part D - Preparation of Methyl (2S)-2-{{{(4-{3-[N-(2-Aminoethyl)carbamoyl]propoxy}-2,6-dimethylphenyl)-sulfonyl}amino)-3-{{[1-methyl-4-oxo-7-{{[1-(triphenylmethyl)imidazol-2-yl]amino)methyl}(3-hydroquinolyl)]carbonylamino}propanoate



- 10 Hydrogenolysis of the product of Part C, above (468 mg, 0.431 mmol) was accomplished in MeOH (100 mL) over 10% Pd/C (95 mg) at 60 psi for 1 h. The catalyst was removed by filtration through Celite® and the filtrate was concentrated to give the title compound as a pale
15 amber oil (405 mg, 98.7%). MS: m/e 953.3 [M+H], 711.3 [M+H-Trityl].

- Part E - Preparation of (2R)-N-{2-[4-(4-{{{(1S)-1-(Methoxycarbonyl)-2-{{[1-methyl-4-oxo-7-{{[1-(triphenylmethyl)imidazol-2-yl]amino)methyl}(3-hydroquinolyl)]carbonylamino}ethyl)amino]sulfonyl}-3,5-dimethylphenoxy)butanoylamino]ethyl}-2-[(tert-butoxy)carbonylamino]propanesulfonic Acid




25

A solution of the product of Part E, above (405 mg, 0.425 mmol), the p-nitrophenyl ester of Boc-L-cysteic

acid (425 mg, 1.03 mmol), and DIEA (0.435 mL, 2.55 mmol) in anhydrous DMF (20 mL) was stirred at ambient temperatures under nitrogen for 3 h. The DMF was removed under vacuum and the resulting oil was purified by HPLC
 5 on a Vydac C-18 column (50 x 250 mm) using a 1.12%/min gradient of 9 to 54% ACN containing 0.1% TFA at a flow rate of 80 mL/min. The main product peak eluting at 37.3 min was lyophilized to give the title compound as a colorless solid (410 mg, 80.2%). MS: m/e 1204.4 [M+H],
 10 962.3 [M+H-Trt].

Part F - Preparation of (2R)-N-(2-[4-(4-(((1S)-1-(Methoxycarbonyl)-2-([1-methyl-4-oxo-7-([1-(triphenylmethyl)imidazol-2-yl]amino)methyl)(3-
 15 hydroquinolyl)]carbonylamino)ethyl)amino)sulfonyl]-3,5-dimethylphenoxy)butanoylamino]ethyl)-2-aminopropanesulfonic Acid




20

The product of Part E, above (410 mg, 0.341 mmol) was dissolved in 50/50 TFA/DCM (20 mL) and allowed to react at ambient temperatures for 10 min. The solution was concentrated and the resulting amber oil was
 25 dissolved in 50% ACN (50 mL) and lyophilized to give the title compound as a colorless solid (371 mg, 98.6%). MS: m/e 1104.4 [M+H], 862.3 [M+H-Trt]; High Resolution MS: Calcd for C₅₅H₆₂N₉O₁₂S₂ [M+H]: 1104.3959; Found: 1104.393.

30 Part G - Preparation of (2R)-N-[(1R)-1-(N-(2-[4-(4-(((1S)-1-(Methoxycarbonyl)-2-([1-methyl-4-oxo-7-([1-(triphenylmethyl)imidazol-2-yl]amino)methyl)(3-hydroquinolyl)]carbonylamino)ethyl)amino)sulfonyl]-3,5-

dimethylphenoxy)butanoylamino]ethyl)carbamoyl)-2-sulfoethyl]-2-[(tert-butoxy)carbonylamino]propanesulfonic Acid




5

A solution of the product of Part F, above (110 mg, 0.100 mmol), the p-nitrophenyl ester of Boc-L-cysteic acid (82.4 mg, 0.200 mmol), and DIEA (0.104 mL, 0.600 mmol) in anhydrous DMF (5.0 mL) was stirred at ambient temperatures under nitrogen for 48 h. The DMF was removed under vacuum and the resulting amber oil was purified by HPLC on a Vydac C-18 column (50 x 250 mm) using a 1.12%/min gradient of 9 to 54% ACN containing 0.1% TFA at a flow rate of 80 mL/min. The main product peak eluting at 37.0 min was lyophilized to give the title compound as a colorless solid (96.0 mg, 70.9%). MS: m/e 1355.3 [M+H], 1113.3 [M-Trt+H], 1013.2 [M-Trt-Boc+H].

20


Part H - Preparation of (2R)-N-[(1R)-1-(N-{2-[4-(4-(((1S)-1-(Methoxycarbonyl)-2-[[1-methyl-4-oxo-7-([1-(triphenylmethyl)imidazol-2-yl]amino)methyl)(3-hydroquinolyl)]carbonylamino)ethyl)amino]sulfonyl)-3,5-dimethylphenoxy]butanoylamino]ethyl)carbamoyl)-2-sulfoethyl]-2-aminopropanesulfonic Acid

25



The product of Part G, above (21 mg, 0.0155 mmol) was dissolved in 50/50 TFA/DCM (5.0 mL) and allowed to react at ambient temperatures for 10 min. The solution was concentrated and the residue was taken up in 50% ACN (15 mL) and lyophilized to give the title compound as a colorless solid (18.7 mg, 96.2%). MS: m/e 1255.3 [M+H], 1013.2 [M+H-Trityl]; High Resolution MS: Calcd for C₅₈H₆₇N₁₀O₁₆S₃ [M+H]: 1255.3899; Found: 1255.391.

Part I - Preparation of (2R)-N-[(1R)-1-(N-(2-[4-(4-(((1S)-1-(Methoxycarbonyl)-2-[[1-methyl-4-oxo-7-([1-(triphenylmethyl)imidazol-2-yl]amino)methyl)(3-hydroquinolyl)]carbonylamino)ethyl)amino)sulfonyl]-3,5-dimethylphenoxy)butanoylamino]ethyl)carbamoyl)-2-sulfoethyl]-2-(2-[1,4,7,10-tetraaza-4,7,10-tris((tert-butoxycarbonyl)methyl)cyclododecyl)acetylamino)propanesulfonic Acid



A solution of 2-(1,4,7,10-tetraaza-4,7,10-tris(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)acetic acid (30.0 mg, 0.0327 mmol) (as described in DM-7003), DIEA (0.034 mL, 0.196 mmol), and HBTU (9.3 mg, 0.0245


mmol) in anhydrous DMF (1.5 mL) was stirred under nitrogen at ambient temperatures for 15 min and treated with the product of Part H, above (18.7 mg, 0.0137 mmol). The DMF was removed under vacuum after 75 min and the
5 resulting amber oil was purified by HPLC on a Vydac C-18 column (22 x 250 mm) using a 0.9%/min gradient of 22.5 to 58.5% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 26.1 min was lyophilized to give the title compound as a colorless
10 fluffy solid (7.5 mg, 53%). MS: m/e 1809.7 [M+H].

Part J - Preparation of (2S)-2-[(2,6-Dimethyl-4-[3-(N-{2-[3-sulfo-2-(3-sulfo-2-(2-[1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)cyclododecyl]acetyl amino)propyl)-propyl]ethyl)carbamoyl]propoxy]phenyl)sulfonyl amino]-3-
15 ({7-[(imidazol-2-ylamino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic Acid
Trifluoroacetate Salt

20 The product of Step I, above (7.5 mg, 0.0039 mmol) was dissolved in a solution of peroxide-free THF (1.40 mL) and water (0.21 mL), and treated with 3 N LiOH (0.14 mL). The mixture was stirred at ambient temperatures under nitrogen for 1 h, and concentrated to dryness under
25 vacuum. The resulting solid residue was dissolved in 95/5 TFA/Et₃SiH (2.0 mL) and heated at 70 °C under nitrogen for 1 h. The solution was concentrated under vacuum and the resulting solid residue was purified by
30 HPLC on a Vydac C-18 column (22 x 250 mm) using a 0.90%/min gradient of 0 to 27% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 20.5 min was lyophilized to give the title compound as a colorless fluffy solid (4.2 mg, 71.9%). MS: m/e 1385.3 [M+H]; High Resolution MS: Calcd for C₅₄H₇₇N₁₄O₂₃S₃ [M+H]:
35 1385.4448; found: 1385.446.


Example 30

Synthesis of DOTA/(2S)-2-[[[4-{3-[N-(2-{2-[(4S)-4-(N-{1-
 {N-(2-{4-[4-(((1S)-1-Carboxy-2-((7-[(imidazol-2-
 5 carbonylamino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))-
 carbonylamino)ethyl]amino)sulfonyl]-3,5-dimethylphenoxy]-
 butanoylamino)ethyl]carbamoyl]-2-sulfoethyl]carbamoyl]-4-
 aminobutanoylamino]-3-
 sulfopropyl)ethyl]carbamoyl]propoxy}-2,6-
 10 dimethylphenyl)sulfonyl]amino]-3-((7-[(imidazol-2-
 ylamino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))-
 carbonylamino)propanoic Acid Conjugate
 Bis(trifluoroacetate) Salt



15

Part A - Preparation of Di-2,3,5,6-tetrafluorophenyl
 (2S)-2-[(tert-Butoxy)carbonylamino]pentane-1,5-dioate




20

To a solution of Boc-L-Glu-OH (28.9 g, 117 mmol) in DMF (500 mL) at ambient temperatures and under nitrogen, was added a solution of 2,3,5,6-tetrafluorophenol (48.2


g, 290 mmol) in DMF (50 mL). After stirring for 10 min, EDC (55.6 g, 290 mmol) was added and the mixture was stirred for 96 h. The volatiles were removed under vacuum and the residue was triturated with 0.1 N HCl (750 mL). To this mixture was added EtOAc (600 mL) and the layers were separated. The aqueous layer was extracted with EtOAc (3 x 500 mL), and all EtOAc extracts were combined, washed consecutively with water (300 mL) and saturated NaCl (300 mL), dried (MgSO₃), and concentrated to give a tan solid (62 g). The tan solid was washed with ACN to give the title compound (45.5 g, 73.0%) in purified form. MS: m/e 566.0 [M+Na].

Part B - Preparation of (2R)-2-[4-(N-((1R)-1-[N-(2-[4-[4-(((1S)-2-((7-[[[1-(Triphenylmethyl)imidazol-2-yl]amino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)-1-(methoxycarbonyl)ethyl]amino)sulfonyl)-3,5-dimethylphenoxy]butanoylamino)ethyl)carbamoyl]-2-sulfoethyl)carbamoyl](4S)-4-[(tert-butoxy)carbonylamino]-butanoylamino]-N-(2-[4-[4-(((1S)-2-((7-[[[1-(imidazol-2-ylamino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)-1-(methoxycarbonyl)ethyl]amino)sulfonyl)-3,5-dimethylphenoxy]butanoylamino)ethyl)propanesulfonic Acid




A solution of the product of Example 29, Part F (130 mg, 0.118 mmol), the product of Part A, above (27.2 mg, 0.050 mmol), and DIEA (0.070 mL, 0.40 mmol) in anhydrous DMF (4.0 mL) was stirred at ambient temperatures under
 5 nitrogen for 29 h. The DMF was removed under vacuum and the resulting amber oil was purified by HPLC on a Vydac C-18 column (50 x 250 mm) using a 0.90%/min gradient of 22.5 to 58.5% ACN containing 0.1% TFA at a flow rate of 80 mL/min. The main product peak eluting at 35.7 min was
 10 lyophilized to give the title compound as a colorless fluffy solid (108 mg, 89.3%). MS: m/e 2419.6 [M+H], 1210.4 [M+2H].

Part C - Preparation of (2R)-2-[4-(N-((1R)-1-[N-(2-{4-[4-
 15 (((1S)-2-((7-[[1-(Triphenylmethyl)imidazol-2-yl]amino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)-1-(methoxycarbonyl)ethyl]amino)sulfonyl)-3,5-dimethylphenoxy]butanoylamino)ethyl)carbamoyl]-2-sulfoethyl)carbamoyl (4S)-4-aminobutanoylamino]-N-(2-{4-
 20 [4-(((1S)-2-((7-[[1-(imidazol-2-ylamino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)-1-(methoxycarbonyl)ethyl]amino)sulfonyl)-3,5-dimethylphenoxy]butanoylamino)ethyl)propanesulfonic Acid



The product of Part B, above (107 mg, 0.0442 mmol) was dissolved in 50/50 TFA/DCM (5.0 mL) and allowed to react at ambient temperatures under nitrogen for 10 min. The solution was concentrated and the resulting amber oil was dissolved in 50% ACN (25 mL) and lyophilized to give the title compound as a pale yellow solid (105 mg, 98.0%). MS: m/e 1159.9 [M+2H], 1039.4 [M+2H-Trt].

Part D - Preparation of DOTA tri-t-Butyl Ester/(2R)-2-[4-(N-((1R)-1-[N-(2-[4-[4-(((1S)-2-((7-[[1-(Triphenylmethyl)imidazol-2-yl]amino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)-1-(methoxycarbonyl)ethyl]amino)sulfonyl)-3,5-dimethylphenoxy]butanoylamino)ethyl)carbamoyl]-2-sulfoethyl)carbamoyl (4S)-4-aminobutanoylamino]-N-(2-[4-[4-(((1S)-2-((7-[[imidazol-2-yl]amino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)-1-(methoxycarbonyl)ethyl]amino)sulfonyl)-3,5-dimethylphenoxy]butanoylamino)ethyl)propanesulfonic Acid Conjugate



A solution of 2-(1,4,7,10-tetraaza-4,7,10-tris(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)acetic acid (31.6 mg, 0.0346 mmol) (as described in DM-7003), DIEA (0.072 mL, 0.416 mmol), and HBTU (9.8 mg, 0.026

mmol) in anhydrous DMF (1.8 mL) was stirred under nitrogen at ambient temperatures for 15 min and treated with the product of Part C, above (40.0 mg, 0.0173 mmol). The DMF was removed under vacuum after 90 min and the
5 resulting pale yellow oil was purified by HPLC on a Vydac C-18 column (50 x 250 mm) using a 1.01%/min gradient of 22.5 to 63.0% ACN containing 0.1% TFA at a flow rate of 80 mL/min. The main product peak eluting at 27.6 min was lyophilized to give the title compound as a colorless
10 solid (29.0 mg, 62.4%). MS: m/e 1437.6 [M+2H], 1316.6 [M+2H-Trt].

Part E - Preparation of DOTA/(2S)-2-[[[4-{3-[N-(2-{2-
15 [(4S)-4-(N-{1-[N-(2-{4-[4-(((1S)-1-Carboxy-2-((7-
[(imidazol-2-ylamino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)ethyl]amino)sulfonyl]-3,5-dimethylphenoxy]butanoylamino)ethyl]carbamoyl]-2-sulfoethyl]carbamoyl]-4-aminobutanoylamino]-3-sulfopropyl]ethyl]carbamoyl]propoxy]-2,6-dimethylphenyl)sulfonyl]amino)-3-((7-[(imidazol-2-ylamino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)propanoic Acid Conjugate
20 Bis(trifluoroacetate) Salt

25 A mixture of the product of Part D, above (30.0 mg, 0.0104 mmol), peroxide-free THF (3.2 mL), water (0.485 mL), and 3 N LiOH (0.320 mL, 0.96 mmol) was stirred at ambient temperatures under nitrogen for 2 h. The solution was concentrated under vacuum and the resulting
30 solid residue was dissolved in 95/5 TFA/Et₃SiH (5.0 mL). The solution was heated at 70 °C under nitrogen for 1 h and concentrated under vacuum. The resulting oily solid was purified by HPLC on a Vydac C-18 column (22 x 250 mm) using a 0.90%/min gradient of 0 to 27% ACN containing
35 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 27.8 min was lyophilized to give the title compound as a colorless fluffy solid (12.8 mg, 48.5%). MS: m/e 1096.8 [M+2H], 731.8 [M+3H]; High


Resolution MS: Calcd for $C_{91}H_{122}N_{23}O_{33}S_4$ [M+H]: 2192.7458;
Found: 2192.741.

5

Example 31

Synthesis of 2-(((4-[3-(N-[2-((2R)-2-((2R)-3-Sulfo-2-{2-[1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)cyclododecyl] acetylamino}propyl)-3-sulfopropyl)ethyl]carbamoyl)propoxy]-2,6-dimethylphenyl)sulfonyl)amino)(2S)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)propanoic Acid Trifluoroacetate Salt


10



15

Part A - Preparation of 2-(((4-[3-(N-[2-((2R)-2-Amino-3-sulfopropyl)ethyl]carbamoyl)propoxy]-2,6-dimethylphenyl)sulfonyl)amino)(2S)-3-((1-methyl-4-oxo-7-((1-(triphenylmethyl)imidazol-2-yl)amino)methyl)(3-hydroquinolyl))]carbonylamino)propanoic Acid


20



A mixture of the product of Example 29, Part F (125 mg, 0.113 mmol), peroxide-free THF (3.8 mL), water (0.57 mL), and 3 N LiOH (0.38 mL, 1.13 mmol) was stirred at ambient temperatures under nitrogen for 1 h. The mixture

was adjusted to pH 1 using 1 N HCl (0.70 mL) and concentrated to dryness under vacuum. The resulting solid was purified by HPLC on a Vydac C-18 column (50 x 250 mm) using a 0.90%/min gradient of 18 to 54% ACN containing 0.1% TFA at a flow rate of 80 mL/min. The main product peak eluting at 21.0 min was lyophilized to give the title compound as a colorless solid (96.0 mg, 77.9%). MS: m/e 1090.3 [M+H], 848.2 [M+H-Trt]; High Resolution MS: Calcd for C₅₄H₆₀N₉O₁₂S₂ [M+H]: 1090.3808; Found: 1090.381.

Part B - Preparation of 2-((4-(3-(N-[2-((2R)-2-((2R)-2-((tert-Butoxy)carbonylamino)-3-sulfopropyl)-3-sulfopropyl)ethyl]carbonyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)(2S)-3-([1-methyl-4-oxo-7-((1-(triphenylmethyl)imidazol-2-yl)amino)methyl)(3-hydroquinolyl)]carbonylamino)propanoic Acid




20

A solution of Boc-L-cysteic acid (37.0 mg, 0.128 mmol), DIEA (0.040 mL, 0.228 mmol), and PyBOP (53.0 mg, 0.102 mmol) in anhydrous DMF (1.0 mL) was stirred at ambient temperatures under nitrogen for 15 min, and added to a solution of the product of Part A, above (93.0 mg, 0.0854 mmol) and DIEA (0.045 mL, 0.256 mmol) in anhydrous DMF (3.0 mL). The resulting solution was stirred at ambient temperatures under nitrogen for 1.5 h and concentrated to a viscous amber oil. Purification by HPLC on a Vydac C-18 column (50 x 250 mm) using a 0.68%/min gradient of 18 to 45% ACN containing 0.1% TFA at a flow rate of 80 mL/min. The main product peak

eluting at 36.4 min was lyophilized to give the title compound as a colorless solid (94.0 mg, 82.1%). MS: m/e 1341.2 [M+H], 1099.1 [M+H-Trt], 999.1 [M+H-Trt-Boc].

- 5 Part C - Preparation of 2-[[[4-{3-[N-(2-{(2R)-2-[(2R)-3-Sulfo-2-(2-{1,4,7,10-tetraaza-4,7,10-tris[(tert-butoxycarbonyl)methyl]cyclododecyl)acetylamino)propyl]-3-sulfopropyl}ethyl)carbamoylethyl]propoxy}-2,6-dimethylphenyl)-sulfonylamino](2S)-3-[[1-methyl-4-oxo-7-([1-(triphenylmethyl)imidazol-2-yl]amino)methyl](3-hydroquinolyl)]carbonylamino]propanoic Acid



- 15 A solution of the product of Part B, above (90.0 mg, 0.0672 mmol) in 50/50 TFA/DCM (10.0 mL) was allowed to react at ambient temperatures under nitrogen for 10 min and concentrated under vacuum to give the intermediate amine as an amber oil. MS: m/e 1241.3 [M+H], 999.3 [M+H-Trt]; High Resolution MS: Calcd for C₅₇H₆₅N₁₀O₁₆S₃ [M+H]: 1241.3742; Found: 1241.375.

- A solution of 2-(1,4,7,10-tetraaza-4,7,10-tris(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)acetic acid (123 mg, 0.134 mmol) (as described in DM-7003), DIEA (0.092 mL, 0.538 mmol), and PyBOP (52.4 mg, 0.101 mmol) in anhydrous DMF (1.5 mL) was stirred under nitrogen at ambient temperatures for 15 min, and added to a solution of the free amine produced above (90.0 mg, 0.0672 mmol) and DIEA (0.046 mL, 0.269 mmol) in anhydrous DMF (1.5 mL). The DMF was removed under vacuum after 1 h and the resulting amber oil was purified by HPLC on a Vydac C-18 column (50 x 250 mm) using a 0.288%/min gradient of 30.6


to 45% ACN containing 0.1% TFA at a flow rate of 80 mL/min. The main product peak eluting at 25.8 min was lyophilized to give the title compound as a colorless solid (92.0 mg, 76.3%). MS: m/e 1795.6 [M+H], 1553.5 [M+H-Trt]; High Resolution MS: Calcd for C₈₅H₁₁₅N₁₄O₂₃S₃ [M+H]: 1795.7422; Found: 1795.744.

Part D - Preparation of 2-([4-[3-(N-(2-((2R)-2-((2R)-3-sulfo-2-(2-[1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)cyclododecyl] acetylamino)propyl)-3-sulfopropyl)ethyl]carbamoyl)propoxy)-2,6-dimethylphenyl)sulfonyl]amino)(2S)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)propanoic Acid Trifluoroacetate Salt


A solution of the product of Part C, above (89.0 mg, 0.0496 mmol) in 97/3 TFA/Et₃SiH (10.0 mL) was heated at 70 °C under nitrogen for 30 min and concentrated under vacuum. The resulting oily solid was purified by HPLC on a Vydac C-18 column (50 x 250 mm) using a 0.45%/min gradient of 4.5 to 22.5% ACN containing 0.1% TFA at a flow rate of 80 mL/min. The main product peak eluting at 19.5 min was lyophilized to give stereochemically pure title compound as a colorless fluffy solid (65.0 mg, 87.5%). MS: m/e 1385.4 [M+H].

Example 32

Alternative Synthesis of Intermediate 2-([4-[3-(N-(2-((2R)-2-Amino-3-sulfopropyl)ethyl]carbamoyl)propoxy)-2,6-dimethylphenyl)sulfonyl]amino)(2S)-3-([1-methyl-4-oxo-7-([1-(triphenylmethyl)imidazol-2-yl]amino)methyl)(3-hydroquinolyl)]carbonylamino)propanoic Acid



Part A - Preparation of (2S)-2-({(4-{3-[N-(2-Aminoethyl)carbamoyl]propoxy}-2,6-dimethylphenyl)-sulfonyl}amino)-3-({[1-methyl-4-oxo-7-({[1-(triphenylmethyl)imidazol-2-yl]amino)methyl}(3-hydroquinolyl)]carbonylamino}propanoic Acid




10

A mixture of the product of Example 29, Part D (956 mg, 1.004 mmol), peroxide-free THF (35 mL), water (5.3 mL), and 3 N LiOH (3.53 mL, 10.6 mmol) was stirred at ambient temperatures under nitrogen for 1 h, and adjusted to pH 5-6 using 1 N HCl (10 mL). The THF was removed under vacuum causing a gummy yellow solid to precipitate. The water layer was removed by decantation and the solid was washed with water (15 mL). The solid was dried under vacuum to give the title compound as a dry yellow solid

20

Part B - Preparation of 2-({(4-{3-[N-(2-((2R)-2-[(tert-Butoxy)carbonylamino]-3-sulfopropyl)ethyl)carbamoyl]-propoxy}-2,6-dimethylphenyl)sulfonyl}amino)(2S)-3-({[1-methyl-4-oxo-7-({[1-(triphenylmethyl)imidazol-2-yl]amino)methyl}(3-hydroquinolyl)]carbonylamino}propanoic Acid

25




A solution of Boc-L-cysteic acid (175 mg, 0.60 mmol), DIEA (0.208 mL, 1.20 mmol), and PyBOP (250 mg, 0.480 mmol) in anhydrous DMF (5.0 mL) was stirred at ambient temperatures under nitrogen for 17 min, and added to a solution of the product of Part A, above (375 mg, 0.400 mmol) and DIEA (0.070 mL, 0.400 mmol) in anhydrous DMF (4.0 mL). The resulting solution was stirred at ambient temperatures under nitrogen for 45 min and concentrated under vacuum to give an amber oil. Purification by PLC on a Vydac C-18 column (50 x 250 mm) using a 0.292%/min gradient of 31.5 to 43.2% ACN containing 0.1% TFA at a flow rate of 80 mL/min. The main product peak eluting at 22.0 min was lyophilized to give the title compound as a colorless solid (430 mg, 90.4%). MS: m/e 1190.3 [M+H], 948.3 [M+H-Trt].

Part C - Preparation of 2-([4-(3-{N-[2-((2R)-2-Amino-3-sulfopropyl)ethyl]carbonyl}propoxy)-2,6-dimethylphenyl]-sulfonyl)amino)(2S)-3-([1-methyl-4-oxo-7-([1-(triphenylmethyl)imidazol-2-yl]amino)methyl)(3-hydroquinolyl)]carbonylamino)propanoic Acid

A solution of the product of Part B, above (430 mg, 0.362 mmol) in 50/50 TFA/DCM (15 mL) was allowed to react at ambient temperatures under nitrogen for 10 min and concentrated under vacuum. The resulting amber oil was taken up in 50% ACN (50 mL) and lyophilized to give the title compound as a pale yellow solid (398 mg, 100%). MS: m/e 1090.3 [M+H], 848.2 [M+H-Trt].

Example 33


Synthesis of DOTA/2-[[[4-{3-[N-(2-{(2R)-2-{(2R)-2-(4-[N-
 [(1R)-1-(N-{(1R)-1-[N-(2-{4-[4-(((1S)-1-Carboxy-2-({7-
 [(imidazol-2-ylamino)methyl]-1-methyl-4-oxo(3-
 5 hydroquinolyl)}carbonylamino)-1-carboxyethyl]amino)-
 sulfonyl)-3,5-dimethylphenoxy]butanoylamino)ethyl)-
 carbamoyl]-2-sulfoethyl}carbamoyl)-2-
 sulfoethyl}carbamoyl](2S)-2-aminobutanoylamino)-3-
 sulfopropyl]-3-sulfopropyl]ethyl}carbamoyl]propoxy}-2,6-
 10 dimethylphenyl)sulfonyl]amino)(2S)-3-({7-[(imidazol-2-
 ylamino)methyl]-1-methyl-4-oxo(3-hydroquinolyl)}-
 carbonylamino)propanoic Acid Conjugate



15

Part A - Preparation of 2-[[[4-{3-[N-(2-{(2R)-2-{(2R)-2-
 (4-[N-[(1R)-1-(N-{(1R)-1-[N-(2-{4-[4-(((1S)-1-Carboxy-2-
 ({7-[[[1-(triphenylmethyl)imidazol-2-yl]amino)methyl]-1-
 methyl-4-oxo(3-hydroquinolyl)}carbonylamino)-1-
 20 carboxyethyl]amino)sulfonyl)-3,5-
 dimethylphenoxy]butanoylamino)ethyl}carbamoyl]-2-
 sulfoethyl}carbamoyl)-2-sulfoethyl}carbamoyl](2S)-2-
 [(tert-butoxy)carbonylamino]butanoylamino)-3-
 sulfopropyl]-3-sulfopropyl]ethyl}carbamoyl]propoxy}-2,6-
 25 dimethylphenyl)sulfonyl]amino)(2S)-3-({7-[[[1-

(triphenylmethyl)imidazol-2-yl)amino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic Acid




5

A solution of the product of the first half of Example 31, Part C (136 mg, 0.110 mmol), DIEA (0.076 mL, 0.44 mmol), and the product of Example 30, Part A (26.2 mg, 0.050 mmol) in anhydrous DMF (3.0 mL) was stirred at ambient temperatures under nitrogen for 7 h. The DMF was removed under vacuum and the viscous amber oil was purified by HPLC on a Vydac C-18 column (50 x 250 mm) using a 0.45%/min gradient of 27 to 45% ACN, followed by a 0.72% gradient of 45-63% ACN containing 0.1% TFA at a flow rate of 80 mL/min. The main product peak eluting at 75.2 min was lyophilized to give the title compound as a colorless solid (129 mg, 47.9%). MS: m/e 1347.3 [M+2H].

Part B - Preparation of DOTA tri-t-Butyl Ester Conjugate of 2-[[[4-{3-[N-(2-{(2R)-2-[(2R)-2-(4-{N-[(1R)-1-(N-[(1R)-1-[N-(2-{4-[4-[[[(1S)-1-Carboxy-2-[(7-[(1-(triphenylmethyl)imidazol-2-yl)amino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)-1-carboxyethyl]amino)sulfonyl]-3,5-dimethylphenoxy]butanoylamino)ethyl)carbamoyl]-2-

sulfoethyl}carbamoyl)-2-sulfoethyl}carbamoyl}(2S)-2-
 aminobutanoylamino)-3-sulfopropyl]-3-sulfopropyl]-
 ethyl}carbamoyl]propoxy}-2,6-dimethylphenyl)sulfonyl]-
 amino}(2S)-3-({7-[(1-(triphenylmethyl)imidazol-2-
 5 yl)amino)methyl]-1-methyl-4-oxo(3-hydroquinolyl))-
 carbonylamino)propanoic Acid



- 10 The product of Part A, above (34.0 mg, 0.0126 mmol)
 was dissolved in 50/50 TFA/DCM (12 mL) and allowed to
 react at ambient temperatures under nitrogen for 10 min.
 The solution was concentrated and the resulting amber oil
 was dried under vacuum.
- 15 A solution of 2-(1,4,7,10-tetraaza-4,7,10-
 tris(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)acetic
 acid (23.1 mg, 0.0253 mmol), DIEA (0.020 mL, 0.115 mmol),
 and PyBOP (9.8 mg, 0.019 mmol) in anhydrous DMF (2.0 mL)
 was stirred under nitrogen at ambient temperatures for 15
 20 min, and added to a solution of the product from the
 deprotection reaction, above and DIEA (0.020 mL, 0.115
 mmol) in anhydrous DMF (2.0 mL). The DMF was removed
 under vacuum after 2 h, and the resulting residue was
 purified by HPLC on a Vydac C-18 column (50 x 250 mm)
 25 using a 0.45%/min gradient of 27 to 49.5% ACN containing

0.1% TFA at a flow rate of 80 mL/min. The main product peak eluting at 43.8 min was lyophilized to give the title compound as a colorless solid (16.0 mg, 40.4%).

MS: m/e 1574.8 [M+2H], 1453.7 [M+2H-Trt], 1332.2 [M+2H-2Trt].

Part C - Preparation of DOTA/2-[[[4-{3-[N-(2-{(2R)-2-[(2R)-2-(4-{N-[(1R)-1-(N-{(1R)-1-[N-(2-{4-[4-([(1S)-1-Carboxy-2-({7-[(imidazol-2-ylamino)methyl]-1-methyl-4-oxo(3-hydroquinolyl)}carbonylamino)-1-carboxyethyl]amino)sulfonyl]-3,5-dimethylphenoxy]butanoylamino)ethyl)carbamoyl]-2-sulfoethyl}carbamoyl)-2-sulfoethyl}carbamoyl](2S)-2-aminobutanoylamino)-3-sulfopropyl]-3-sulfopropyl]-ethyl)carbamoyl]propoxy}-2,6-dimethylphenyl)sulfonyl]amino)(2S)-3-({7-[(imidazol-2-ylamino)methyl]-1-methyl-4-oxo(3-hydroquinolyl)}carbonylamino)propanoic Acid Conjugate


The product of Part B, above (14.0 mg, 0.00445 mmol) was dissolved in 95/5 TFA/Et₃SiH (8.0 mL) and heated at 70 °C under nitrogen for 1 h. The solution was concentrated under vacuum and the resulting yellow solid was purified by HPLC on a Vydac C-18 column (22 x 250 mm) using a 0.9%/min gradient of 0 to 27% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 24.5 min was lyophilized to give the title compound as a colorless solid (8.2 mg, 73.9%). MS: m/e 1247.7 [M+2H].

30

Example 34


Synthesis of (2S)-3-([7-[(Imidazol-2-ylamino)methyl]-4-oxo-1-(3-{2-[1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)cyclododecyl]acetylamino)propyl}(3-hydroquinolyl)}carbonylamino)-2-[(2,4,6-trimethylphenyl)sulfonyl]amino)propanoic Acid Tris(trifluoroacetate) Salt

35



Part A - Preparation of (2S)-3-({7-[(Imidazol-2-ylamino)methyl]-4-oxo-1-[3-(2-{1,4,7,10-tetraaza-4,7,10-
 5 tris[(tert-butoxycarbonyl)methyl]cyclododecyl)acetylaminopropyl](3-hydroquinolyl)}carbonylamino)-2-[[2,4,6-trimethylphenyl)sulfonyl]amino]propanoic Acid
 Tris(trifluoroacetate) Salt

10



A solution of 2-(1,4,7,10-tetraaza-4,7,10-tris(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)acetic
 15 acid (89 mg, 0.0974 mmol) (as described in DM-7003), DIEA (0.103 mL, 0.607 mmol), and HBTU (28.0 mg, 0.0735 mmol) in anhydrous DMF (1.0 mL) was stirred under nitrogen at ambient temperatures for 15 min and treated with a
 solution of the product of Example 4, Part H (30.0 mg,
 20 0.049 mmol) in anhydrous DMF (1.0 mL). The DMF was removed under vacuum after 3 h and the residue was purified by HPLC on a Vydac C-18 column (22 x 250 mm) using a 1.08%/min gradient of 18 to 72% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product


peak eluting at 17.5 min was lyophilized to give the title compound as a colorless solid (48.0 mg, 65.0%). MS: m/e 1164.7 [M+H].

- 5 Part B - Preparation of (2S)-3-[[7-[(Imidazol-2-ylamino)methyl]-4-oxo-1-(3-{2-[1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)cyclododecyl]acetylamino}propyl)(3-hydroquinolyl)]carbonylamino]-2-[[{(2,4,6-trimethylphenyl)sulfonyl]amino}propanoic Acid
- 10 Tris(trifluoroacetate) Salt


- A solution of the product of Part A, above (48.0 mg, 0.0375 mmol) in 95/5 TFA/Et₃SiH (2.1 mL) was stirred at 50 °C under nitrogen for 2 h. The solution was
- 15 concentrated under vacuum and the oily residue was purified by HPLC on a Vydac C-18 column (22 x 250 mm) using a 1.2%/min gradient of 0 to 36% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 18.6 min was lyophilized to give the title
- 20 compound as a colorless solid (25.7 mg, 51.2%). MS: m/e 996.5 [M+H]; High Resolution MS: Calcd for C₄₅H₆₂N₁₁O₁₃S [M+H]: 996.4249; Found: 996.4278.

Example 35

- 25 Synthesis of 3-((1-[3-((2R)-3-Sulfo-2-{2-[1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)cyclododecyl]-acetylamino}propyl)propyl]-7-[(imidazol-2-ylamino)methyl]-4-oxo(3-hydroquinolyl))carbonylamino)(2S)-2-[[{(2,4,6-trimethylphenyl)sulfonyl]amino}propanoic Acid
- 30 Bis(trifluoroacetate) Salt



Part A - Preparation of 3-([1-(3-((2R)-2-[(tert-Butoxy)carbonylamino]-3-sulfopropyl)propyl)-7-[(imidazol-2-ylamino)methyl]-4-oxo(3-hydroquinolyl)]carbonylamino)(2S)-2-[(2,4,6-trimethylphenyl)sulfonylamino]propanoic Acid




10

A solution of the product of Example 4, Part H (105 mg, 0.125 mmol), the N-hydroxysuccinimide ester of Boc-Cysteic acid (as described in *Liebigs Ann. Chem.* 1979, 776-783) (146 mg, 0.467 mmol), and DIEA (0.120 mL, 0.69 mmol) in anhydrous DMF (1.5 mL) was stirred at ambient temperatures under nitrogen for 24 h. The DMF was removed under vacuum and the resulting solid residue was purified by HPLC on a Vydac C-18 column (22 x 250 mm) using a 0.68%/min gradient of 9 to 36% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 30.3 min was lyophilized to give the title compound as a colorless solid (73.0 mg, 67.9%). MS: m/e 861.3 [M+H].

Part B - Preparation of 3-([1-[3-((2R)-2-Amino-3-sulfopropyl)propyl]-7-[(imidazol-2-ylamino)methyl]-4-oxo(3-hydroquinolyl)]carbonylamino)(2S)-2-[(2,4,6-trimethylphenyl)sulfonylamino]propanoic Acid


5 Trifluoroacetate Salt



The product of Part B, above (70.0 mg, 0.0814 mmol)
 10 was dissolved in 2:1 DCM/TFA (1.5 mL) and allowed to
 react at ambient temperatures under nitrogen for 30 min.
 The solution was concentrated under vacuum and the amber
 oil was dissolved in 50% ACN (25 mL) and lyophilized to
 give the title compound as a colorless solid (70.8 mg,
 15 99.5%). MS: m/e 761.2 [M+H]; High Resolution MS: Calcd
 for C₃₂H₄₁N₈O₁₀S₂ [M+H]: 761.2387; Found: 761.2393.

Part C - Preparation of 3-([1-[3-[(2R)-3-Sulfo-2-(2-
 {1,4,7,10-tetraaza-4,7,10-tris[(tert-butoxycarbonyl)-
 20 methyl]cyclododecyl)acetylamino]propyl]propyl]-7-
 [(imidazol-2-ylamino)methyl]-4-oxo(3-
 hydroquinolyl)]carbonylamino)(2S)-2-[(2,4,6-
 trimethylphenyl)sulfonylamino]propanoic Acid
 Bis(trifluoroacetate) Salt

25




A solution of 2-(1,4,7,10-tetraaza-4,7,10-tris(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)acetic acid (20.8 mg, 0.0228 mmol) (as described in DM-7003), DIEA (0.006 mL, 0.034 mmol), and HBTU (6.5 mg, 0.0171 mmol) in anhydrous DMF (0.5 mL) was stirred under nitrogen at ambient temperatures for 5 min and treated with a solution of the product of Part B, above (10.0 mg, 0.0114 mmol) and DIEA (0.006 mL, 0.034 mmol) in anhydrous DMF (0.5 mL). Stirring was continued at ambient temperatures for 24 h, and the reaction was diluted with water (3.0 mL), treated with concentrated ammonium hydroxide (0.003 mL), and stirred an additional 10 min. The solution was adjusted to pH 3 using 0.1 N HCl (6.0 mL) and diluted further with 10% ACN (5.5 mL). This solution was purified directly by HPLC on a Vydac C-18 column (22 x 250 mm) using a 0.68%/min gradient of 9 to 36% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 36.0 min was lyophilized to give the title compound as a colorless solid (12.0 mg, 68.3%). MS: m/e 1315.6 [M+H].

Part D - Preparation of 3-({1-[3-((2R)-3-Sulfo-2-{2-
25 [1,4,7,10-tetraaza-4,7,10-
tris(carboxymethyl)cyclododecyl]-
acetyl amino)propyl)propyl]-7-[(imidazol-2-
ylamino)methyl]-4-oxo(3-
hydroquinolyl)}carbonylamino)(2S)-2-[(2,4,6-
30 trimethylphenyl)sulfonyl]amino)propanoic Acid
Bis(trifluoroacetate) Salt

A solution of the product of Part C, above (12.0 mg, 0.00778 mmol) in 95/5 TFA/Et₃SiH (1.0 mL) was stirred at ambient temperatures under nitrogen for 18 h. The solution was concentrated under vacuum and the oily residue was purified by HPLC on a Vydac C-18 column (22 x 250 mm) using a 1.2%/min gradient of 0 to 36% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 21.1 min was lyophilized to give the title compound as a colorless solid (8.1 mg, 75.7%). MS: m/e 1147.3 [M+H]; High Resolution MS: Calcd for C₄₈H₆₇N₁₂O₁₇S₂ [M+H]: 1147.4189; Found: 1147.418.

Example 36

Synthesis of 3-[[[1-(3-{2-[(6-[(1E)-1-Aza-2-(2-sulfophenyl)vinyl]amino)(3-pyridyl)carbonylamino](2R)-3-sulfopropyl}propyl)-7-[(imidazol-2-ylamino)methyl]-4-oxo(3-hydroquinolyl)]carbonylamino](2S)-2-[(2,4,6-trimethylphenyl)sulfonyl]amino)propanoic Acid




A solution of the product of Example 35, Part B (10.0 mg, 0.0101 mmol), DIEA (0.007 mL, 0.040 mmol), and 2-(2-aza-2-((5-((2,5-dioxopyrrolidinyl)carbonyl)(2-pyridyl))amino)vinyl)benzenesulfonic acid (5.3 mg, 0.0120 mmol) in anhydrous DMF (0.5 mL) was allowed to stand at ambient temperatures under a nitrogen atmosphere for 48 h. Additional 2-(2-aza-2-((5-((2,5-dioxopyrrolidinyl)-carbonyl)(2-pyridyl))amino)vinyl)benzenesulfonic acid (2.0 mg, 0.00455 mmol) was added and stirring was continued an additional 48 h. The DMF was removed under vacuum and the residue was purified by HPLC on a Vydac C-

18 column (22 x 250 mm) using a 0.9%/min gradient of 0 to 36% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 30.0 min was lyophilized to give the title compound as a colorless solid (2.5 mg, 23.3%). MS: m/e 1064.3 [M+H]; High Resolution MS: Calcd for C₄₅H₅₀N₁₁O₁₄S₃ [M+H]: 1064.27005; Found: 1064.272.

Example 37

Synthesis of 3-[[1-(3-[(2R)-2-[4-(N-[(1R)-1-[N-(3-{3-[N-
 10 ((2S)-2-Carboxy-2-[(2,4,6-trimethylphenyl)sulfonyl]-
 amino)ethyl)carbamoyl]-7-[(imidazol-2-ylamino)methyl]-4-
 oxohydroquinoly]propyl)carbamoyl]-2-sulfoethyl)-
 carbamoyl)(2S)-2-(2-[1,4,7,10-tetraaza-4,7,10-
 tris(carboxymethyl)cyclododecyl]acetylaminobutanoylamino
 15]-3-sulfopropyl)propyl)-7-[(imidazol-2-ylamino)methyl]-4-
 oxo(3-hydroquinoly)]carbonylamino)(2S)-2-[(2,4,6-
 trimethylphenyl)sulfonyl]amino]propanoic Acid
 Bis(trifluoroacetate) Salt




20

Part A - Preparation of 3-[[1-(3-[(2R)-2-[4-(N-[(1R)-1-[N-(3-{3-[N-
 [N-(3-{3-[N-((2S)-2-Carboxy-2-[(2,4,6-trimethylphenyl)-
 sulfonyl]amino)ethyl)carbamoyl]-7-[(imidazol-2-
 25 ylamino)methyl]-4-oxohydroquinoly]propyl)carbamoyl]-2-
 sulfoethyl)carbamoyl)(2S)-2-[(tert-butoxy)carbonylamino]-

butanoylamino]-3-sulfopropyl)propyl)-7-[(imidazol-2-ylamino)methyl]-4-oxo(3-hydroquinolyl)]carbonylamino)(2S)-2-[[[2,4,6-trimethylphenyl)sulfonyl]amino]propanoic Acid

5




A solution of the product of Example 35, Part B (38.0 mg, 0.0434 mmol), DIEA (0.015 mL, 0.0869 mmol), and the product of Example 30, Part A (10.9 mg, 0.0202 mmol) in anhydrous DMF (1.0 mL) was stirred at ambient temperatures under nitrogen for 48 h. The DMF was removed under vacuum and the amber oil was purified by HPLC on a Vydac C-18 column (22 x 250 mm) using a 0.68%/min gradient of 9 to 36% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 36.1 min was lyophilized to give the title compound as a colorless solid (13.5 mg, 38.6%). MS: m/e 1732.4 [M+H]⁺, 1632.2 [M+H-Boc]⁺.

20

Part B - Preparation of 3-[[[1-(3-[(2R)-2-[4-(N-[(1R)-1-[N-(3-[3-[N-[(2S)-2-Carboxy-2-[(2,4,6-trimethylphenyl)-sulfonyl]amino)ethyl)carbamoyl]-7-[(imidazol-2-ylamino)methyl]-4-oxohydroquinolyl]propyl)carbamoyl]-2-sulfoethyl)carbamoyl](2S)-2-aminobutanoylamino]-3-sulfopropyl]propyl]-7-[(imidazol-2-ylamino)methyl]-4-

oxo(3-hydroquinolyl)]carbonylamino)(2S)-2-[[(2,4,6-trimethylphenyl)sulfonylamino]propanoic Acid
Trifluoroacetate Salt




5

The product of Part A, above (13.5 mg, 0.00779 mmol), was dissolved in 50/50 TFA/DCM (1.0 mL) and allowed to react at ambient temperatures under nitrogen for 45 min.
10 The solution was concentrated under vacuum to give the title compound as a pale amber oil. MS: m/e 1633.3 [M+H].

Part C - Preparation of 3-[[1-(3-[(2R)-2-[4-(N-[(1R)-1-[N-(3-{3-[N-((2S)-2-Carboxy-2-[[(2,4,6-trimethylphenyl)sulfonylamino]ethyl)carbamoyl]-7-[(imidazol-2-ylamino)methyl]-4-oxohydroquinolyl)propyl)carbamoyl]-2-sulfoethyl)carbamoyl)(2S)-2-{2-[1,4,7,10-tetraaza-4,7,10-tris[(tert-
20 butoxycarbonyl)methyl]cyclododecyl]acetyl amino}-butanoylamino]-3-sulfopropyl)propyl)-7-[(imidazol-2-ylamino)methyl]-4-oxo(3-hydroquinolyl)]carbonylamino)-(2S)-2-[[(2,4,6-trimethylphenyl)sulfonylamino]propanoic Acid Bis(trifluoroacetate) Salt

25



A solution of 2-(1,4,7,10-tetraaza-4,7,10-tris(((tert-butyl)oxycarbonyl)methyl)cyclododecyl)acetic acid (15.0 mg, 0.0164 mmol) (as described in DM-7003),
 5 DIEA (0.004 mL), and HBTU (4.7 mg, 0.0124 mmol) in anhydrous DMF (0.5 mL) was stirred under nitrogen at ambient temperatures for 8 min and treated with a solution of the product of Part B, above (0.00779 mmol) and DIEA (0.004 mL) in anhydrous DMF (0.5 mL). The
 10 solution was stirred at ambient temperatures for 24 h, treated with 0.1 N NaOH (0.33 mL), stirred an additional 5 min, and adjusted to pH 3 with 0.1 N HCl (0.60 mL). This solution was diluted with water (4.5 mL) and
 15 purified directly by HPLC on a Vydac C-18 column (22 x 250 mm) using a 1.01%/min gradient of 9 to 49.5% ACN containing 0.1% TFA at a flow rate of 20 mL/min. The main product peak eluting at 26.7 min was lyophilized to give the title compound as a colorless solid (7.0 mg, 37.2%). MS: m/e 1094.4 [M+2H]; High Resolution MS: Calcd
 20 for C₉₇H₁₃₆N₂₁O₂₉S₄ [M+H]: 2186.8696; Found: 2186.867.

Part D - Preparation of 3-[[1-(3-[(2R)-2-[4-(N-[(1R)-1-[N-(3-{3-[N-((2S)-2-Carboxy-2-[(2,4,6-trimethylphenyl)-sulfonyl]amino)ethyl)carbamoyl]-7-[(imidazol-2-ylamino)methyl]-4-oxohydroquinolyl)propyl)carbamoyl]-2-

5 sulfoethyl)carbamoyl) (2S)-2-(2-[1,4,7,10-tetraaza-4,7,10-
tris(carboxymethyl)cyclododecyl]acetylamino)butanoylamino
]-3-sulfopropyl}propyl)-7-[(imidazol-2-ylamino)methyl]-4-
oxo(3-hydroquinolyl)]carbonylamino) (2S)-2-[[2,4,6-
trimethylphenyl)sulfonyl]amino}propanoic Acid
Bis(trifluoroacetate) Salt

10 A solution of the product of Step C, above (7.0 mg,
0.00290 mmol) in 95/5 TFA/Et₃SiH (1.0 mL) was heated to
reflux under nitrogen for 3 h. The solution was
concentrated under vacuum and the oily residue was
purified by HPLC on a Vydac C-18 column (22 x 250 mm)
using a 1.2%/min gradient of 0 to 36% ACN containing 0.1%
TFA at a flow rate of 20 mL/min. The main product peak
15 eluting at 26.5 min was lyophilized to give the title
compound as a colorless solid (4.5 mg, 66.1%). High
Resolution MS: Calcd for C₈₅H₁₁₂N₂₁O₂₉S₄ [M+H]: 2018.6818;
Found: 2018.683.

20

Example 38

Synthesis of the In-111 Complex of the Conjugate Example
29

To a shielded and crimped 2 cc autosampler vial was
added 70 µg of the conjugate of Example 29 dissolved in
25 140 µl 0.5 M ammonium acetate buffer (pH 4.8) followed by
the addition of 2 mg gentisic acid sodium salt and 2.6
mCi (7 µl) In-111 in 0.05M HCl. The reaction mixture
(specific activity was heated at 85°C for 20 minutes and
analyzed by HPLC. Yield: 87.9% (total for the two
30 isomers); Ret. Time: 12.5, 13.1 min.

HPLC Method

Column: Zorbax Rx C18, 25 cm x 4.6 mm

Column Temperature: Ambient

35 Flow: 1.0 ml/min

Solvent A: 10 mM ammonium acetate

Solvent B: Acetonitrile

Detector: IN-US β -ram, and UV at 220 nm wavelength.

Gradient

t (min)	0	25	26	35	36	45
%B	7	7	60	60	7	7

5

Example 39

Synthesis of the In-111 Complex of the Conjugate Example 30

- To a lead shielded and crimped 2 cc autosampler vial was added 120 μ g of the conjugate of Example 30 dissolved in 240 μ L ammonium acetate buffer (0.5 M, pH 4.7) followed by the addition of 2 mg of gentisic acid (sodium salt) dissolved in 20 μ L of H₂O, and 2.3 mCi, (10 μ L) In-111 (NEN) in 0.05 N HCl (specific activity: 52 μ g/mCi). The reaction mixture was heated at 100 °C for 20 min and analyzed by HPLC. Yield: 94.7% (total for the two isomers), Ret. Time: 16.6 and 17.3 min.

HPLC Method

- Column: Zorbax Rx C18, 25 cm x 4.6 mm
 Column Temperature: Ambient
 Flow: 1.0 ml/min
 Solvent A: 10 mM ammonium acetate
 Solvent B: Acetonitrile
 Detector: IN-US β -ram, and UV at 220 nm wavelength.
 Gradient
- | | | | | | | |
|---------|----|----|----|----|----|----|
| t (min) | 0 | 25 | 26 | 35 | 36 | 45 |
| %B | 10 | 15 | 60 | 60 | 10 | 10 |

30

Example 40

Synthesis of the In-111 Complex of the Conjugate Example 31

- To a shielded and crimped 2 cc autosampler vial was added 70 μ g of the conjugate of Example 31 dissolved in 140 μ L 0.5 M ammonium acetate buffer (pH 4.8) followed by the addition of 2 mg gentisic acid sodium salt and 2.6

mCi (7 μ l) In-111 in 0.05M HCl. The reaction mixture (specific activity was heated at 85°C for 20 minutes and analyzed by HPLC. Yield: 92.2%; Ret. Time: 12.9 min.

5 HPLC Method

Column: Zorbax Rx C18, 25 cm x 4.6 mm

Column Temperature: Ambient

Flow: 1.0 ml/min

Solvent A: 10 mM ammonium acetate

10 Solvent B: Acetonitrile

Detector: IN-US β -ram, and UV at 220 nm wavelength.

Gradient

t (min)	0	25	26	35	36	45
%B	7	7	60	60	7	7

15

Example 41

Synthesis of the In-111 Complex of the Conjugate Example 33

- 20 To a shielded and crimped 2 cc autosampler vial was added 107 μ g of the conjugate of Example 33 dissolved in 140 μ l 0.5 M ammonium acetate buffer (pH 4.8) followed by the addition of 2 mg gentisic acid sodium salt and 2.6 mCi (7 μ l) In-111 in 0.05M HCl. The reaction mixture
- 25 (specific activity was heated at 85°C for 20 minutes and analyzed by HPLC. Yield: 77.9%; Ret. Time: 17.8 min.

HPLC Method

Column: Zorbax Rx C18, 25 cm x 4.6 mm

30 Column Temperature: Ambient

Flow: 1.0 ml/min

Solvent A: 10 mM ammonium acetate

Solvent B: Acetonitrile

Detector: IN-US β -ram, and UV at 220 nm wavelength.

35 Gradient

t (min)	0	25	26	35	36	45
%B	9	11	60	60	9	9

Example 42

Synthesis of the In-111 Complex of the Conjugate Example

5 34

To a lead shielded and crimped autosampler vial was added 25 µg of the conjugate of Example 34 and 1.0 mg gentisic acid, sodium salt dissolved in 50 µL ammonium acetate buffer (0.4 M, pH 4.7) followed by the addition
10 of 1.2 mCi, (5 µL) In-111 in 0.05 N HCl (specific activity: 21 µg/mCi). The reaction mixture was heated at 80 °C for 45 min and analyzed by HPLC and ITLC. 93.5% yield by HPLC, Ret. Time: 16.7 min.

15 HPLC Method

Column: Zorbax Rx C18, 25 cm x 4.6 mm

Column Temperature: Ambient

Flow: 1.0 ml/min

Solvent A: 25 mM sodium phosphate buffer at pH 6

20 Solvent B: Acetonitrile

Detector: Sodium iodide (NaI) radiometric probe, and UV at 220 nm wavelength.

Gradient

t (min)	0	25	26	35	36	45
25 %B	10	20	60	60	10	10

Example 43

Synthesis of the In-111 Complex of the Conjugate Example

35

30 To a lead shielded and crimped 1 cc autosampler vial was added 40-50 µg of the conjugate of Example 35 dissolved in 100 µL ammonium citrate buffer (0.4 M, pH 4.7) followed by the addition of 2 mCi, (5 µL) In-111 in 0.05 N HCl (specific activity: 25 µg/mCi). The reaction
35 mixture was heated at 90-100 C for 30 min and analyzed by HPLC. Yield: 95%; Ret. Time 12.5 min.

HPLC Method

Column: Zorbax Rx C18, 25 cm x 4.6 mm

Column Temperature: Ambient

Flow: 1.0 ml/min

5 Solvent A: 25 mM sodium phosphate buffer at pH 6

Solvent B: Acetonitrile

Detector: Sodium iodide (NaI) radiometric probe, and UV
at 220 nm wavelength.

Gradient

10	t (min)	0	25	26	35	36	45
	%B	10	20	60	60	10	10

Example 4415 Synthesis of the In-111 Complex of the Conjugate Example
37

To a lead shielded and crimped 2 cc autosampler vial was added 150 µg of the conjugate of Example 37 dissolved in 300 µL ammonium citrate buffer (0.3 M, pH 4.8) followed by the addition of 4.5 mCi, (25 µL) In-111 (NEN) in 0.05 N HCl (specific activity: 33 µg/mCi). The reaction mixture was heated at 100 °C for 20 min and analyzed by HPLC. RCP: 80%, Ret. Time: 21 min.

HPLC Method

25 Column: Zorbax Rx C18, 25 cm x 4.6 mm

Column Temperature: Ambient

Flow: 1.0 ml/min

Solvent A: 25 mM sodium phosphate buffer at pH 6

Solvent B: Acetonitrile

30 Detector: Sodium iodide (NaI) radiometric probe, and UV
at 220 nm wavelength.

Gradient

35	t (min)	0	25	26	35	36	45
	%B	17	19	60	60	17	17

Examples 45 - 51

Synthesis of Y-90 and Lu-177 Complexes of the Conjugates of Examples 30, 31, 34, 35 and 37

- To a clean sealed 5 mL vial was added 0.5 -1.0 mL of the appropriate conjugate solution (200 µg/mL in 0.5 M ammonium acetate buffer, pH 7.0-8.0), followed by 0.05 mL of sodium gentisate (10 mg/mL in 0.5 M ammonium acetate buffer, pH 7.0-8.0) solution, and 10 - 40 µL of ⁹⁰YCl₃ or ¹⁷⁷LuCl₃ solution (10 - 20 mCi) in 0.05 N HCl. The reaction mixture was heated at 100 °C for 5-10 min. After cooling to room temperature, a sample of the resulting solution was analyzed by HPLC and by ITLC.

Complex Ex #	Isotope	Conjugate Ex. #	Ret. Time (min)	% Yield	HPLC Method
45	Y-90	30	14.0, 16.0	90	D
46	Y-90	31	14.0	90.5	F
47	Lu-177	31	13.0	85	D
48	Y-90	34	8.0	81.9	A
49	Y-90	35	16.0	89	B
50	Y-90	37	8.2	83.5	B
51	Lu-177	37	14.0	70	G

- HPLC Method A: The HPLC method using a reverse phase C₁₈ Zorbax column (4.6 mm x 25 cm, 80 Å pore size) at a flow rate of 1.0 mL/min with a gradient mobile phase from 85% A (25 mM pH 6.0 phosphate buffer) and 15% B (acetonitrile) to 75% A and 25% B at 20 min.

20

HPLC Method B: The HPLC method using a reverse phase C₁₈ Zorbax column (4.6 mm x 25 cm, 80 Å pore size) at a flow rate of 1.0 mL/min with a gradient mobile phase from 90% A (25 mM pH 6.0 phosphate buffer) and 10% B

- (acetonitrile) to 80% A and 20% B at 20 min.

- HPLC Method D: The HPLC method using a reverse phase C₁₈ Zorbax column (4.6 mm x 25 cm, 80 Å pore size) at a flow rate of 1.0 mL/min with a gradient mobile phase from 87% A (25 mM pH 6.0 phosphate buffer) and 13% B (acetonitrile) to 86% A and 14% B at 20 min.

HPLC Method F: The HPLC method using a reverse phase C₁₈ Zorbax column (4.6 mm x 25 cm, 80 Å pore size) at a flow rate of 1.0 mL/min with a gradient mobile phase from 92% A (25 mM ammonium acetate buffer, pH = 6.8) and 8% B (acetonitrile) to 90% A and 10% B at 20 min.

HPLC Method G: The HPLC method using a reverse phase C₁₈ Zorbax column (4.6 mm x 25 cm, 80 Å pore size) at a flow rate of 1.0 mL/min with an isocratic mobile phase of 87% A (25 mM ammonium acetate buffer, pH = 6.8) and 13% B (acetonitrile) from 0 to 20 min.

Example 52

15 Synthesis of ^{99m}Tc(3-[[1-(3-{2-[(6-(diazenido)(3-pyridyl))carbonylamino](2R)-3-sulfopropyl)propyl]-7-[(imidazol-2-ylamino)methyl]-4-oxo(3-hydroquinolyl)]carbonylamino}(2S)-2-[[2,4,6-trimethylphenyl)sulfonyl]amino}propanoic acid)
20 (tricine) (TPPTS)

To a lyophilized vial containing 4.84 mg TPPTS, 6.3 mg tricine, 40 mg mannitol, succinic acid buffer, pH 4.8, and 0.1% Pluronic F-64 surfactant, was added 1.1 mL sterile water for injection, 0.2 mL (20 µg) of the the
25 conjugate of Example 36 in deionized water or 50% aqueous ethanol, and 0.2 mL of ^{99m}TcO₄⁻ (50±5 mCi) in saline. The reconstituted kit was heated in a 100 °C water bath for 15 minutes, and was allowed to cool 10 minutes at room temperature. A sample of the reaction mixture was
30 analyzed by HPLC. The yield was 89.0% and the retention time 12.8, 13.2 min (2 isomers).

HPLC Method

Column: Zorbax C₁₈ , 25 cm x 4.6 mm

Flow rate : 1.0 mL/min

35 Solvent A: 10 mM sodium phosphate buffer, pH 6.0

Solvent B : 100 % CH₃CN

Gradient 0 - 25% B over 20 min.

Utility

The pharmaceuticals of the present invention are useful for imaging angiogenic tumor vasculature, therapeutic cardiovascular angiogenesis, and cardiac

5 pathologies associated with the expression of vitronectin receptors in a patient or for treating cancer in a patient. The radiopharmaceuticals of the present invention comprised of a gamma ray or positron emitting isotope are useful for imaging of pathological processes

10 involving angiogenic neovasculature, including cancer, diabetic retinopathy, macular degeneration, restenosis of blood vessels after angioplasty, and wound healing, as well as atherosclerotic plaque, myocardial reperfusion injury, and myocardial ischemia, stunning or infarction.

15 The radiopharmaceuticals of the present invention comprised of a beta, alpha or Auger electron emitting isotope are useful for treatment of pathological processes involving angiogenic neovasculature, by delivering a cytotoxic dose of radiation to the locus of

20 the angiogenic neovasculature. The treatment of cancer is affected by the systemic administration of the radiopharmaceuticals resulting in a cytotoxic radiation dose to tumors.

The compounds of the present invention comprised of

25 one or more paramagnetic metal ions selected from gadolinium, dysprosium, iron, and manganese, are useful as contrast agents for magnetic resonance imaging (MRI) of pathological processes involving angiogenic neovasculature, as well as atherosclerotic plaque,

30 myocardial reperfusion injury, and myocardial ischemia, stunning or infarction.

The compounds of the present invention comprised of one or more heavy atoms with atomic number of 20 or greater are useful as X-ray contrast agents for X-ray

35 imaging of pathological processes involving angiogenic neovasculature, as well as atherosclerotic plaque, myocardial reperfusion injury, and myocardial ischemia, stunning or infarction.

The compounds of the present invention comprised of an echogenic gas containing surfactant microsphere are useful as ultrasound contrast agents for sonography of pathological processes involving angiogenic
5 neovasculature, as well as atherosclerotic plaque, myocardial reperfusion injury, and myocardial ischemia, stunning or infarction.

Representative compounds of the present invention
10 were tested in the following in vitro assays and in vivo models and were found to be active.

Immobilized Human Placental avb3 Receptor Assay

The assay conditions were developed and validated
15 using [I-125]vitronectin. Assay validation included Scatchard format analysis (n=3) where receptor number (Bmax) and Kd (affinity) were determined. Assay format is such that compounds are preliminarily screened at 10 and 100 nM final concentrations prior to IC50
20 determination. Three standards (vitronectin, anti-avB3 antibody, LM609, and anti-avB5, P1F6) and five reference peptides have been evaluated for IC50 determination. Briefly, the method involves immobilizing previously isolated receptors in 96 well plates and incubating
25 overnight. The receptors were isolated from normal, fresh, non-infectious (HIV, hepatitis B and C, syphilis, and HTLV free) human placenta. The tissue was lysed and tissue debris removed via centrifugation. The lysate was filtered. The receptors were isolated by affinity
30 chromatography using the immobilized avb3 antibody. The plates are then washed 3x with wash buffer. Blocking buffer is added and plates incubated for 120 minutes at room temperature. During this time compounds to be tested and [I-125]vitronectin are premixed in a reservoir
35 plate. Blocking buffer is removed and compound mixture pipetted. Competition is carried out for 60 minutes at room temperature. Unbound material is then removed and wells are separated and counted via gamma scintillation.

Oncomouse® Imaging

The study involves the use of the c-Neu Oncomouse® and FVB mice simultaneously as controls. The mice are anesthetized with sodium pentobarbital and injected with approximately 0.5 mCi of radiopharmaceutical. Prior to injection, the tumor locations on each Oncomouse® are recorded and tumor size measured using calipers. The animals are positioned on the camera head so as to image the anterior or posterior of the animals. 5 Minute dynamic images are acquired serially over 2 hours using a 256x256 matrix and a zoom of 2x. Upon completion of the study, the images are evaluated by circumscribing the tumor as the target region of interest (ROI) and a background site in the neck area below the carotid salivary glands.

This model can also be used to assess the effectiveness of the radiopharmaceuticals of the present invention comprised of a beta, alpha or Auger electron emitting isotope. The radiopharmaceuticals are administered in appropriate amounts and the uptake in the tumors can be quantified either non-invasively by imaging for those isotopes with a coincident imageable gamma emission, or by excision of the tumors and counting the amount of radioactivity present by standard techniques. The therapeutic effect of the radiopharmaceuticals can be assessed by monitoring the rate of growth of the tumors in control mice versus those in the mice administered the radiopharmaceuticals of the present invention.

This model can also be used to assess the compounds of the present invention comprised of paramagnetic metals as MRI contrast agents. After administration of the appropriate amount of the paramagnetic compounds, the whole animal can be placed in a commercially available magnetic resonance imager to image the tumors. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

This model can also be used to assess the compounds of the present invention comprised of heavy atoms as X-

ray contrast agents. After administration of the appropriate amount of the X-ray absorbing compounds, the whole animal can be placed in a commercially available X-ray imager to image the tumors. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

This model can also be used to assess the compounds of the present invention comprised of an echogenic gas containing surfactant microsphere as ultrasound contrast agents. After administration of the appropriate amount of the echogenic compounds, the tumors in the animal can be imaging using an ultrasound probe held proximate to the tumors. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

Rabbit Matrigel Model

This model was adapted from a matrigel model intended for the study of angiogenesis in mice. Matrigel (Becton & Dickinson, USA) is a basement membrane rich in laminin, collagen IV, entactin, HSPG and other growth factors. When combined with growth factors such as bFGF [500 ng/ml] or VEGF [2 µg/ml] and injected subcutaneously into the mid-abdominal region of the mice, it solidifies into a gel and stimulates angiogenesis at the site of injection within 4-8 days. In the rabbit model, New Zealand White rabbits (2.5-3.0 kg) are injected with 2.0 ml of matrigel, plus 1 µg bFGF and 4 µg VEGF. The radiopharmaceutical is then injected 7 days later and the images obtained.

This model can also be used to assess the effectiveness of the radiopharmaceuticals of the present invention comprised of a beta, alpha or Auger electron emitting isotope. The radiopharmaceuticals are administered in appropriate amounts and the uptake at the angiogenic sites can be quantified either non-invasively by imaging for those isotopes with a coincident imageable gamma emission, or by excision of the angiogenic sites

and counting the amount of radioactivity present by standard techniques. The therapeutic effect of the radiopharmaceuticals can be assessed by monitoring the rate of growth of the angiogenic sites in control rabbits
5 versus those in the rabbits administered the radiopharmaceuticals of the present invention.

This model can also be used to assess the compounds of the present invention comprised of paramagnetic metals as MRI contrast agents. After administration of the
10 appropriate amount of the paramagnetic compounds, the whole animal can be placed in a commercially available magnetic resonance imager to image the angiogenic sites. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that
15 are not administered a contrast agent.

This model can also be used to assess the compounds of the present invention comprised of heavy atoms as X-ray contrast agents. After administration of the appropriate amount of the X-ray absorbing compounds, the
20 whole animal can be placed in a commercially available X-ray imager to image the angiogenic sites. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

25 This model can also be used to assess the compounds of the present invention comprised of an echogenic gas containing surfactant microsphere as ultrasound contrast agents. After administration of the appropriate amount of the echogenic compounds, the angiogenic sites in the
30 animal can be imaging using an ultrasound probe held proximate to the tumors. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

35

Canine Spontaneous Tumor Model

Adult dogs with spontaneous mammary tumors were sedated with xylazine (20 mg/kg)/atropine (1 ml/kg). Upon sedation the animals were intubated using ketamine

(5 mg/kg)/diazepam (0.25 mg/kg) for full anesthesia. Chemical restraint was continued with ketamine (3 mg/kg)/xylazine (6 mg/kg) titrating as necessary. If required the animals were ventilated with room air via an endotracheal tube (12 strokes/min, 25 ml/kg) during the study. Peripheral veins were catheterized using 20G I.V. catheters, one to serve as an infusion port for compound while the other for exfusion of blood samples. Heart rate and EKG were monitored using a cardiometer (Biotech, Grass Quincy, MA) triggered from a lead II electrocardiogram generated by limb leads. Blood samples are generally taken at ~10 minutes (control), end of infusion, (1 minute), 15 min, 30 min, 60 min, 90 min, and 120 min for whole blood cell number and counting. Radiopharmaceutical dose was 300 μ Ci/kg administered as an i.v. bolus with saline flush. Parameters were monitored continuously on a polygraph recorder (Model 7E Grass) at a paper speed of 10 mm/min or 10 mm/sec.

Imaging of the laterals were for 2 hours with a 256x256 matrix, no zoom, 5 minute dynamic images. A known source is placed in the image field (20-90 μ Ci) to evaluate region of interest (ROI) uptake. Images were also acquired 24 hours post injection to determine retention of the compound in the tumor. The uptake is determined by taking the fraction of the total counts in an inscribed area for ROI/source and multiplying the known μ Ci. The result is μ Ci for the ROI.

This model can also be used to assess the effectiveness of the radiopharmaceuticals of the present invention comprised of a beta, alpha or Auger electron emitting isotope. The radiopharmaceuticals are administered in appropriate amounts and the uptake in the tumors can be quantified either non-invasively by imaging for those isotopes with a coincident imageable gamma emission, or by excision of the tumors and counting the amount of radioactivity present by standard techniques. The therapeutic effect of the radiopharmaceuticals can be assessed by monitoring the size of the tumors over time.

This model can also be used to assess the compounds

of the present invention comprised of paramagnetic metals as MRI contrast agents. After administration of the appropriate amount of the paramagnetic compounds, the whole animal can be placed in a commercially available magnetic resonance imager to image the tumors. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

This model can also be used to assess the compounds of the present invention comprised of heavy atoms as X-ray contrast agents. After administration of the appropriate amount of the X-ray absorbing compounds, the whole animal can be placed in a commercially available X-ray imager to image the tumors. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

This model can also be used to assess the compounds of the present invention comprised of an echogenic gas containing surfactant microsphere as ultrasound contrast agents. After administration of the appropriate amount of the echogenic compounds, the tumors in the animal can be imaging using an ultrasound probe held proximate to the tumors. The effectiveness of the contrast agents can be readily seen by comparison to the images obtain from animals that are not administered a contrast agent.

Cardiovascular disease models that can be used to assess the diagnostic radiopharmaceuticals, magnetic resonance, X-ray and ultrasound contrast agents of the present invention are reviewed in J. Nucl. Cardiol., 1998, 5, 167-83. There are several well established rabbit models of atherosclerosis; one model produces predominantly proliferating smooth muscle cells by balloon deendothelialization of infradiaphragmatic abdominal aorta to simulate restenotic lesions; another model that produces simulated advanced human atherosclerotic plaque by balloon deendothelialization followed by a high cholesterol diet.

A model of congestive heart failure is described in Am. J. Physiol., 1998, 274, H1516-23. In general, Yorkshire pigs are randomly assigned to undergo 3 wks of rapid atrial pacing at 240 beats/min. or to be sham controls.

- 5 The pigs are chronically instrumented to measure left ventricular function in the conscious state. The pigs are anesthetized.


- A shielded stimulating electrode is sutured onto the left atrium, connected to a modified programmable pace maker
10 and buried in a subcutaneous pocket. The pericardium is closed loosely, the thoracotomy is closed, and the pleural space is evacuated of air. After a recovery period of 7-10 days, the pacemaker is activated in the animals selected to undergo chronic rapid pacing. The
15 animals are sedated, the pacemaker is deactivated (pacing groups only. After a 30 min stabilization period, indexes of LV function and geometry are determined (by echocardiography as a control) by injecting the radiolabeled compound. For biodistribution, the animals
20 are anesthetized, the heart extirpate and the LV apex and midventricular regions are evaluated.

A rat model of reversible coronary occlusion and reperfusion is described in McNulty et al., J. Am. Physiol., 1996, H2283-9.


- 25 Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically described
30 herein.

WHAT IS CLAIMED IS DESCRIBED BELOW:

1. A compound, comprising: a targeting moiety and a chelator, wherein the targeting moiety is bound to the chelator, is a quinolone nonpeptide, and binds to a receptor that is upregulated during angiogenesis and the compound has 0-1 linking groups between the targeting moiety and chelator.
2. A compound according to Claim 1, wherein the receptor is the integrin $\alpha_v\beta_3$ or $\alpha_v\beta_5$ and the compound is of the formula:




- 15 wherein, Q is a compound of Formula (II):



(II)

- including stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, or pharmaceutically acceptable salt or prodrug forms thereof wherein:

- 25 R^{1e} is selected from:



5

A^e is $-CH_2-$ or $-N(R^{10e})-$;

A^{1e} and B^e are independently $-CH_2-$ or $-N(R^{10e})-$;

10

D^e is $-N(R^{10e})-$ or $-S-$;

E^e-F^e is $-C(R^{2e})=C(R^{3e})-$ or $-C(R^{2e})_2C(R^{3e})_2-$;

15

J^e is $-C(R^{2e})-$ or $-N-$;

K^e , L^e and M^e are independently $-C(R^{2e})-$ or $-C(R^{3e})-$;

R^{2e} and R^{3e} are independently selected from:

H, C₁-C₄ alkoxy, NR^{11e}R^{12e}, halogen, NO₂, CN, CF₃,
 C₁-C₆ alkyl, C₃-C₆ alkenyl, C₃-C₇ cycloalkyl,
 C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl(C₁-C₆ alkyl)-,
 (C₁-C₆ alkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl,
 5 arylcarbonyl, and aryl substituted with 0-4 R^{7e},

alternatively, when R^{2e} and R^{3e} are substituents on
 adjacent atoms, they can be taken together with the
 carbon atoms to which they are attached to form a 5-
 10 7 membered carbocyclic or 5-7 membered heterocyclic
 aromatic or nonaromatic ring system, said
 carbocyclic or heterocyclic ring being substituted
 with 0-2 groups selected from C₁-C₄ alkyl, C₁-C₄
 alkoxy, halo, cyano, amino, CF₃ and NO₂;

15

R^{2ae} is selected from:

H, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl,
 C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl, aryl(C₁-C₄
 alkyl)-, (C₂-C₇ alkyl)carbonyl, arylcarbonyl,
 20 (C₂-C₁₀ alkoxy)carbonyl, C₃-C₇ cycloalkoxycarbonyl,
 C₇-C₁₁ bicycloalkoxycarbonyl, aryloxycarbonyl,
 aryl(C₁-C₁₀ alkoxy)carbonyl,
 C₁-C₆ alkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl,
 arylcarbonyloxy(C₁-C₄ alkoxy)carbonyl, and
 25 C₃-C₇ cycloalkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl;

R^{7e} is selected from:

H, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, aryl, aryl(C₁-
 C₄ alkyl)-, (C₁-C₄ alkyl)carbonyl, CO₂R^{18ae}, SO₂R^{11e},
 30 SO₂NR^{10e}R^{11e}, OR^{10e}, and N(R^{11e})R^{12e};

U^e is selected from:

-(CH₂)_n^e-, -(CH₂)_n^eO(CH₂)_m^e-, -(CH₂)_n^eN(R¹²)(CH₂)_m^e-,
 -NH(CH₂)_n^e-, -(CH₂)_n^eC(=O)(CH₂)_m^e-,
 35 -(CH₂)_n^eS(O)_p^e(CH₂)_m^e-, -(CH₂)_n^eNHNH(CH₂)_m^e-,
 -N(R^{10e})C(=O)-, -NHC(=O)(CH₂)_n^e-, -C(=O)N(R^{10e})-, and
 -N(R^{10e})S(O)_p^e-;

G^e is N or CR^{19e};

W^e is -C(=O)-N(R^{10e})-(C₁-C₃ alkylene)-, in which the
alkylene group is substituted by R^{8e} and by R^{9e}:

5

R^{8e} and R^{9e} are independently selected from:

- H, CO₂R^{18be}, C(=O)R^{18be}, CONR¹⁷R^{18be},
- C₁-C₁₀ alkyl substituted with 0-1 R^{6e},
- C₂-C₁₀ alkenyl substituted with 0-1 R^{6e},
- 10 C₂-C₁₀ alkynyl substituted with 0-1 R^{6e},
- C₃-C₈ cycloalkyl substituted with 0-1 R^{6e},
- C₅-C₆ cycloalkenyl substituted with 0-1 R^{6e},
- (C₁-C₁₀ alkyl)carbonyl,
- C₃-C₁₀ cycloalkyl(C₁-C₄ alkyl)-,
- 15 phenyl substituted with 0-3 R^{6e},
- naphthyl substituted with 0-3 R^{6e},
- a 5-10 membered heterocyclic ring containing 1-3 N,
O, or S heteroatoms, wherein said heterocyclic
ring may be saturated, partially saturated, or
- 20 fully unsaturated, said heterocyclic ring being
substituted with 0-2 R^{7e},
- C₁-C₁₀ alkoxy substituted with 0-2 R^{7e},
- hydroxy, nitro, -N(R^{10e})R^{11e}, -N(R^{16e})R^{17e}, aryl(C₀-C₆
alkyl)carbonyl, aryl(C₃-C₆ alkyl),
- 25 heteroaryl(C₁-C₆ alkyl), CONR^{18ae}R^{20e}, SO₂R^{18ae},
and SO₂NR^{18ae}R^{20e},
- providing that any of the above alkyl, cycloalkyl,
aryl or heteroaryl groups may be unsubstituted
or substituted independently with 1-2 R^{7e};

30

R^{6e} is selected from:

- H, C₁-C₁₀ alkyl, hydroxy, C₁-C₁₀ alkoxy, nitro, C₁-C₁₀
alkylcarbonyl, -N(R^{11e})R^{12e}, cyano, halo, CF₃,
- CHO, CO₂R^{18be}, C(=O)R^{18be}, CONR^{17e}R^{18be},
- 35 OC(=O)R^{10e}, OR^{10e}, OC(=O)NR^{10e}R^{11e},
- NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},
- NR^{10e}C(=O)NR^{10e}R^{11e}, NR^{10e}SO₂NR^{10e}R^{11e},
- NR^{10e}SO₂R^{21e}, S(O)_pR^{11e}, SO₂NR^{10e}R^{11e},

aryl substituted with 0-3 groups selected from
halogen, C₁-C₆ alkoxy, C₁-C₆ alkyl, CF₃,
S(O)_m^eMe, and -NMe₂,
aryl(C₁-C₄ alkyl)-, said aryl being substituted with
5 0-3 groups selected from halogen, C₁-C₆ alkoxy,
C₁-C₆ alkyl, CF₃, S(O)_p^eMe, and -NMe₂, and
a 5-10 membered heterocyclic ring containing 1-3 N, O, or
S heteroatoms, wherein said heterocyclic ring may be
saturated, partially saturated, or fully unsaturated,
10 said heterocyclic ring being substituted with 0-2 R^{7e};

R^{10e} is selected from:
H, CF₃, C₃-C₆ alkenyl, C₃-C₁₁ cycloalkyl, aryl,
(C₃-C₁₁ cycloalkyl)methyl, aryl(C₁-C₄ alkyl), and C₁-
15 C₁₀ alkyl substituted with 0-2 R^{6e};

R^{11e} is selected from:
H, hydroxy, C₁-C₈ alkyl, C₃-C₆ alkenyl, C₃-C₁₁
cycloalkyl, (C₃-C₁₁ cycloalkyl)methyl, C₁-C₆ alkoxy,
20 benzyloxy, aryl, heteroaryl, heteroaryl(C₁-C₄ alkyl)-
, aryl(C₁-C₄ alkyl), adamantylmethyl, and
C₁-C₁₀ alkyl substituted with 0-2 R^{4e};

R^{4e} is selected from:
25 H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇
cycloalkyl(C₁-C₄ alkyl)-, (C₁-C₁₀ alkyl)carbonyl,
aryl, heteroaryl, aryl(C₁-C₆ alkyl)-, and
heteroaryl(C₁-C₆ alkyl)-, wherein said aryl or
heteroaryl groups are substituted with 0-2
30 substituents independently selected from the group
consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, F, Cl, Br,
CF₃, and NO₂,

alternatively, when R^{10e} and R^{11e} are both substituents on
35 the same nitrogen atom (as in -NR^{10e}R^{11e}) they may be
taken together with the nitrogen atom to which they
are attached to form a heterocycle selected from:
3-azabicyclononyl, 1,2,3,4-tetrahydro-1-quinolinyl,

1,2,3,4-tetrahydro-2-isoquinolinyl, 1-piperidinyl, 1-morpholinyl, 1-pyrrolidinyl, thiamorpholinyl, thiazolidinyl, and 1-piperazinyl;

5 said heterocycle being substituted with 0-3 groups selected from: C₁-C₆ alkyl, aryl, heteroaryl, aryl(C₁-C₄ alkyl)-, (C₁-C₆ alkyl)carbonyl, (C₃-C₇ cycloalkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl, aryl(C₁-C₄ alkoxy)carbonyl, C₁-C₆ alkylsulfonyl, and arylsulfonyl;

10

R^{12e} is selected from:

H, C₁-C₆ alkyl, triphenylmethyl, methoxymethyl, methoxyphenyldiphenylmethyl, trimethylsilylethoxymethyl, (C₁-C₆ alkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl, (C₁-C₆ alkyl)aminocarbonyl, 15 C₃-C₆ alkenyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-, aryl, heteroaryl(C₁-C₆ alkyl)carbonyl, heteroarylcarbonyl, aryl(C₁-C₆ alkyl)-, (C₁-C₆ alkyl)carbonyl, arylcarbonyl, C₁-C₆ 20 alkylsulfonyl, arylsulfonyl, aryl(C₁-C₆ alkyl)sulfonyl, heteroarylsulfonyl, heteroaryl(C₁-C₆ alkyl)sulfonyl, aryloxy carbonyl, and aryl(C₁-C₆ alkoxy)carbonyl, wherein said aryl groups are substituted with 0-2 substituents selected from the 25 group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, halo, CF₃, and nitro;

R^{16e} is selected from:

-C(=O)OR^{18ae}, -C(=O)R^{18be}, -C(=O)N(R^{18be})₂,
30 -C(=O)NHSO₂R^{18ae}, -C(=O)NHC(=O)R^{18be},
-C(=O)NHC(=O)OR^{18ae}, -C(=O)NHSO₂NHR^{18be}, -SO₂R^{18ae},
-SO₂N(R^{18be})₂, and -SO₂NHC(=O)OR^{18be};

R^{17e} is selected from:

35 H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-, aryl, aryl(C₁-C₆ alkyl)-, and heteroaryl(C₁-C₆ alkyl);

R^{18ae} is selected from:

5 C₁-C₈ alkyl optionally substituted with a bond to L_n,
 C₃-C₁₁ cycloalkyl optionally substituted with a
 bond to L_n, aryl(C₁-C₆ alkyl)- optionally
 10 substituted with a bond to L_n, heteroaryl(C₁-C₆
 alkyl)- optionally substituted with a bond to
 L_n, (C₁-C₆ alkyl)heteroaryl optionally
 substituted with a bond to L_n, biaryl(C₁-C₆
 15 alkyl) optionally substituted with a bond to
 L_n, heteroaryl optionally substituted with a
 bond to L_n, phenyl substituted with 3-4 R^{19e} and
 optionally substituted with a bond to L_n,
 naphthyl substituted with 0-4 R^{19e} and
 optionally substituted with a bond to L_n, and a
 bond to L_n, wherein said aryl or heteroaryl
 groups are optionally substituted with 0-4 R^{19e};

R^{18be} is H or R^{18ae};

20 R^{19e} is selected from:

H, halogen, CF₃, CO₂H, CN, NO₂, -NR^{11e}R^{12e}, OCF₃,
 C₁-C₈ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl,
 C₃-C₁₁ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-,
 25 aryl(C₁-C₆ alkyl)-, C₁-C₆ alkoxy, C₁-C₄
 alkoxycarbonyl, aryl, aryl-O-, aryl-SO₂-,
 heteroaryl, and heteroaryl-SO₂-, wherein said aryl
 and heteroaryl groups are substituted with 0-4
 groups selected from hydrogen, halogen, CF₃, C₁-C₃
 alkyl, and C₁-C₃ alkoxy;

30

R^{20e} is selected from:

hydroxy, C₁-C₁₀ alkyloxy, C₃-C₁₁ cycloalkyloxy,
 aryloxy, aryl(C₁-C₄ alkyl)oxy,
 C₂-C₁₀ alkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 35 C₂-C₁₀ alkoxycarbonyloxy(C₁-C₂ alkyl)oxy-,
 C₂-C₁₀ alkoxycarbonyl(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkoxycarbonyloxy(C₁-C₂ alkyl)oxy-,

- C₃-C₁₀ cycloalkoxycarbonyl(C₁-C₂ alkyl)oxy-,
 aryloxy carbonyl(C₁-C₂ alkyl)oxy-,
 aryloxy carbonyloxy(C₁-C₂ alkyl)oxy-,
 aryl carbonyloxy(C₁-C₂ alkyl)oxy-,
 5 C₁-C₅ alkoxy(C₁-C₅ alkyl) carbonyloxy(C₁-C₂ alkyl)oxy,
 (5-(C₁-C₅ alkyl)-1,3-dioxo-cyclopenten-2-one-
 yl)methyloxy,
 (5-aryl-1,3-dioxo-cyclopenten-2-one-yl)methyloxy,
 and
 10 (R^{10e}) (R^{11e})N-(C₁-C₁₀ alkoxy)-;

R^{21e} is selected from:


- C₁-C₈ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl, (C₃-C₁₁
 cycloalkyl)methyl, aryl, aryl(C₁-C₄ alkyl)-, and C₁-
 15 C₁₀ alkyl substituted with 0-2 R^{7e};

R^{22e} is selected from:

- C(=O)-R^{18be}, -C(=O)N(R^{18be})₂, -C(=O)NHSO₂R^{18ae}, -
 C(=O)NHC(=O)R^{18be}, -C(=O)NHC(=O)OR^{18ae}, and -
 20 C(=O)NHSO₂NHR^{18be};

Y^e is selected from:

- COR^{20e}, -SO₃H, -PO₃H, -CONHNHSO₂CF₃, -CONHSO₂R^{18ae},
 -CONHSO₂NHR^{18be}, -NHCOCF₃, -NHCONHSO₂R^{18ae},
 25 -NHSO₂R^{18ae}, -OPO₃H₂, -OSO₃H, -PO₃H₂, -SO₂NHCOR^{18ae},
 -SO₂NHCO₂R^{18ae},



- 30 m^e is 0-2;

n^e is 0-4;

p^e is 0-2;

35

r^e is 0-2;


with the following proviso: n^e and m^e are chosen such that the number of atoms connecting R^{1e} and Y^e is in the range of 8-14;

5

d is selected from 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

d' is 1-100;

10 L_n is a linking group having the formula:



15 W is independently selected at each occurrence from the group: O, S, NH, NHC(=O), C(=O)NH, $NR^8C(=O)$, C(=O)N R^8 , C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, SO₂NH, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'}, (OCH₂CH₂CH₂)_{s''}, (CH₂CH₂CH₂O)_t, and (aa)_{t'};

20 aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-3 R^{10} , C₃₋₁₀ cycloalkyl substituted with 0-3 R^{10} , and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R^{10} ;

25 R^6 , R^{6a} , R^7 , R^{7a} , and R^8 are independently selected at each occurrence from the group: H, =O, COOH, SO₃H, PO₃H, C₁₋₅ alkyl substituted with 0-3 R^{10} , aryl substituted with 0-3 R^{10} , benzyl substituted with 0-3 R^{10} , and C₁₋₅ alkoxy substituted with 0-3 R^{10} , NHC(=O) R^{11} , C(=O)NHR¹¹, NHC(=O)NHR¹¹, NHR¹¹, R^{11} , and a bond to C_h ;

35

R^{10} is independently selected at each occurrence from the group: a bond to C_h , COOR¹¹, C(=O)NHR¹¹, NHC(=O) R^{11} , OH, NHR¹¹, SO₃H, PO₃H, -OPO₃H₂, -OSO₃H, aryl


substituted with 0-3 R¹¹, C₁₋₅ alkyl substituted with
0-1 R¹², C₁₋₅ alkoxy substituted with 0-1 R¹², and a
5-10 membered heterocyclic ring system containing
1-4 heteroatoms independently selected from N, S,
5 and O and substituted with 0-3 R¹¹;

R¹¹ is independently selected at each occurrence from the
group: H, alkyl substituted with 0-1 R¹², aryl
substituted with 0-1 R¹², a 5-10 membered
10 heterocyclic ring system containing 1-4 heteroatoms
independently selected from N, S, and O and
substituted with 0-1 R¹², C₃₋₁₀ cycloalkyl
substituted with 0-1 R¹², polyalkylene glycol
substituted with 0-1 R¹², carbohydrate substituted
15 with 0-1 R¹², cyclodextrin substituted with 0-1 R¹²,
amino acid substituted with 0-1 R¹², polycarboxyalkyl
substituted with 0-1 R¹², polyazaalkyl substituted
with 0-1 R¹², peptide substituted with 0-1 R¹²,
wherein the peptide is comprised of 2-10 amino
20 acids, 3,6-O-disulfo-B-D-galactopyranosyl,
bis(phosphonomethyl)glycine, and a bond to C_H;

R¹² is a bond to C_H;

25 k is selected from 0, 1, and 2;
h is selected from 0, 1, and 2;
h' is selected from 0, 1, and 2;
g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
30 s is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
s' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
s" is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
t is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
35 x is selected from 0, 1, 2, 3, 4, and 5;
x' is selected from 0, 1, 2, 3, 4, and 5;

C_h is a metal bonding unit having a formula selected from the group:



10 $A^1, A^2, A^3, A^4, A^5, A^6, A^7,$ and A^8 are independently selected at each occurrence from the group: $NR^{13}, NR^{13}R^{14}, S, SH, S(Pg), O, OH, PR^{13}, PR^{13}R^{14}, P(O)R^{15}R^{16},$ and a bond to L_n ;

15 E is a bond, CH , or a spacer group independently selected at each occurrence from the group: C_1 - C_{10} alkyl substituted with 0-3 R^{17} , aryl substituted with 0-3 R^{17} , C_3 - C_{10} cycloalkyl substituted with 0-3 R^{17} , heterocyclo- C_1 - C_{10} alkyl substituted with 0-3 R^{17} , wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from $N, S,$ and O , C_6 - C_{10} aryl- C_1 - C_{10} alkyl substituted with 0-3 R^{17} , C_1 - C_{10} alkyl- C_6 - C_{10} aryl- substituted with 0-3 R^{17} , and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from $N, S,$ and O and substituted with 0-3 R^{17} ;

20

25

R^{13} and R^{14} are each independently selected from the group: a bond to L_n , hydrogen, C_1 - C_{10} alkyl substituted with 0-3 R^{17} , aryl substituted with 0-3

R^{17} , C_{1-10} cycloalkyl substituted with 0-3 R^{17} ,
 heterocyclo- C_{1-10} alkyl substituted with 0-3 R^{17} ,
 wherein the heterocyclo group is a 5-10 membered
 heterocyclic ring system containing 1-4 heteroatoms
 5 independently selected from N, S, and O, C_{6-10}
 aryl- C_{1-10} alkyl substituted with 0-3 R^{17} , C_{1-10}
 alkyl- C_{6-10} aryl- substituted with 0-3 R^{17} , a 5-10
 membered heterocyclic ring system containing 1-4
 heteroatoms independently selected from N, S, and O
 10 and substituted with 0-3 R^{17} , and an electron,
 provided that when one of R^{13} or R^{14} is an electron,
 then the other is also an electron;

alternatively, R^{13} and R^{14} combine to form $=C(R^{20})(R^{21})$;

15 R^{15} and R^{16} are each independently selected from the
 group: a bond to L_n , -OH, C_{1-10} alkyl substituted
 with 0-3 R^{17} , C_{1-10} alkyl substituted with 0-3 R^{17} ,
 aryl substituted with 0-3 R^{17} , C_{3-10} cycloalkyl
 20 substituted with 0-3 R^{17} , heterocyclo- C_{1-10} alkyl
 substituted with 0-3 R^{17} , wherein the heterocyclo
 group is a 5-10 membered heterocyclic ring system
 containing 1-4 heteroatoms independently selected
 from N, S, and O, C_{6-10} aryl- C_{1-10} alkyl substituted
 25 with 0-3 R^{17} , C_{1-10} alkyl- C_{6-10} aryl- substituted with
 0-3 R^{17} , and a 5-10 membered heterocyclic ring
 system containing 1-4 heteroatoms independently
 selected from N, S, and O and substituted with 0-3
 R^{17} ;

30 R^{17} is independently selected at each occurrence from the
 group: a bond to L_n , =O, F, Cl, Br, I, -CF₃, -CN,
 -CO₂ R^{18} , -C(=O) R^{18} , -C(=O)N(R^{18})₂, -CHO, -CH₂OR¹⁸,
 -OC(=O) R^{18} , -OC(=O)OR^{18a}, -OR¹⁸, -OC(=O)N(R^{18})₂,
 35 -NR¹⁹C(=O) R^{18} , -NR¹⁹C(=O)OR^{18a}, -NR¹⁹C(=O)N(R^{18})₂,
 -NR¹⁹SO₂N(R^{18})₂, -NR¹⁹SO₂R^{18a}, -SO₃H, -SO₂R^{18a},
 -SR¹⁸, -S(=O) R^{18a} , -SO₂N(R^{18})₂, -N(R^{18})₂,
 -NHC(=S)NHR¹⁸, =NOR¹⁸, NO₂, -C(=O)NHOR¹⁸,


-C(=O)NHN¹⁸R^{18a}, -OCH₂CO₂H, 2-(1-morpholino)ethoxy, C₁-C₅ alkyl, C₂-C₄ alkenyl, C₃-C₆ cycloalkyl, C₃-C₆ cycloalkylmethyl, C₂-C₆ alkoxyalkyl, aryl substituted with 0-2 R¹⁸, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O;

R¹⁸, R^{18a}, and R¹⁹ are independently selected at each occurrence from the group: a bond to L_n, H, C₁-C₆ alkyl, phenyl, benzyl, C₁-C₆ alkoxy, halide, nitro, cyano, and trifluoromethyl;

Pg is a thiol protecting group;

R²⁰ and R²¹ are independently selected from the group: H, C₁-C₁₀ alkyl, -CN, -CO₂R²⁵, -C(=O)R²⁵, -C(=O)N(R²⁵)₂, C₂-C₁₀ 1-alkene substituted with 0-3 R²³, C₂-C₁₀ 1-alkyne substituted with 0-3 R²³, aryl substituted with 0-3 R²³, unsaturated 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R²³, and unsaturated C₃-10 carbocycle substituted with 0-3 R²³;

alternatively, R²⁰ and R²¹, taken together with the divalent carbon radical to which they are attached form:



30

R²² and R²³ are independently selected from the group: H, R²⁴, C₁-C₁₀ alkyl substituted with 0-3 R²⁴, C₂-C₁₀ alkenyl substituted with 0-3 R²⁴, C₂-C₁₀ alkynyl substituted with 0-3 R²⁴, aryl substituted with 0-3

R²⁴, a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R²⁴, and C₃-10 carbocycle substituted with 0-3 R²⁴;

5

alternatively, R²², R²³ taken together form a fused aromatic or a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O;

10


a and b indicate the positions of optional double bonds and n is 0 or 1;

R²⁴ is independently selected at each occurrence from the group: =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R²⁵, -C(=O)R²⁵, -C(=O)N(R²⁵)₂, -N(R²⁵)₃⁺, -CH₂OR²⁵, -OC(=O)R²⁵, -OC(=O)OR^{25a}, -OR²⁵, -OC(=O)N(R²⁵)₂, -NR²⁶C(=O)R²⁵, -NR²⁶C(=O)OR^{25a}, -NR²⁶C(=O)N(R²⁵)₂, -NR²⁶SO₂N(R²⁵)₂, -NR²⁶SO₂R^{25a}, -SO₃H, -SO₂R^{25a}, -SR²⁵, -S(=O)R^{25a}, -SO₂N(R²⁵)₂, -N(R²⁵)₂, =NOR²⁵, -C(=O)NHOR²⁵, -OCH₂CO₂H, and 2-(1-morpholino)ethoxy; and,

R²⁵, R^{25a}, and R²⁶ are each independently selected at each occurrence from the group: hydrogen and C₁-C₆ alkyl;

and a pharmaceutically acceptable salt thereof.

30 3. A compound according to Claim 2, wherein Q is a compound of Formula (IV):





(IV)



- 5 including stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, or pharmaceutically acceptable salt or prodrug forms thereof wherein:

R^{1e} is selected from:

10



15



- 20 R^{2e} and R^{3e} are independently selected from:
H, C₁-C₄ alkoxy, NR^{11e}R^{12e}, halogen, NO₂, CN, CF₃,
C₁-C₆ alkyl, C₃-C₆ alkenyl, C₃-C₇ cycloalkyl,
C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl(C₁-C₆ alkyl)-,
(C₁-C₆ alkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl,
25 arylcarbonyl, and aryl substituted with 0-4 R^{7e},

alternatively, when R^{2e} and R^{3e} are substituents on adjacent atoms, they can be taken together with the carbon atoms to which they are attached to form a 5-7 membered carbocyclic or 5-7 membered heterocyclic aromatic or nonaromatic ring system, said carbocyclic or heterocyclic ring being substituted with 0-2 groups selected from C_1 - C_4 alkyl, C_1 - C_4 alkoxy, halo, cyano, amino, CF_3 and NO_2 ;

10 R^{2ae} is selected from:
 H, C_1 - C_{10} alkyl, C_2 - C_6 alkenyl, C_3 - C_{11} cycloalkyl, C_3 - C_7 cycloalkyl(C_1 - C_4 alkyl), aryl, aryl(C_1 - C_4 alkyl)-, (C_2 - C_7 alkyl)carbonyl, arylcarbonyl,
 15 (C_2 - C_{10} alkoxy)carbonyl, C_3 - C_7 cycloalkoxycarbonyl, C_7 - C_{11} bicycloalkoxycarbonyl, aryloxycarbonyl, aryl(C_1 - C_{10} alkoxy)carbonyl, C_1 - C_6 alkylcarbonyloxy(C_1 - C_4 alkoxy)carbonyl, arylcarbonyloxy(C_1 - C_4 alkoxy)carbonyl, and
 20 C_3 - C_7 cycloalkylcarbonyloxy(C_1 - C_4 alkoxy)carbonyl;

R^{7e} is selected from:
 H, hydroxy, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, aryl, aryl(C_1 - C_4 alkyl)-, (C_1 - C_4 alkyl)carbonyl, CO_2R^{18ae} , SO_2R^{11e} ,
 25 $SO_2NR^{10e}R^{11e}$, OR^{10e} , and $N(R^{11e})R^{12e}$;

U^e is selected from:
 $-(CH_2)_n^e-$, $-(CH_2)_n^eO(CH_2)_m^e-$, $-NH(CH_2)_n^e-$,
 $-N(R^{10e})C(=O)-$, $-NHC(=O)(CH_2)_n^e-$, and $-C(=O)N(R^{10e})-$;

30 G^e is N or CR^{19e} ;

R^{8e} is selected from:
 H, CO_2R^{18be} , $C(=O)R^{18be}$, $CONR^{17e}R^{18be}$,
 35 C_1 - C_{10} alkyl substituted with 0-1 R^{6e} ,
 C_2 - C_{10} alkenyl substituted with 0-1 R^{6e} ,
 C_2 - C_{10} alkynyl substituted with 0-1 R^{6e} ,
 C_3 - C_8 cycloalkyl substituted with 0-1 R^{6e} ,

C₅-C₆ cycloalkenyl substituted with 0-1 R^{6e},
 (C₁-C₁₀ alkyl)carbonyl,
 C₃-C₁₀ cycloalkyl(C₁-C₄ alkyl)-,
 phenyl substituted with 0-3 R^{6e},
 5 naphthyl substituted with 0-3 R^{6e},
 a 5-10 membered heterocyclic ring containing 1-3 N,
 O, or S heteroatoms, wherein said heterocyclic
 ring may be saturated, partially saturated, or
 fully unsaturated, said heterocyclic ring being
 10 substituted with 0-2 R^{7e};

R^{9e} is selected from:

C₁-C₁₀ alkyl substituted with 0-1 R^{6e},
 C₁-C₁₀ alkoxy substituted with 0-2 R^{7e},
 15 H, nitro, N(R^{11e})R^{12e}, OC(=O)R^{10e}, OR^{10e},
 OC(=O)NR^{10e}R^{11e}, NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},
 NR^{10e}C(=O)NR^{10e}R^{11e}, NR^{10e}SO₂NR^{10e}R^{11e},
 NR^{10e}SO₂R^{21e}, hydroxy, OR^{22e}, -N(R^{10e})R^{11e}, -
 N(R^{16e})R^{17e}, aryl(C₀-C₆ alkyl)carbonyl, aryl(C₁-
 20 C₆ alkyl), heteroaryl(C₁-C₆ alkyl), CONR^{18ae}R^{20e},
 SO₂R^{18ae}, and SO₂NR^{18ae}R^{20e},

providing that any of the above alkyl, cycloalkyl,
 aryl or heteroaryl groups may be unsubstituted
 or substituted independently with 1-2 R^{7e};

R^{6e} is selected from:

H, C₁-C₁₀ alkyl, hydroxy, C₁-C₁₀ alkoxy, nitro, C₁-C₁₀
 alkylcarbonyl, -N(R^{11e})R^{12e}, cyano, halo, CF₃,
 CHO, CO₂R^{18be}, C(=O)R^{18be}, CONR^{17e}R^{18be},
 30 OC(=O)R^{10e}, OR^{10e}, OC(=O)NR^{10e}R^{11e},
 NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},
 NR^{10e}C(=O)NR^{10e}R^{11e}, NR^{10e}SO₂NR^{10e}R^{11e},
 NR^{10e}SO₂R^{21e}, S(O)_pR^{11e}, SO₂NR^{10e}R^{11e},
 aryl substituted with 0-3 groups selected from
 35 halogen, C₁-C₆ alkoxy, C₁-C₆ alkyl, CF₃,
 S(O)_mMe, and -NMe₂,

aryl(C₁-C₄ alkyl)-, said aryl being substituted with
0-3 groups selected from halogen, C₁-C₆ alkoxy,
C₁-C₆ alkyl, CF₃, S(O)_pMe, and -NMe₂, and
a 5-10 membered heterocyclic ring containing 1-3 N, O, or
5 S heteroatoms, wherein said heterocyclic ring may be
saturated, partially saturated, or fully unsaturated,
said heterocyclic ring being substituted with 0-2 R^{7e};

R^{10e} is selected from:

10 H, CF₃, C₃-C₆ alkenyl, C₃-C₁₁ cycloalkyl, aryl,
(C₃-C₁₁ cycloalkyl)methyl, aryl(C₁-C₄ alkyl), and C₁-
C₁₀ alkyl substituted with 0-2 R^{6e};

R^{11e} is selected from:

15 H, hydroxy, C₁-C₈ alkyl, C₃-C₆ alkenyl, C₃-C₁₁
cycloalkyl, (C₃-C₁₁ cycloalkyl)methyl, C₁-C₆ alkoxy,
benzyloxy, aryl, heteroaryl, heteroaryl(C₁-C₄ alkyl)-
, aryl(C₁-C₄ alkyl), adamantylmethyl, and
C₁-C₁₀ alkyl substituted with 0-2 R^{4e};

20

R^{4e} is selected from:

H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇
cycloalkyl(C₁-C₄ alkyl)-, aryl, heteroaryl, aryl(C₁-
C₆ alkyl)-, and heteroaryl(C₁-C₆ alkyl)-, wherein
25 said aryl or heteroaryl groups are substituted with
0-2 substituents independently selected from the
group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, F, Cl,
Br, CF₃, and NO₂,

30 R^{12e} is selected from:

H, C₁-C₆ alkyl, triphenylmethyl, methoxymethyl,
methoxyphenyldiphenylmethyl,
trimethylsilylethoxymethyl, (C₁-C₆ alkyl)carbonyl,
(C₁-C₆ alkoxy)carbonyl, (C₁-C₆ alkyl)aminocarbonyl,
35 C₃-C₆ alkenyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-
C₄ alkyl)-, aryl, heteroaryl(C₁-C₆ alkyl)carbonyl,
heteroarylcarbonyl, aryl(C₁-C₆ alkyl)-,
(C₁-C₆ alkyl)carbonyl, arylcarbonyl, C₁-C₆

alkylsulfonyl, arylsulfonyl, aryl(C₁-C₆
alkyl)sulfonyl, heteroarylsulfonyl, heteroaryl(C₁-C₆
alkyl)sulfonyl, aryloxy carbonyl, and aryl(C₁-C₆
alkoxy)carbonyl, wherein said aryl groups are
5 substituted with 0-2 substituents selected from the
group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, halo,
CF₃, and nitro;

R^{16e} is selected from:
10 -C(=O)OR^{18ae}, -C(=O)R^{18be}, -C(=O)N(R^{18be})₂, -SO₂R^{18ae},
and -SO₂N(R^{18be})₂;

R^{17e} is selected from:
H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇
15 cycloalkyl(C₁-C₄ alkyl)-, aryl, aryl(C₁-C₆ alkyl)-,
and heteroaryl(C₁-C₆ alkyl);

R^{18ae} is selected from:
C₁-C₈ alkyl optionally substituted with a bond to L_n,
20 C₃-C₁₁ cycloalkyl optionally substituted with a
bond to L_n, aryl(C₁-C₆ alkyl)- optionally
substituted with a bond to L_n, heteroaryl(C₁-C₆
alkyl)- optionally substituted with a bond to
L_n, (C₁-C₆ alkyl)heteroaryl optionally
25 substituted with a bond to L_n, biaryl(C₁-C₆
alkyl) optionally substituted with a bond to
L_n, heteroaryl optionally substituted with a
bond to L_n, phenyl substituted with 3-4 R^{19e} and
optionally substituted with a bond to L_n,
30 naphthyl substituted with 0-4 R^{19e} and
optionally substituted with a bond to L_n, and a
bond to L_n, wherein said aryl or heteroaryl
groups are optionally substituted with 0-4 R^{19e};

35 R^{18be} is H or R^{18ae};

R^{19e} is selected from:

H, halogen, CF₃, CO₂H, CN, NO₂, -NR^{11e}R^{12e}, OCF₃,
 C₁-C₈ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl,
 C₃-C₁₁ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-,
 aryl(C₁-C₆ alkyl)-, C₁-C₆ alkoxy, C₁-C₄
 5 alkoxycarbonyl, aryl, aryl-O-, aryl-SO₂-,
 heteroaryl, and heteroaryl-SO₂-, wherein said aryl
 and heteroaryl groups are substituted with 0-4
 groups selected from hydrogen, halogen, CF₃, C₁-C₃
 alkyl, and C₁-C₃ alkoxy;

10

R^{20e} is selected from:
 hydroxy, C₁-C₁₀ alkyloxy, C₃-C₁₁ cycloalkyloxy,
 aryloxy, aryl(C₁-C₄ alkyl)oxy,
 C₂-C₁₀ alkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 15 C₂-C₁₀ alkoxycarbonyloxy(C₁-C₂ alkyl)oxy-,
 C₂-C₁₀ alkoxycarbonyl(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkoxycarbonyloxy(C₁-C₂ alkyl)oxy-,
 C₃-C₁₀ cycloalkoxycarbonyl(C₁-C₂ alkyl)oxy-,
 20 aryloxycarbonyl(C₁-C₂ alkyl)oxy-,
 aryloxycarbonyloxy(C₁-C₂ alkyl)oxy-,
 arylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 C₁-C₅ alkoxy(C₁-C₅ alkyl)carbonyloxy(C₁-C₂ alkyl)oxy,
 (5-(C₁-C₅ alkyl)-1,3-dioxa-cyclopenten-2-one-
 25 yl)methyloxy,
 (5-aryl-1,3-dioxa-cyclopenten-2-one-yl)methyloxy,
 and
 (R^{10e})(R^{11e})N-(C₁-C₁₀ alkoxy)-;

30 R^{21e} is selected from:

C₁-C₈ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl, (C₃-C₁₁
 cycloalkyl)methyl, aryl, aryl(C₁-C₄ alkyl)-; and
 C₁-C₁₀ alkyl substituted with 0-2 R^{7e};

35 R^{22e} is selected from:

-C(=O)-R^{18be}, -C(=O)N(R^{18be})₂, -C(=O)NHSO₂R^{18ae},
 -C(=O)NHC(=O)R^{18be}, -C(=O)NHC(=O)OR^{18ae}, and
 -C(=O)NHSO₂NHR^{18be};

m^e is 0-2;

n^e is 0-4; and

5

p^e is 0-2;

with the following proviso: n^e and m^e are chosen such
that the number of atoms connecting R^1 and $-COR^{20e}$ in
10 Formula (IV) is in the range of 8-14;

d is selected from 1, 2, 3, 4, and 5;

d' is 1-50;

15

W is independently selected at each occurrence from the
group: O, NH, $NHC(=O)$, $C(=O)NH$, $NR^8C(=O)$, $C(=O)N R^8$,
 $C(=O)$, $C(=O)O$, $OC(=O)$, $NHC(=S)NH$, $NHC(=O)NH$, SO_2 ,
(OCH_2CH_2) $_s$, (CH_2CH_2O) $_s$, ($OCH_2CH_2CH_2$) $_s$, ($CH_2CH_2CH_2O$) $_t$,
20 and (aa) $_t$;

aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-1
25 R^{10} , C_{3-10} cycloalkyl substituted with 0-1 R^{10} , and a
5-10 membered heterocyclic ring system containing
1-4 heteroatoms independently selected from N, S,
and O and substituted with 0-1 R^{10} ;

30 R^6 , R^{6a} , R^7 , R^{7a} , and R^8 are independently selected at
each occurrence from the group: H, $=O$, $COOH$, SO_3H ,
 C_1-C_5 alkyl substituted with 0-1 R^{10} , aryl
substituted with 0-1 R^{10} , benzyl substituted with 0-1
 R^{10} , and C_1-C_5 alkoxy substituted with 0-1 R^{10} ,
35 $NHC(=O)R^{11}$, $C(=O)NHR^{11}$, $NHC(=O)NHR^{11}$, NHR^{11} , R^{11} , and
a bond to Ch ;

k is 0 or 1;

s is selected from 0, 1, 2, 3, 4, and 5;
 s' is selected from 0, 1, 2, 3, 4, and 5;
 s" is selected from 0, 1, 2, 3, 4, and 5;
 t is selected from 0, 1, 2, 3, 4, and 5;

5

A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ are independently
 selected at each occurrence from the group: NR¹³,
 NR¹³R¹⁴, S, SH, S(Pg), OH, and a bond to L_n;

10 E is a bond, CH, or a spacer group independently selected
 at each occurrence from the group: C₁-C₁₀ alkyl
 substituted with 0-3 R¹⁷, aryl substituted with 0-3
 R¹⁷, C₃-₁₀ cycloalkyl substituted with 0-3 R¹⁷, and a
 5-10 membered heterocyclic ring system containing
 15 1-4 heteroatoms independently selected from N, S,
 and O and substituted with 0-3 R¹⁷;

R¹³, and R¹⁴ are each independently selected from the
 group: a bond to L_n, hydrogen, C₁-C₁₀ alkyl
 20 substituted with 0-3 R¹⁷, aryl substituted with 0-3
 R¹⁷, a 5-10 membered heterocyclic ring system
 containing 1-4 heteroatoms independently selected
 from N, S, and O and substituted with 0-3 R¹⁷, and
 an electron, provided that when one of R¹³ or R¹⁴ is
 25 an electron, then the other is also an electron;


alternatively, R¹³ and R¹⁴ combine to form =C(R²⁰)(R²¹);

R¹⁷ is independently selected at each occurrence from the
 30 group: a bond to L_n, =O, F, Cl, Br, I, -CF₃, -CN,
 -CO₂R¹⁸, -C(=O)R¹⁸, -C(=O)N(R¹⁸)₂, -CH₂OR¹⁸,
 -OC(=O)R¹⁸, -OC(=O)OR^{18a}, -OR¹⁸, -OC(=O)N(R¹⁸)₂,
 -NR¹⁹C(=O)R¹⁸, -NR¹⁹C(=O)OR^{18a}, -NR¹⁹C(=O)N(R¹⁸)₂,
 -NR¹⁹SO₂N(R¹⁸)₂, -NR¹⁹SO₂R^{18a}, -SO₃H, -SO₂R^{18a},
 35 -S(=O)R^{18a}, -SO₂N(R¹⁸)₂, -N(R¹⁸)₂, -NHC(=S)NHR¹⁸,
 =NOR¹⁸, -C(=O)NHN(R¹⁸)R^{18a}, -OCH₂CO₂H, and
 2-(1-morpholino)ethoxy;

R^{18} , R^{18a} , and R^{19} are independently selected at each occurrence from the group: a bond to L_n , H, and C₁-C₆ alkyl;

5 R^{20} and R^{21} are independently selected from the group: H, C₁-C₅ alkyl, $-CO_2R^{25}$, C₂-C₅ 1-alkene substituted with 0-3 R^{23} , C₂-C₅ 1-alkyne substituted with 0-3 R^{23} , aryl substituted with 0-3 R^{23} , and unsaturated 5-10 membered heterocyclic ring system containing 1-4
10 heteroatoms independently selected from N, S, and O and substituted with 0-3 R^{23} ;

alternatively, R^{20} and R^{21} , taken together with the divalent carbon radical to which they are attached
15 form:



20 R^{22} and R^{23} are independently selected from the group: H, and R^{24} ;

alternatively, R^{22} , R^{23} taken together form a fused aromatic or a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected
25 from N, S, and O;


R^{24} is independently selected at each occurrence from the group: $-CO_2R^{25}$, $-C(=O)N(R^{25})_2$, $-CH_2OR^{25}$, $-OC(=O)R^{25}$, $-OR^{25}$, $-SO_3H$, $-N(R^{25})_2$, and $-OCH_2CO_2H$; and,
30

R^{25} is independently selected at each occurrence from the group: H and C₁-C₃ alkyl.

4. A compound according to Claim 3, including stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, or pharmaceutically acceptable salt or prodrug forms thereof wherein:


5

R^{1e} is selected from:



10

and



R^{2e} and R^{3e} are independently selected from:

15 H, C₁-C₄ alkoxy, NR^{11e}R^{12e}, halogen, NO₂, CN, CF₃, C₁-C₆ alkyl, C₃-C₆ alkenyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl(C₁-C₆ alkyl)-, (C₁-C₆ alkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl, arylcarbonyl, and aryl substituted with 0-4 R^{7e},

20

alternatively, when R^{2e} and R^{3e} are substituents on adjacent atoms, they can be taken together with the carbon atoms to which they are attached to form a 5-7 membered carbocyclic or 5-7 membered heterocyclic aromatic or nonaromatic ring system, said carbocyclic or heterocyclic ring being

25

substituted with 0-2 groups selected from C₁-C₄ alkyl, C₁-C₄ alkoxy, halo, cyano, amino, CF₃ and NO₂;

R^{2ae} is selected from:

30 H, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl, aryl(C₁-C₄ alkyl)-, (C₂-C₇ alkyl)carbonyl, arylcarbonyl, (C₂-C₁₀ alkoxy)carbonyl, C₃-C₇ cycloalkoxycarbonyl, C₇-C₁₁ bicycloalkoxycarbonyl, aryloxy carbonyl, 35 aryl(C₁-C₁₀ alkoxy)carbonyl,

C₁-C₆ alkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl,
 arylcarbonyloxy(C₁-C₄ alkoxy)carbonyl, and
 C₃-C₇ cycloalkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl;

5 R^{7e} is selected from:

H, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, aryl, aryl(C₁-
 C₄ alkyl)-, (C₁-C₄ alkyl)carbonyl, CO₂R^{18ae}, SO₂R^{11e},
 SO₂NR^{10e}R^{11e}, OR^{10e}, and N(R^{11e})R^{12e};

10 U^e is selected from:

-(CH₂)_n^e-, -NH(CH₂)_n^e-, -N(R^{10e})C(=O)-, and
 -NHC(=O)(CH₂)_n^e;

G^e is N or CR^{19e};

15

R^{8e} is H;

R^{9e} is selected from:

20 H, nitro, N(R^{11e})R^{12e}, OC(=O)R^{10e}, OR^{10e},
 OC(=O)NR^{10e}R^{11e}, NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},
 NR^{10e}C(=O)NR^{10e}R^{11e}, NR^{10e}SO₂NR^{10e}R^{11e},
 NR^{10e}SO₂R^{21e}, hydroxy, OR^{22e}, -N(R^{10e})R^{11e},
 -N(R^{16e})R^{17e}, aryl(C₀-C₄ alkyl)carbonyl, aryl(C₁-
 C₄ alkyl), heteroaryl(C₁-C₄ alkyl), CONR^{18aer}R^{20e},
 25 SO₂R^{18ae}, and SO₂NR^{18aer}R^{20e},

providing that any of the above alkyl, cycloalkyl,
 aryl or heteroaryl groups may be unsubstituted
 or substituted independently with 1-2 R^{7e};

30 R^{10e} is selected from:

H, CF₃, C₃-C₆ alkenyl, C₃-C₆ cycloalkyl, aryl,
 (C₃-C₆ cycloalkyl)methyl, aryl(C₁-C₄ alkyl), and
 C₁-C₄ alkyl substituted with 0-2 R^{6e};

35 R^{6e} is selected from:

H, C₁-C₄ alkyl, hydroxy, C₁-C₄ alkoxy, nitro, C₁-C₄
 alkylcarbonyl, -N(R^{11e})R^{12e}, cyano, halo, CF₃,
 CHO, CO₂R^{18be}, C(=O)R^{18be}, CONR^{17e}R^{18be},

- $\text{OC}(=\text{O})\text{R}^{10\text{e}}$, $\text{OR}^{10\text{e}}$, $\text{OC}(=\text{O})\text{NR}^{10\text{e}}\text{R}^{11\text{e}}$,
 $\text{NR}^{10\text{e}}\text{C}(=\text{O})\text{R}^{10\text{e}}$, $\text{NR}^{10\text{e}}\text{C}(=\text{O})\text{OR}^{21\text{e}}$,
 $\text{NR}^{10\text{e}}\text{C}(=\text{O})\text{NR}^{10\text{e}}\text{R}^{11\text{e}}$, $\text{NR}^{10\text{e}}\text{SO}_2\text{NR}^{10\text{e}}\text{R}^{11\text{e}}$,
 $\text{NR}^{10\text{e}}\text{SO}_2\text{R}^{21\text{e}}$, $\text{S}(\text{O})_p\text{R}^{11\text{e}}$, $\text{SO}_2\text{NR}^{10\text{e}}\text{R}^{11\text{e}}$,
5 aryl substituted with 0-3 groups selected from
halogen, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_1\text{-C}_4$ alkyl, CF_3 ,
 $\text{S}(\text{O})_m\text{Me}$, and $-\text{NMe}_2$,
aryl($\text{C}_1\text{-C}_4$ alkyl)-, said aryl being substituted with
0-3 groups selected from halogen, $\text{C}_1\text{-C}_4$ alkoxy,
10 $\text{C}_1\text{-C}_4$ alkyl, CF_3 , $\text{S}(\text{O})_p\text{Me}$, and $-\text{NMe}_2$, and
a 5-10 membered heterocyclic ring containing 1-3 N,
O, or S heteroatoms, wherein said heterocyclic ring
may be saturated, partially saturated, or fully
unsaturated, said heterocyclic ring being
15 substituted with 0-2 $\text{R}^{7\text{e}}$;

- $\text{R}^{11\text{e}}$ is selected from:
H, hydroxy, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_3\text{-C}_6$ alkenyl, $\text{C}_3\text{-C}_6$
cycloalkyl, ($\text{C}_3\text{-C}_6$ cycloalkyl)methyl, $\text{C}_1\text{-C}_4$ alkoxy,
20 benzyloxy, aryl, heteroaryl,
heteroaryl($\text{C}_1\text{-C}_4$ alkyl)-, aryl($\text{C}_1\text{-C}_4$ alkyl),
adamantylmethyl, and
 $\text{C}_1\text{-C}_4$ alkyl substituted with 0-2 $\text{R}^{4\text{e}}$;
25 $\text{R}^{4\text{e}}$ is selected from:
H, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_3\text{-C}_7$ cycloalkyl, $\text{C}_3\text{-C}_7$
cycloalkyl($\text{C}_1\text{-C}_4$ alkyl)-, aryl, heteroaryl, aryl($\text{C}_1\text{-C}_4$
 C_4 alkyl)-, and heteroaryl($\text{C}_1\text{-C}_4$ alkyl)-, wherein
said aryl or heteroaryl groups are substituted with
30 0-2 substituents independently selected from the
group consisting of $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, F, Cl,
Br, CF_3 , and NO_2 ,

- $\text{R}^{12\text{e}}$ is selected from:
35 H, $\text{C}_1\text{-C}_4$ alkyl, ($\text{C}_1\text{-C}_4$ alkyl)carbonyl, ($\text{C}_1\text{-C}_4$
alkoxy)carbonyl, phenyl($\text{C}_1\text{-C}_4$ alkyl)-,
phenylsulfonyl, phenyloxycarbonyl, and phenyl($\text{C}_1\text{-C}_4$
alkoxy)carbonyl, wherein said phenyl groups are

substituted with 0-2 substituents selected from the group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, halo, CF₃, and nitro;

5 R^{16e} is selected from:

-C(=O)OR^{18ae} -C(=O)R^{18be}, -C(=O)N(R^{18be})₂, -SO₂R^{18ae},
and -SO₂N(R^{18be})₂;

R^{17e} is selected from:

10 H, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, C₃-C₆
cycloalkyl(C₁-C₄ alkyl)-, aryl, aryl(C₁-C₆ alkyl)-,
and heteroaryl(C₁-C₆ alkyl);

R^{18ae} is selected from:

15 C₁-C₈ alkyl optionally substituted with a bond to L_n,
C₃-C₁₁ cycloalkyl optionally substituted with a
bond to L_n, aryl(C₁-C₆ alkyl)- optionally
substituted with a bond to L_n, heteroaryl(C₁-C₆
20 alkyl)- optionally substituted with a bond to
L_n, (C₁-C₆ alkyl)heteroaryl optionally
substituted with a bond to L_n, biaryl(C₁-C₆
alkyl) optionally substituted with a bond to
L_n, heteroaryl optionally substituted with a
bond to L_n, phenyl substituted with 3-4 R^{19e} and
25 optionally substituted with a bond to L_n,
naphthyl substituted with 0-4 R^{19e} and
optionally substituted with a bond to L_n, and a
bond to L_n, wherein said aryl or heteroaryl
groups are optionally substituted with 0-4 R^{19e};

30

R^{18be} is H or R^{18ae};

R^{19e} is selected from:

35 H, halogen, CF₃, CO₂H, CN, NO₂, -NR^{11e}R^{12e}, OCF₃,
C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl,
C₃-C₆ cycloalkyl, C₃-C₆ cycloalkyl(C₁-C₄ alkyl)-,
aryl(C₁-C₄ alkyl)-, C₁-C₆ alkoxy, C₁-C₄
alkoxycarbonyl, aryl, aryl-O-, aryl-SO₂-,

heteroaryl, and heteroaryl-SO₂-, wherein said aryl and heteroaryl groups are substituted with 0-4 groups selected from hydrogen, halogen, CF₃, C₁-C₃ alkyl, and C₁-C₃ alkoxy;

5

R^{20e} is selected from:

hydroxy, C₁-C₆ alkyloxy, C₃-C₆ cycloalkyloxy, aryloxy, aryl(C₁-C₄ alkyl)oxy, C₂-C₁₀ alkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
 10 C₂-C₁₀ alkoxy carbonyloxy(C₁-C₂ alkyl)oxy-, C₂-C₁₀ alkoxy carbonyl(C₁-C₂ alkyl)oxy-, C₃-C₁₀ cycloalkylcarbonyloxy(C₁-C₂ alkyl)oxy-, C₃-C₁₀ cycloalkoxy carbonyloxy(C₁-C₂ alkyl)oxy-, C₃-C₁₀ cycloalkoxy carbonyl(C₁-C₂ alkyl)oxy-,
 15 aryloxy carbonyl(C₁-C₂ alkyl)oxy-, aryloxy carbonyloxy(C₁-C₂ alkyl)oxy-, arylcarbonyloxy(C₁-C₂ alkyl)oxy-, C₁-C₅ alkoxy(C₁-C₅ alkyl)carbonyloxy(C₁-C₂ alkyl)oxy, (5-(C₁-C₅ alkyl)-1,3-dioxa-cyclopenten-2-one-yl)methyloxy,
 20 (5-aryl-1,3-dioxa-cyclopenten-2-one-yl)methyloxy, and (R^{10e})(R^{11e})N-(C₁-C₁₀ alkoxy)-;

25 R^{21e} is selected from:

C₁-C₄ alkyl, C₂-C₆ alkenyl, C₃-C₆ cycloalkyl, (C₃-C₆ cycloalkyl)methyl, aryl, aryl(C₁-C₄ alkyl)-, and C₁-C₁₀ alkyl substituted with 0-2 R^{7e};

30 R^{22e} is selected from:


-C(=O)-R^{18be}, -C(=O)N(R^{18be})₂, -C(=O)NHSO₂R^{18ae}, -C(=O)NHC(=O)R^{18be}, -C(=O)NHC(=O)OR^{18ae}, and -C(=O)NHSO₂NHR^{18be};

35 m^e is 0-2;

n^e is 0-4;

p^e is 0-2;

C_h is



A^1 is selected from the group: OH, and a bond to L_n ;

A^2 , A^4 , and A^6 are each N;

10

A^3 , A^5 , and A^8 are each OH;


A^7 is a bond to L_n or NH-bond to L_n ;

15 E is a C_2 alkyl substituted with 0-1 R^{17} ;

R^{17} is =O;

alternatively, C_h is

20



A^1 is selected from the group: OH, and a bond to L_n ;

25

A^2 , A^3 and A^4 are each N;

A^5 , A^6 and A^8 are each OH;

30 A^7 is a bond to L_n ;

E is a C₂ alkyl substituted with 0-1 R¹⁷;

R¹⁷ is =O;

5

alternatively, C_h is $A^1 \begin{array}{c} \diagup \\ E-A^2 \end{array}$;

A¹ is NH₂ or N=C(R²⁰)(R²¹);

10 E is a bond;

A² is NHR¹³;

15 R¹³ is a heterocycle substituted with R¹⁷, the heterocycle being selected from pyridine and pyrimidine;

R¹⁷ is selected from a bond to L_n, C(=O)NHR¹⁸ and C(=O)R¹⁸;

20 R¹⁸ is a bond to L_n;

R²⁴ is selected from the group: -CO₂R²⁵, -OR²⁵, -SO₃H, and -N(R²⁵)₂; and,

25 R²⁵ is independently selected at each occurrence from the group: hydrogen and methyl.

5. A compound according to Claim 2, including enantiomeric or diastereomeric forms thereof, or mixtures
30 of enantiomeric or diastereomeric forms thereof, or pharmaceutically acceptable salt or prodrug forms thereof, wherein Q is selected from the group:

35 3-[7-[(imidazolin-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(3,5-dimethylisoxazol-4-ylsulfonylamino)propionic acid,

- 3-[7-[(imidazolin-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
- 5 3-[7-[(imidazolin-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butyloxycarbonylamino)propionic acid,
- 3-[7-[(imidazolin-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butylsulfonylamino)propionic acid,
- 10 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
- 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butyloxycarbonylamino)propionic acid,
- 15 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(phenylsulfonylamino)propionic acid,
- 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butylsulfonyl)aminopropionic acid,
- 20 3-[7-[(2-aminothiazol-4-yl)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
- 25 3-[7-[(imidazolin-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 30 3-[7-[(imidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(3,5-dimethylisoxazol-4-ylsulfonylamino)propionic acid,
- 35

- 3-[7-[(imidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
- 5 3-[7-[(imidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 10 3-[7-[(imidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((4-biphenyl)sulfonylamino)propionic acid,
- 3-[7-[(imidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(1-naphthylsulfonylamino)propionic acid,
- 15 3-[7-[(benzimidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 20 3-[7-[(4-methylimidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 3-[7-[(4,5-dimethylimidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 25 3-[7-[(4,5,6,7-tetrahydrobenzimidazol-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 30 3-[7-[(pyridin-2-ylamino)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 35 3-[7-(2-aminopyridin-6-yl)-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,

- 3-[7-[(7-azabenzimidazol-2-yl)methyl]-1-methyl-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
- 5 3-[7-[(benzimidazol-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]propionic acid,
- 3-[7-[(pyridin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]propionic acid,
- 10 3-[7-[(imidazolin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]propionic acid,
- 3-[7-[(imidazol-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]propionic acid,
- 15 3-[7-[(imidazolin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
- 20 3-[7-[(imidazolin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butyloxycarbonylamino)propionic acid,
- 25 3-[7-[(imidazolin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(phenylsulfonylamino)propionic acid,
- 30 3-[7-[(imidazolin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butylsulfonylamino)propionic acid,
- 35 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
- 3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-

- ylcarbonylamino]-2-(n-butylloxycarbonylamino)propionic acid,
3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(phenylsulfonylamino)propionic acid,
3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(n-butylsulfonyl)aminopropionic acid,
3-[7-[(2-aminothiazol-4-yl)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(phenylsulfonylamino)propionic acid,
3-[7-[(2-aminothiazol-4-yl)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
3-[7-[(imidazolin-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
3-[7-[(tetrahydropyrimid-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfon-ylamino)propionic acid,
3-[7-[(imidazol-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(benzyloxycarbonylamino)propionic acid,
3-[7-[(imidazol-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-(phenylsulfonylamino)propionic acid,
3-[7-[(imidazol-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-2-((2,6-dichlorophenyl)sulfonylamino)propionic acid,
3-[7-[(imidazol-2-ylamino)methyl]-1-(2-phenylethyl)-6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-


- 2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
3-[7-[(imidazol-2-ylamino)methyl]-1-(2-phenylethyl)-
6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-
2-((4-biphenyl)sulfonylamino)propionic acid,
3-[7-[(benzimidazol-2-ylamino)methyl]-1-(2-
phenylethyl)-6,8-difluoroquinoline-4-one-3-
ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
3-[7-[(4-methylimidazol-2-ylamino)methyl]-1-(2-
phenylethyl)-6,8-difluoroquinoline-4-one-3-
ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
3-[7-[(4,5-dimethylimidazol-2-ylamino)methyl]-1-(2-
phenylethyl)-6,8-difluoroquinoline-4-one-3-
ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
3-[7-[(4,5,6,7-tetrahydrobenzimidazol-2-
ylamino)methyl]-1-(2-phenylethyl)-6,8-
difluoroquinoline-4-one-3-ylcarbonylamino]-2-
((2,4,6-trimethylphenyl)sulfonylamino)propionic
acid,
3-[7-[(pyridin-2-ylamino)methyl]-1-(2-phenylethyl)-
6,8-difluoroquinoline-4-one-3-ylcarbonylamino]-
2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid,
3-[7-(2-aminopyridin-6-yl)-1-(2-phenylethyl)-6,8-
difluoroquinoline-4-one-3-ylcarbonylamino]-2-
((2,4,6-trimethylphenyl)sulfonylamino)propionic
acid, and
3-[7-[(7-azabenzimidazol-2-yl)methyl]-1-(2-
phenylethyl)-6,8-difluoroquinoline-4-one-3-
ylcarbonylamino]-2-((2,4,6-trimethylphenyl)sulfonylamino)propionic acid.


6. A compound according to Claim 2, wherein the compound is selected from the group:

- 2-(((4-(4-(((3-(2-(2-(3-((6-((1-aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)-propoxy)ethoxy)-ethoxy)propyl)amino)sulfonyl)phenyl)phenyl)-sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic acid;
- 3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((4-(4-(((3-(2-(2-(3-(2-(1,4,7,10-tetraaza-4,7,10-tris(carboxymethyl)cyclododecyl)acetyl)amino)-propoxy)ethoxy)ethoxy)propyl)amino)sulfonyl)-phenyl)phenyl)sulfonyl)amino)propanoic acid;
- 2-(((4-(3-(N-(3-(2-(2-(3-((6-((1-aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)-propoxy)ethoxy)ethoxy)propyl)carbamoyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)propanoic;
- 3-((1-(3-((6-((1-aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)propyl)-7-((imidazole-2-ylamino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic acid;
- 3-((1-(3-((6-((1-aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)propyl)-7-(((1-hydroxyimidazole-2-yl)amino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic acid;
- 3-((1-(3-(3-(N-(3-(2-(2-(3-((6-((1-aza-2-(2-sulfophenyl)vinyl)amino)(3-pyridyl))carbonylamino)propoxy)ethoxy)-ethoxy)propyl)carbamoyl)propanoylamino)propyl)-7-((imidazole-2-ylamino)methyl)-4-oxo(3-



hydroquinolyl))carbonylamino)-2-((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic acid;

2-(2-aza-2-(5-(N-(1,3-bis(3-(2-(2-(3-(3-(N-(3-(3-(N-(3-carboxy-2-((2,4,6-trimethylphenyl)sulfonyl)amino)-ethyl)carbamoyl)-7-((imidazole-2-ylamino)methyl)4-oxohydroquinolyl)propyl)carbamoyl)propanoylamino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)(2-pyridyl))amino)vinyl)benzenesulfonic acid;








5



10





5

2-(((4-(3-(N-(3-(2-(2-(3-(2-(1,4,7,10-tetraaza-4,7,10-
 tris(carboxymethyl)cyclododecylacetyl-amino)-6-
 amino-hexanoyl-amino)propoxy)ethoxy)ethoxy)-
 propyl)carbamoyl)propoxy)-2,6-
 10 dimethylphenyl)sulfonyl-amino)-3-((7-((imidazol-2-
 ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-
 carbonylamino)propanoic acid ;




15



2-(((4-(3-(N-(3-(2-(2-(3-(2-(1,4,7,10-tetraaza-4,7,10-
 tris(carboxymethyl)cyclododecylacetyl-amino)-6-(2-
 (bis(phosphonomethyl)amino)acetyl-amino)hexanoyl-amino
)propoxy)ethoxy)ethoxy)propyl)carbamoyl)propoxy)-
 2,6-dimethylphenyl)sulfonyl-amino)-3-((7-((imidazol-
 2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-
 20 carbonylamino)propanoic acid conjugate; and

25



2-(((4-(3-(N-(3-(2-(2-(3-(2-(2-((2-((2-
 (bis(carboxymethyl)-
 amino)ethyl)(carboxymethyl)amino)ethyl)(carboxymethy
 1)amino)acetyl-amino)-3-sulfo-propyl)propoxy)ethoxy)-
 ethoxy)propyl)carbamoyl)propoxy)-2,6-

dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-carbonylamino)propanoic acid;






5



; and



- 5 or a pharmaceutically acceptable salt form thereof.
7. A kit comprising a compound of Claim 2, or a pharmaceutically acceptable salt form thereof and a pharmaceutically acceptable carrier.
- 10
8. A kit according to Claim 7, wherein the kit further comprises one or more ancillary ligands and a reducing agent.
- 15 9. A kit according to Claim 8, wherein the ancillary ligands are tricine and TPPTS.
10. A kit according to Claim 8, wherein the reducing agent is tin(II).
- 20
11. A diagnostic or therapeutic metallopharmaceutical composition, comprising: a metal, a chelator capable of chelating the metal and a targeting moiety, wherein the targeting moiety is bound to the chelator, is a quinolone
- 25 non-peptide and binds to a receptor that is upregulated

during angiogenesis and the compound has 0-1 linking groups between the targeting moiety and chelator.

12. A composition according to Claim 11, wherein the
5 metallopharmaceutical is a diagnostic
radiopharmaceutical, the metal is a radioisotope selected from the group: ^{99m}Tc , ^{95}Tc , ^{111}In , ^{62}Cu , ^{64}Cu , ^{67}Ga , and ^{68}Ga , and the linking group is present between the non-peptide targeting moiety and chelator.

10

13. A composition according to Claim 12, wherein the targeting moiety is a quinolone non-peptide and the receptor is $\alpha_v\beta_3$ or $\alpha_v\beta_5$.

- 15 14. A composition according to Claim 13, wherein the radioisotope is ^{99m}Tc or ^{95}Tc , the radiopharmaceutical further comprises a first ancillary ligand and a second ancillary ligand capable of stabilizing the radiopharmaceutical.

20

15. A composition according to Claim 14, wherein the radioisotope is ^{99m}Tc .

16. A composition according to Claim 15, wherein the
25 radiopharmaceutical is selected from the group:

^{99m}Tc (2-(((4-(4-((3-(2-(2-(3-((6-(diazenido)(3-pyridyl))carbonylamino)propoxy)ethoxy)-ethoxy)propyl)amino)sulfonyl)phenyl)phenyl)-sulfonyl)amino)-3-((7-((imidazol-2-ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))carbonylamino)propanoic acid)(tricine)(TPPTS);

35 ^{99m}Tc (2-(((4-(3-(N-(3-(2-(2-(3-((6-(diazenido)(3-pyridyl))carbonylamino)propoxy)ethoxy)-ethoxy)propyl)carbonyl)propoxy)-2,6-dimethylphenyl)sulfonyl)amino)-3-((7-((imidazol-2-



ylamino)methyl)-1-methyl-4-oxo(3-hydroquinolyl))-
carbonylamino)propanoic acid)(tricine)(TPPDS);



- 5 ^{99m}Tc (3-((1-(3-((6-(diazenido)(3-pyridyl))carbonylamino)propyl)-7-((imidazole-2-ylamino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic acid)(tricine)(TPPTS);
- 10 ^{99m}Tc (3-((1-(3-((6-(diazenido)(3-pyridyl))carbonylamino)propyl)-7-(((1-hydroxyimidazole-2-yl)amino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic acid)(tricine)(TPPTS);
- 15 ^{99m}Tc (3-((1-(3-(3-(N-(3-(2-(2-(3-((6-(diazenido)(3-pyridyl))carbonylamino)propoxy)ethoxy)-ethoxy)propyl)carbamoyl)propanoylamino)propyl)-7-((imidazole-2-ylamino)methyl)-4-oxo(3-hydroquinolyl))carbonylamino)-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)propanoic acid)(tricine)(TPPTS);
- 20 ^{99m}Tc (2-(2-(5-(N-(1,3-bis(3-(2-(2-(3-(3-(N-(3-(3-(N-(3-carboxy-2-(((2,4,6-trimethylphenyl)sulfonyl)amino)-ethyl)carbamoyl)-7-((imidazole-2-ylamino)methyl)4-oxohydroquinolyl)propyl)carbamoyl)propanoylamino)propoxy)ethoxy)ethoxy)propyl)carbamoyl)(2-pyridyl)diazenido)))(tricine)(TPPTS);
- 25 ^{99m}Tc (3-([1-(3-{2-[(6-(diazenido)(3-pyridyl))carbonylamino](2R)-3-sulfopropyl}propyl)-7-[(imidazol-2-ylamino)methyl]-4-oxo(3-hydroquinolyl))carbonylamino)(2S)-2-[[2,4,6-
- 30

trimethylphenyl)sulfonyl]amino}propanoic acid)
(tricine) (TPPTS).


17. A composition according to Claim 13, wherein the
5 radioisotope is ^{111}In .


18. A composition according to Claim 17, wherein the
radiopharmaceutical is selected from the group:






5





; and



5

19. A composition according to Claim 11, wherein the metallopharmaceutical is a therapeutic radiopharmaceutical, the metal is a radioisotope selected
- 10 from the group: ^{186}Re , ^{188}Re , ^{153}Sm , ^{166}Ho , ^{177}Lu , ^{149}Pm , ^{90}Y , ^{212}Bi , ^{103}Pd , ^{109}Pd , ^{159}Gd , ^{140}La , ^{198}Au , ^{199}Au , ^{169}Yb , ^{175}Yb , ^{165}Dy , ^{166}Dy , ^{67}Cu , ^{105}Rh , ^{111}Ag , and ^{192}Ir , and the

linking group is present between the non-peptide targeting moiety and chelator.


20. A composition according to Claim 19, wherein the
5 targeting moiety is a quinolone non-peptide and the receptor is $\alpha_v\beta_3$ or $\alpha_v\beta_5$.




21. A composition according to Claim 20, wherein the radioisotope is ^{153}Sm .

10

22. A composition according to Claim 20, wherein the radioisotope is ^{177}Lu .

23. A composition according to Claim 22, wherein the
15 radiopharmaceutical is selected from the group:









5

24. A composition according to Claim 20, wherein the radioisotope is ^{90}Y .

10


25. A composition according to Claim 24, wherein the radiopharmaceutical is selected from the group;





26. A composition according to Claim 11, wherein the
 5 metallopharmaceutical is a MRI contrast agent, the metal
 is a paramagnetic metal ion selected from the group:
 Gd(III), Dy(III), Fe(III), and Mn(II), the targeting
 moiety is a quinolone nonpeptide and the linking group is
 present between the targeting moiety and chelator.
- 10 27. A composition according to Claim 26, wherein the
 targeting moiety is quinolone non-peptide and the
 receptor is $\alpha_v\beta_3$ or $\alpha_v\beta_5$.
- 15 28. A composition according to Claim 27, wherein the
 metal ion is Gd(III).
29. A composition according to Claim 28, wherein the
 contrast agent is

20



30. A composition according to Claim 11, wherein the metallopharmaceutical is a X-ray contrast agent, the metal is selected from the group: Re, Sm, Ho, Lu, Pm, Y, Bi, Pd, Gd, La, Au, Au, Yb, Dy, Cu, Rh, Ag, and Ir, the targeting moiety is a quinolone non-peptide, the receptor is $\alpha_v\beta_3$ or $\alpha_v\beta_5$, and the linking group is present between the targeting moiety and chelator.
31. A method of treating rheumatoid arthritis in a patient comprising: administering a therapeutic radiopharmaceutical of Claim 19 capable of localizing in new angiogenic vasculature to a patient by injection or infusion.
32. A method of treating cancer in a patient comprising: administering to a patient in need thereof a therapeutic radiopharmaceutical of Claim 19 by injection or infusion.
33. A method of imaging therapeutic angiogenesis in a patient comprising: (1) administering a diagnostic radiopharmaceutical, a MRI contrast agent, or a X-ray contrast agent of Claim 11 to a patient by injection or infusion; (2) imaging the area of the patient wherein the desired formation of new blood vessels is located.

34. A method of imaging cancer in a patient comprising:
(1) administering a diagnostic radiopharmaceutical of
Claim 12 to a patient by injection or infusion; (2)
imaging the patient using planar or SPECT gamma
5 scintigraphy, or positron emission tomography.

35. A method of imaging cancer in a patient comprising:
(1) administering a MRI contrast agent of Claim 26; and
(2) imaging the patient using magnetic resonance imaging.

10


36. A method of imaging cancer in a patient comprising:
(1) administering a X-ray contrast agent of Claim 30; and
(2) imaging the patient using X-ray computed tomography.

15 37. A compound, comprising: a targeting moiety and a
surfactant, wherein the targeting moiety is bound to the
surfactant, is a nonpeptide, and binds to a receptor that
is upregulated during angiogenesis and the compound has
0-1 linking groups between the targeting moiety and
20 surfactant.

38. A compound according to Claim 37, wherein the
targeting moiety comprises a quinolone non-peptide and
the linking group is present between the targeting moiety
25 and surfactant.


39. A compound according to Claim 38, wherein the
receptor is the integrin $\alpha_v\beta_3$ or $\alpha_v\beta_5$ and the compound is
of the formula:

30



wherein, Q is a compound of Formula (II):

35




(II)

- 5 including stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, or pharmaceutically acceptable salt or prodrug forms thereof wherein:

R^{1e} is selected from:

10



15

A^e is -CH₂- or -N(R^{10e})-;

A^{1e} and B^e are independently -CH₂- or -N(R^{10e})-;

5

D^e is -N(R^{10e})- or -S-;

E^e-F^e is -C(R^{2e})=C(R^{3e})- or -C(R^{2e})₂C(R^{3e})₂-;

10 J^e is -C(R^{2e})- or -N-;

K^e, L^e and M^e are independently -C(R^{2e})- or -C(R^{3e})-;

R^{2e} and R^{3e} are independently selected from:

15 H, C₁-C₄ alkoxy, NR^{11e}R^{12e}, halogen, NO₂, CN, CF₃,
C₁-C₆ alkyl, C₃-C₆ alkenyl, C₃-C₇ cycloalkyl,
C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl(C₁-C₆ alkyl)-,
(C₁-C₆ alkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl,
arylcarbonyl, and aryl substituted with 0-4 R^{7e},

20

alternatively, when R^{2e} and R^{3e} are substituents on
adjacent atoms, they can be taken together with the
carbon atoms to which they are attached to form a 5-
7 membered carbocyclic or 5-7 membered heterocyclic
aromatic or nonaromatic ring system, said
25 carbocyclic or heterocyclic ring being substituted
with 0-2 groups selected from C₁-C₄ alkyl, C₁-C₄
alkoxy, halo, cyano, amino, CF₃ and NO₂;

30 R^{2ae} is selected from:

H, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl,
C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl, aryl(C₁-C₄
alkyl)-, (C₂-C₇ alkyl)carbonyl, arylcarbonyl,
(C₂-C₁₀ alkoxy)carbonyl, C₃-C₇ cycloalkoxycarbonyl,
35 C₇-C₁₁ bicycloalkoxycarbonyl, aryloxy carbonyl,
aryl(C₁-C₁₀ alkoxy)carbonyl,
C₁-C₆ alkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl,
arylcarbonyloxy(C₁-C₄ alkoxy)carbonyl, and

C₃-C₇ cycloalkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl;

R^{7e} is selected from:

- 5 H, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, aryl, aryl(C₁-C₄ alkyl)-, (C₁-C₄ alkyl)carbonyl, CO₂R^{18ae}, SO₂R^{11e}, SO₂NR^{10e}R^{11e}, OR^{10e}, and N(R^{11e})R^{12e};

U^e is selected from:

- 10 -(CH₂)_n^e-, -(CH₂)_n^eO(CH₂)_m^e-, -(CH₂)_n^eN(R¹²)(CH₂)_m^e-,
 -NH(CH₂)_n^e-, -(CH₂)_n^eC(=O)(CH₂)_m^e-,
 -(CH₂)_n^eS(O)_p^e(CH₂)_m^e-, -(CH₂)_n^eNHNH(CH₂)_m^e-,
 -N(R^{10e})C(=O)-, -NHC(=O)(CH₂)_n^e-, -C(=O)N(R^{10e})-, and
 -N(R^{10e})S(O)_p^e-;

- 15 G^e is N or CR^{19e};

W^e is -C(=O)-N(R^{10e})-(C₁-C₃ alkylene)-, in which the
 alkylene group is substituted by R^{8e} and by R^{9e}:

- 20 R^{8e} and R^{9e} are independently selected from:

- H, CO₂R^{18be}, C(=O)R^{18be}, CONR¹⁷R^{18be},
 C₁-C₁₀ alkyl substituted with 0-1 R^{6e},
 C₂-C₁₀ alkenyl substituted with 0-1 R^{6e},
 C₂-C₁₀ alkynyl substituted with 0-1 R^{6e},
 25 C₃-C₈ cycloalkyl substituted with 0-1 R^{6e},
 C₅-C₆ cycloalkenyl substituted with 0-1 R^{6e},
 (C₁-C₁₀ alkyl)carbonyl,
 C₃-C₁₀ cycloalkyl(C₁-C₄ alkyl)-,
 phenyl substituted with 0-3 R^{6e},
 30 naphthyl substituted with 0-3 R^{6e},
 a 5-10 membered heterocyclic ring containing 1-3 N,
 O, or S heteroatoms, wherein said heterocyclic
 ring may be saturated, partially saturated, or
 fully unsaturated, said heterocyclic ring being
 35 substituted with 0-2 R^{7e},
 C₁-C₁₀ alkoxy substituted with 0-2 R^{7e},
 hydroxy, nitro, -N(R^{10e})R^{11e}, -N(R^{16e})R^{17e}, aryl(C₀-C₆
 alkyl)carbonyl, aryl(C₃-C₆ alkyl),

heteroaryl(C₁-C₆ alkyl), CONR^{18ae}R^{20e}, SO₂R^{18ae},
and SO₂NR^{18ae}R^{20e},

providing that any of the above alkyl, cycloalkyl,
aryl or heteroaryl groups may be unsubstituted
5 or substituted independently with 1-2 R^{7e};

R^{6e} is selected from:

H, C₁-C₁₀ alkyl, hydroxy, C₁-C₁₀ alkoxy, nitro, C₁-C₁₀
alkylcarbonyl, -N(R^{11e})R^{12e}, cyano, halo, CF₃,
10 CHO, CO₂R^{18be}, C(=O)R^{18be}, CONR^{17e}R^{18be},
OC(=O)R^{10e}, OR^{10e}, OC(=O)NR^{10e}R^{11e},
NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},
NR^{10e}C(=O)NR^{10e}R^{11e}, NR^{10e}SO₂NR^{10e}R^{11e},
NR^{10e}SO₂R^{21e}, S(O)_pR^{11e}, SO₂NR^{10e}R^{11e},
15 aryl substituted with 0-3 groups selected from
halogen, C₁-C₆ alkoxy, C₁-C₆ alkyl, CF₃,
S(O)_m^eMe, and -NMe₂,
aryl(C₁-C₄ alkyl)-, said aryl being substituted with
0-3 groups selected from halogen, C₁-C₆ alkoxy,
20 C₁-C₆ alkyl, CF₃, S(O)_p^eMe, and -NMe₂, and
a 5-10 membered heterocyclic ring containing 1-3 N, O, or
S heteroatoms, wherein said heterocyclic ring may be
saturated, partially saturated, or fully unsaturated,
said heterocyclic ring being substituted with 0-2 R^{7e};

25

R^{10e} is selected from:

H, CF₃, C₃-C₆ alkenyl, C₃-C₁₁ cycloalkyl, aryl,
(C₃-C₁₁ cycloalkyl)methyl, aryl(C₁-C₄ alkyl), and C₁-
C₁₀ alkyl substituted with 0-2 R^{6e};

30

R^{11e} is selected from:

H, hydroxy, C₁-C₈ alkyl, C₃-C₆ alkenyl, C₃-C₁₁
cycloalkyl, (C₃-C₁₁ cycloalkyl)methyl, C₁-C₆ alkoxy,
benzyloxy, aryl, heteroaryl, heteroaryl(C₁-C₄ alkyl)-
35 , aryl(C₁-C₄ alkyl), adamantylmethyl, and
C₁-C₁₀ alkyl substituted with 0-2 R^{4e};

R^{4e} is selected from:

- H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-, (C₁-C₁₀ alkyl)carbonyl, aryl, heteroaryl, aryl(C₁-C₆ alkyl)-, and heteroaryl(C₁-C₆ alkyl)-, wherein said aryl or heteroaryl groups are substituted with 0-2 substituents independently selected from the group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, F, Cl, Br, CF₃, and NO₂,
- 10 alternatively, when R^{10e} and R^{11e} are both substituents on the same nitrogen atom (as in -NR^{10e}R^{11e}) they may be taken together with the nitrogen atom to which they are attached to form a heterocycle selected from: 3-azabicyclononyl, 1,2,3,4-tetrahydro-1-quinolinyl, 1,2,3,4-tetrahydro-2-isoquinolinyl, 1-piperidinyl, 1-morpholinyl, 1-pyrrolidinyl, thiamorpholinyl, thiazolidinyl, and 1-piperazinyl; said heterocycle being substituted with 0-3 groups selected from: C₁-C₆ alkyl, aryl, heteroaryl, aryl(C₁-C₄ alkyl)-, (C₁-C₆ alkyl)carbonyl, (C₃-C₇ cycloalkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl, aryl(C₁-C₄ alkoxy)carbonyl, C₁-C₆ alkylsulfonyl, and arylsulfonyl;
- 25 R^{12e} is selected from: H, C₁-C₆ alkyl, triphenylmethyl, methoxymethyl, methoxyphenyldiphenylmethyl, trimethylsilylethoxymethyl, (C₁-C₆ alkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl, (C₁-C₆ alkyl)aminocarbonyl, C₃-C₆ alkenyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-, aryl, heteroaryl(C₁-C₆ alkyl)carbonyl, heteroarylcabonyl, aryl(C₁-C₆ alkyl)-, (C₁-C₆ alkyl)carbonyl, arylcarbonyl, C₁-C₆ alkylsulfonyl, arylsulfonyl, aryl(C₁-C₆ alkyl)sulfonyl, heteroarylsulfonyl, heteroaryl(C₁-C₆ alkyl)sulfonyl, aryloxy carbonyl, and aryl(C₁-C₆ alkoxy)carbonyl, wherein said aryl groups are substituted with 0-2 substituents selected from the

group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, halo, CF₃, and nitro;

R^{16e} is selected from:

- 5 -C(=O)OR^{18ae}, -C(=O)R^{18be}, -C(=O)N(R^{18be})₂,
 -C(=O)NHSO₂R^{18ae}, -C(=O)NHC(=O)R^{18be},
 -C(=O)NHC(=O)OR^{18ae}, -C(=O)NHSO₂NHR^{18be}, -SO₂R^{18ae},
 -SO₂N(R^{18be})₂, and -SO₂NHC(=O)OR^{18be};

10 R^{17e} is selected from:

H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇
 cycloalkyl(C₁-C₄ alkyl)-, aryl, aryl(C₁-C₆ alkyl)-,
 and heteroaryl(C₁-C₆ alkyl);

15 R^{18ae} is selected from:

- C₁-C₈ alkyl optionally substituted with a bond to L_n,
 C₃-C₁₁ cycloalkyl optionally substituted with a
 bond to L_n, aryl(C₁-C₆ alkyl)- optionally
 substituted with a bond to L_n, heteroaryl(C₁-C₆
 20 alkyl)- optionally substituted with a bond to
 L_n, (C₁-C₆ alkyl)heteroaryl optionally
 substituted with a bond to L_n, biaryl(C₁-C₆
 alkyl) optionally substituted with a bond to
 L_n, heteroaryl optionally substituted with a
 25 bond to L_n, phenyl substituted with 3-4 R^{19e} and
 optionally substituted with a bond to L_n,
 naphthyl substituted with 0-4 R^{19e} and
 optionally substituted with a bond to L_n, and a
 bond to L_n, wherein said aryl or heteroaryl
 30 groups are optionally substituted with 0-4 R^{19e};

R^{18be} is H or R^{18ae};

R^{19e} is selected from:

- 35 H, halogen, CF₃, CO₂H, CN, NO₂, -NR^{11e}R^{12e}, OCF₃,
 C₁-C₈ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl,
 C₃-C₁₁ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-,
 aryl(C₁-C₆ alkyl)-, C₁-C₆ alkoxy, C₁-C₄

alkoxycarbonyl, aryl, aryl-O-, aryl-SO₂-,
heteroaryl, and heteroaryl-SO₂-, wherein said aryl
and heteroaryl groups are substituted with 0-4
groups selected from hydrogen, halogen, CF₃, C₁-C₃
5 alkyl, and C₁-C₃ alkoxy;

R^{20e} is selected from:
hydroxy, C₁-C₁₀ alkyloxy, C₃-C₁₁ cycloalkyloxy,
aryloxy, aryl(C₁-C₄ alkyl)oxy,
10 C₂-C₁₀ alkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
C₂-C₁₀ alkoxycarbonyloxy(C₁-C₂ alkyl)oxy-,
C₂-C₁₀ alkoxycarbonyl(C₁-C₂ alkyl)oxy-,
C₃-C₁₀ cycloalkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
C₃-C₁₀ cycloalkoxycarbonyloxy(C₁-C₂ alkyl)oxy-,
15 C₃-C₁₀ cycloalkoxycarbonyl(C₁-C₂ alkyl)oxy-,
aryloxy carbonyl(C₁-C₂ alkyl)oxy-,
aryloxy carbonyloxy(C₁-C₂ alkyl)oxy-,
arylcarbonyloxy(C₁-C₂ alkyl)oxy-,
C₁-C₅ alkoxy(C₁-C₅ alkyl)carbonyloxy(C₁-C₂ alkyl)oxy,
20 (5-(C₁-C₅ alkyl)-1,3-dioxo-cyclopenten-2-one-
yl)methyloxy,
(5-aryl-1,3-dioxo-cyclopenten-2-one-yl)methyloxy,
and
(R^{10e})(R^{11e})N-(C₁-C₁₀ alkoxy)-;


25 R^{21e} is selected from:
C₁-C₈ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl, (C₃-C₁₁
cycloalkyl)methyl, aryl, aryl(C₁-C₄ alkyl)-, and C₁-
C₁₀ alkyl substituted with 0-2 R^{7e};

30 R^{22e} is selected from:
-C(=O)-R^{18be}, -C(=O)N(R^{18be})₂, -C(=O)NHSO₂R^{18ae}, -
C(=O)NHC(=O)R^{18be}, -C(=O)NHC(=O)OR^{18ae}, and -
C(=O)NHSO₂NHR^{18be};

35 Y^e is selected from:

-COR^{20e}, -SO₃H, -PO₃H, -CONHNHSO₂CF₃, -CONHSO₂R^{18ae},
 -CONHSO₂NHR^{18be}, -NHCOCF₃, -NHCONHSO₂R^{18ae},
 -NHSO₂R^{18ae}, -OPO₃H₂, -OSO₃H, -PO₃H₂, -SO₂NHCOR^{18ae},
 -SO₂NHCO₂R^{18ae},

5



m^e is 0-2;

10 n^e is 0-4;

p^e is 0-2;


r^e is 0-2;

15

with the following proviso: n^e and m^e are chosen such
 that the number of atoms connecting R^{1e} and Y^e is in
 the range of 8-14;

20 d is selected from 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

L_n is a linking group having the formula:



25

W is independently selected at each occurrence from the
 group: O, S, NH, NHC(=O), C(=O)NH, NR⁸C(=O), C(=O)N
 R⁸, C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂,
 SO₂NH, (OCH₂CH₂)₂₀₋₂₀₀, (CH₂CH₂O)₂₀₋₂₀₀, (OCH₂CH₂CH₂)₂₀₋
 30 200, (CH₂CH₂CH₂O)₂₀₋₂₀₀, and (aa)_t;

aa is independently at each occurrence an amino acid;

35 Z is selected from the group: aryl substituted with 0-3
 R¹⁰, C₃₋₁₀ cycloalkyl substituted with 0-3 R¹⁰, and a
 5-10 membered heterocyclic ring system containing

1-4 heteroatoms independently selected from N, S,
and O and substituted with 0-3 R¹⁰;

5 R⁶, R^{6a}, R⁷, R^{7a}, and R⁸ are independently selected at
each occurrence from the group: H, =O, COOH, SO₃H,
PO₃H, C₁-C₅ alkyl substituted with 0-3 R¹⁰, aryl
substituted with 0-3 R¹⁰, benzyl substituted with 0-3
R¹⁰, and C₁-C₅ alkoxy substituted with 0-3 R¹⁰,
10 NHC(=O)R¹¹, C(=O)NHR¹¹, NHC(=O)NHR¹¹, NHR¹¹, R¹¹, and
a bond to S_f;

R¹⁰ is independently selected at each occurrence from the
group: a bond to S_f, COOR¹¹, C(=O)NHR¹¹, NHC(=O)R¹¹,
OH, NHR¹¹, SO₃H, PO₃H, -OPO₃H₂, -OSO₃H, aryl
15 substituted with 0-3 R¹¹, C₁-5 alkyl substituted with
0-1 R¹², C₁-5 alkoxy substituted with 0-1 R¹², and a
5-10 membered heterocyclic ring system containing
1-4 heteroatoms independently selected from N, S,
and O and substituted with 0-3 R¹¹;

20 R¹¹ is independently selected at each occurrence from the
group: H, alkyl substituted with 0-1 R¹², aryl
substituted with 0-1 R¹², a 5-10 membered
heterocyclic ring system containing 1-4 heteroatoms
25 independently selected from N, S, and O and
substituted with 0-1 R¹², C₃-10 cycloalkyl
substituted with 0-1 R¹², polyalkylene glycol
substituted with 0-1 R¹², carbohydrate substituted
with 0-1 R¹², cyclodextrin substituted with 0-1 R¹²,
30 amino acid substituted with 0-1 R¹², polycarboxyalkyl
substituted with 0-1 R¹², polyazaalkyl substituted
with 0-1 R¹², peptide substituted with 0-1 R¹²,
wherein the peptide is comprised of 2-10 amino
acids, 3,6-O-disulfo-B-D-galactopyranosyl,
35 bis(phosphonomethyl)glycine, and a bond to S_f;

R¹² is a bond to S_f;

k is selected from 0, 1, and 2;
 h is selected from 0, 1, and 2;
 h' is selected from 0, 1, and 2;
 g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 5 g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 x is selected from 0, 1, 2, 3, 4, and 5;
 x' is selected from 0, 1, 2, 3, 4, and 5;

10

S_f is a surfactant which is a lipid or a compound of the

formula: $A^9-E^1-A^{10}$;

15

A⁹ is selected from the group: OH and OR²⁷;

A¹⁰ is OR²⁷;

20 R²⁷ is C(=O)C₁₋₂₀ alkyl;

E¹ is C₁₋₁₀ alkylene substituted with 1-3 R²⁸;

25 R²⁸ is independently selected at each occurrence from the
 group: R³⁰, -PO₃H-R³⁰, =O, -CO₂R²⁹, -C(=O)R²⁹,
 -C(=O)N(R²⁹)₂, -CH₂OR²⁹, -OR²⁹, -N(R²⁹)₂, C₁-C₅
 alkyl, and C₂-C₄ alkenyl;


30 R²⁹ is independently selected at each occurrence from the
 group: R³⁰, H, C₁-C₆ alkyl, phenyl, benzyl, and
 trifluoromethyl;

R³⁰ is a bond to L_n;

35


and a pharmaceutically acceptable salt thereof.

40. A compound according to Claim 39, wherein the compound is of the formula:



5

wherein, Q is a compound of Formula (IV):




10

(IV)

including stereoisomeric forms thereof, or mixtures of stereoisomeric forms thereof, or pharmaceutically acceptable salt or prodrug forms thereof wherein:

15

R1e is selected from:



25

R^{2e} and R^{3e} are independently selected from:

H, C₁-C₄ alkoxy, NR^{11e}R^{12e}, halogen, NO₂, CN, CF₃,
C₁-C₆ alkyl, C₃-C₆ alkenyl, C₃-C₇ cycloalkyl,
C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl(C₁-C₆ alkyl)-,
5 (C₁-C₆ alkyl)carbonyl, (C₁-C₆ alkoxy)carbonyl,
arylcabonyl, and aryl substituted with 0-4 R^{7e},

alternatively, when R^{2e} and R^{3e} are substituents on
adjacent atoms, they can be taken together with the
10 carbon atoms to which they are attached to form a 5-
7 membered carbocyclic or 5-7 membered heterocyclic
aromatic or nonaromatic ring system, said
carbocyclic or heterocyclic ring being substituted
with 0-2 groups selected from C₁-C₄ alkyl, C₁-C₄
15 alkoxy, halo, cyano, amino, CF₃ and NO₂;

R^{2ae} is selected from:

H, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl,
C₃-C₇ cycloalkyl(C₁-C₄ alkyl), aryl, aryl(C₁-C₄
20 alkyl)-, (C₂-C₇ alkyl)carbonyl, arylcarbonyl,
(C₂-C₁₀ alkoxy)carbonyl, C₃-C₇ cycloalkoxycarbonyl,
C₇-C₁₁ bicycloalkoxycarbonyl, aryloxycarbonyl,
aryl(C₁-C₁₀ alkoxy)carbonyl,
C₁-C₆ alkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl,
25 arylcarbonyloxy(C₁-C₄ alkoxy)carbonyl, and
C₃-C₇ cycloalkylcarbonyloxy(C₁-C₄ alkoxy)carbonyl;

R^{7e} is selected from:

H, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy, aryl, aryl(C₁-
30 C₄ alkyl)-, (C₁-C₄ alkyl)carbonyl, CO₂R^{18ae}, SO₂R^{11e},
SO₂NR^{10e}R^{11e}, OR^{10e}, and N(R^{11e})R^{12e};

U^e is selected from:

-(CH₂)_n^e-, -(CH₂)_n^eO(CH₂)_m^e-, -NH(CH₂)_n^e-,
35 -N(R^{10e})C(=O)-, -NHC(=O)(CH₂)_n^e-, and -C(=O)N(R^{10e})-;

G^e is N or CR^{19e};

R^{8e} is selected from:

- H, CO₂R^{18be}, C(=O)R^{18be}, CONR^{17e}R^{18be},
- C₁-C₁₀ alkyl substituted with 0-1 R^{6e},
- C₂-C₁₀ alkenyl substituted with 0-1 R^{6e},
- 5 C₂-C₁₀ alkynyl substituted with 0-1 R^{6e},
- C₃-C₈ cycloalkyl substituted with 0-1 R^{6e},
- C₅-C₆ cycloalkenyl substituted with 0-1 R^{6e},
- (C₁-C₁₀ alkyl)carbonyl,
- C₃-C₁₀ cycloalkyl(C₁-C₄ alkyl)-,
- 10 phenyl substituted with 0-3 R^{6e},
- naphthyl substituted with 0-3 R^{6e},
- a 5-10 membered heterocyclic ring containing 1-3 N,
- O, or S heteroatoms, wherein said heterocyclic
- ring may be saturated, partially saturated, or
- 15 fully unsaturated, said heterocyclic ring being
- substituted with 0-2 R^{7e};

R^{9e} is selected from:

- C₁-C₁₀ alkyl substituted with 0-1 R^{6e},
- 20 C₁-C₁₀ alkoxy substituted with 0-2 R^{7e},
- H, nitro, N(R^{11e})R^{12e}, OC(=O)R^{10e}, OR^{10e},
- OC(=O)NR^{10e}R^{11e}, NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},
- NR^{10e}C(=O)NR^{10e}R^{11e}, NR^{10e}SO₂NR^{10e}R^{11e},
- NR^{10e}SO₂R^{21e}, hydroxy, OR^{22e}, -N(R^{10e})R^{11e}, -
- 25 N(R^{16e})R^{17e}, aryl(C₀-C₆ alkyl)carbonyl, aryl(C₁-
- C₆ alkyl), heteroaryl(C₁-C₆ alkyl), CONR^{18aer}R^{20e},
- SO₂R^{18ae}, and SO₂NR^{18aer}R^{20e},
- providing that any of the above alkyl, cycloalkyl,
- aryl or heteroaryl groups may be unsubstituted
- 30 or substituted independently with 1-2 R^{7e};

R^{6e} is selected from:

- H, C₁-C₁₀ alkyl, hydroxy, C₁-C₁₀ alkoxy, nitro, C₁-C₁₀
- alkylcarbonyl, -N(R^{11e})R^{12e}, cyano, halo, CF₃,
- 35 CHO, CO₂R^{18be}, C(=O)R^{18be}, CONR^{17e}R^{18be},
- OC(=O)R^{10e}, OR^{10e}, OC(=O)NR^{10e}R^{11e},
- NR^{10e}C(=O)R^{10e}, NR^{10e}C(=O)OR^{21e},

- $\text{NR}^{10\text{e}}\text{C}(=\text{O})\text{NR}^{10\text{e}}\text{R}^{11\text{e}}$, $\text{NR}^{10\text{e}}\text{SO}_2\text{NR}^{10\text{e}}\text{R}^{11\text{e}}$,
 $\text{NR}^{10\text{e}}\text{SO}_2\text{R}^{21\text{e}}$, $\text{S}(\text{O})_{\text{p}}\text{R}^{11\text{e}}$, $\text{SO}_2\text{NR}^{10\text{e}}\text{R}^{11\text{e}}$,
 aryl substituted with 0-3 groups selected from
 halogen, $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_1\text{-C}_6$ alkyl, CF_3 ,
 5 $\text{S}(\text{O})_{\text{m}}\text{Me}$, and $-\text{NMe}_2$,
 aryl($\text{C}_1\text{-C}_4$ alkyl)-, said aryl being substituted with
 0-3 groups selected from halogen, $\text{C}_1\text{-C}_6$ alkoxy,
 $\text{C}_1\text{-C}_6$ alkyl, CF_3 , $\text{S}(\text{O})_{\text{p}}\text{Me}$, and $-\text{NMe}_2$, and
 a 5-10 membered heterocyclic ring containing 1-3 N, O, or
 10 S heteroatoms, wherein said heterocyclic ring may be
 saturated, partially saturated, or fully unsaturated,
 said heterocyclic ring being substituted with 0-2 $\text{R}^{7\text{e}}$;
- $\text{R}^{10\text{e}}$ is selected from:
 15 H, CF_3 , $\text{C}_3\text{-C}_6$ alkenyl, $\text{C}_3\text{-C}_{11}$ cycloalkyl, aryl,
 ($\text{C}_3\text{-C}_{11}$ cycloalkyl)methyl, aryl($\text{C}_1\text{-C}_4$ alkyl), and $\text{C}_1\text{-C}_{10}$
 alkyl substituted with 0-2 $\text{R}^{6\text{e}}$;
- $\text{R}^{11\text{e}}$ is selected from:
 20 H, hydroxy, $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_3\text{-C}_6$ alkenyl, $\text{C}_3\text{-C}_{11}$
 cycloalkyl, ($\text{C}_3\text{-C}_{11}$ cycloalkyl)methyl, $\text{C}_1\text{-C}_6$ alkoxy,
 benzyloxy, aryl, heteroaryl, heteroaryl($\text{C}_1\text{-C}_4$ alkyl)-,
 , aryl($\text{C}_1\text{-C}_4$ alkyl), adamantylmethyl, and
 $\text{C}_1\text{-C}_{10}$ alkyl substituted with 0-2 $\text{R}^{4\text{e}}$;
- 25 $\text{R}^{4\text{e}}$ is selected from:
 H, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_7$ cycloalkyl, $\text{C}_3\text{-C}_7$
 cycloalkyl($\text{C}_1\text{-C}_4$ alkyl)-, aryl, heteroaryl, aryl($\text{C}_1\text{-C}_6$
 alkyl)-, and heteroaryl($\text{C}_1\text{-C}_6$ alkyl)-, wherein
 30 said aryl or heteroaryl groups are substituted with
 0-2 substituents independently selected from the
 group consisting of $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, F, Cl,
 Br, CF_3 , and NO_2 ,
- 35 $\text{R}^{12\text{e}}$ is selected from:
 H, $\text{C}_1\text{-C}_6$ alkyl, triphenylmethyl, methoxymethyl,
 methoxyphenyldiphenylmethyl,
 trimethylsilylethoxymethyl, ($\text{C}_1\text{-C}_6$ alkyl)carbonyl,

(C₁-C₆ alkoxy)carbonyl, (C₁-C₆ alkyl)aminocarbonyl, C₃-C₆ alkenyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-, aryl, heteroaryl(C₁-C₆ alkyl)carbonyl, heteroarylcarbonyl, aryl(C₁-C₆ alkyl)-, 5 (C₁-C₆ alkyl)carbonyl, arylcarbonyl, C₁-C₆ alkylsulfonyl, arylsulfonyl, aryl(C₁-C₆ alkyl)sulfonyl, heteroarylsulfonyl, heteroaryl(C₁-C₆ alkyl)sulfonyl, aryloxy carbonyl, and aryl(C₁-C₆ alkoxy)carbonyl, wherein said aryl groups are 10 substituted with 0-2 substituents selected from the group consisting of C₁-C₄ alkyl, C₁-C₄ alkoxy, halo, CF₃, and nitro;

R^{16e} is selected from:

15 -C(=O)OR^{18ae}, -C(=O)R^{18be}, -C(=O)N(R^{18be})₂, -SO₂R^{18ae}, and -SO₂N(R^{18be})₂;

R^{17e} is selected from:

20 H, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-, aryl, aryl(C₁-C₆ alkyl)-, and heteroaryl(C₁-C₆ alkyl);

R^{18ae} is selected from:

25 C₁-C₈ alkyl optionally substituted with a bond to L_n, C₃-C₁₁ cycloalkyl optionally substituted with a bond to L_n, aryl(C₁-C₆ alkyl)- optionally substituted with a bond to L_n, heteroaryl(C₁-C₆ alkyl)- optionally substituted with a bond to L_n, (C₁-C₆ alkyl)heteroaryl optionally 30 substituted with a bond to L_n, biaryl(C₁-C₆ alkyl) optionally substituted with a bond to L_n, heteroaryl optionally substituted with a bond to L_n, phenyl substituted with 3-4 R^{19e} and optionally substituted with a bond to L_n, 35 naphthyl substituted with 0-4 R^{19e} and optionally substituted with a bond to L_n, and a bond to L_n, wherein said aryl or heteroaryl groups are optionally substituted with 0-4 R^{19e};

R^{18be} is H or R^{18ae};

R^{19e} is selected from:

- 5 H, halogen, CF₃, CO₂H, CN, NO₂, -NR^{11e}R^{12e}, OCF₃,
C₁-C₈ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl,
C₃-C₁₁ cycloalkyl, C₃-C₇ cycloalkyl(C₁-C₄ alkyl)-,
aryl(C₁-C₆ alkyl)-, C₁-C₆ alkoxy, C₁-C₄
alkoxycarbonyl, aryl, aryl-O-, aryl-SO₂-,
10 heteroaryl, and heteroaryl-SO₂-, wherein said aryl
and heteroaryl groups are substituted with 0-4
groups selected from hydrogen, halogen, CF₃, C₁-C₃
alkyl, and C₁-C₃ alkoxy;
- 15 R^{20e} is selected from:
hydroxy, C₁-C₁₀ alkyloxy, C₃-C₁₁ cycloalkyloxy,
aryloxy, aryl(C₁-C₄ alkyl)oxy,
C₂-C₁₀ alkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
C₂-C₁₀ alkoxycarbonyloxy(C₁-C₂ alkyl)oxy-,
20 C₂-C₁₀ alkoxycarbonyl(C₁-C₂ alkyl)oxy-,
C₃-C₁₀ cycloalkylcarbonyloxy(C₁-C₂ alkyl)oxy-,
C₃-C₁₀ cycloalkoxycarbonyloxy(C₁-C₂ alkyl)oxy-,
C₃-C₁₀ cycloalkoxycarbonyl(C₁-C₂ alkyl)oxy-,
aryloxycarbonyl(C₁-C₂ alkyl)oxy-,
25 aryloxycarbonyloxy(C₁-C₂ alkyl)oxy-,
arylcarbonyloxy(C₁-C₂ alkyl)oxy-,
C₁-C₅ alkoxy(C₁-C₅ alkyl)carbonyloxy(C₁-C₂ alkyl)oxy,
(5-(C₁-C₅ alkyl)-1,3-dioxa-cyclopenten-2-one-
yl)methyloxy,
30 (5-aryl-1,3-dioxa-cyclopenten-2-one-yl)methyloxy,
and
(R^{10e})(R^{11e})N-(C₁-C₁₀ alkoxy)-;

R^{21e} is selected from:

- 35 C₁-C₈ alkyl, C₂-C₆ alkenyl, C₃-C₁₁ cycloalkyl, (C₃-C₁₁
cycloalkyl)methyl, aryl, aryl(C₁-C₄ alkyl)-, and
C₁-C₁₀ alkyl substituted with 0-2 R^{7e};

R^{22e} is selected from:

-C(=O)-R^{18be}, -C(=O)N(R^{18be})₂, -C(=O)NHSO₂R^{18ae},
 -C(=O)NHC(=O)R^{18be}, -C(=O)NHC(=O)OR^{18ae}, and
 -C(=O)NHSO₂NHR^{18be};

5

m^e is 0-2;

n^e is 0-4; and

10 p^e is 0-2;

with the following proviso: n^e and m^e are chosen such
 that the number of atoms connecting R¹ and -COR^{20e} in
 Formula (IV) is in the range of 8-14;

15

W is independently selected at each occurrence from the
 group: O, S, NH, NHC(=O), C(=O)NH, NR⁸C(=O), C(=O)N R⁸,
 C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, SO₂NH,
 (OCH₂CH₂)₂₀₋₂₀₀, (CH₂CH₂O)₂₀₋₂₀₀, (OCH₂CH₂CH₂)₂₀₋₂₀₀,

20 (CH₂CH₂CH₂O)₂₀₋₂₀₀, and (aa)_t;

aa is independently at each occurrence an amino acid;

25 Z is selected from the group: aryl substituted with 0-1
 R¹⁰, C₃₋₁₀ cycloalkyl substituted with 0-1 R¹⁰, and a
 5-10 membered heterocyclic ring system containing
 1-4 heteroatoms independently selected from N, S,
 and O and substituted with 0-1 R¹⁰;

30 R⁶, R^{6a}, R⁷, R^{7a}, and R⁸ are independently selected at
 each occurrence from the group: H, =O, COOH, SO₃H,
 C₁-C₅ alkyl substituted with 0-1 R¹⁰, aryl
 substituted with 0-1 R¹⁰, benzyl substituted with 0-1
 R¹⁰, and C₁-C₅ alkoxy substituted with 0-1 R¹⁰,
 35 NHC(=O)R¹¹, C(=O)NHR¹¹, NHC(=O)NHR¹¹, NHR¹¹, R¹¹, and
 a bond to S_f;

k is 0 or 1;

S_f is a surfactant which is a lipid or a compound of the

formula: $A^9-E^1-A^{10}$;

5

A⁹ is OR²⁷;

A¹⁰ is OR²⁷;

10 R²⁷ is C(=O)C₁₋₁₅ alkyl;

E¹ is C₁₋₄ alkylene substituted with 1-3 R²⁸;

15 R²⁸ is independently selected at each occurrence from the group: R³⁰, -PO₃H-R³⁰, =O, -CO₂R²⁹, -C(=O)R²⁹, -CH₂OR²⁹, -OR²⁹, and C₁-C₅ alkyl;

R²⁹ is independently selected at each occurrence from the group: R³⁰, H, C₁-C₆ alkyl, phenyl, and benzyl;

20

R³⁰ is a bond to L_n;

and a pharmaceutically acceptable salt thereof.

25 41. An ultrasound contrast agent composition, comprising:

(a) a compound of Claim 39, comprising: an quinolone that binds to the integrin α_vβ₃, a surfactant and a linking group between the quinolone and the
30 surfactant;

(b) a parenterally acceptable carrier; and,
(c) an echogenic gas.

42. An ultrasound contrast agent composition of Claim
35 41, further comprising: 1,2-dipalmitoyl-sn-glycero-3-phosphotidic acid, 1,2-dipalmitoyl-sn-glycero-3-phosphatidylcholine, and N-(methoxypolyethylene glycol

5000 carbamoyl)-1,2-dipalmitoyl-sn-glycero-3-phosphatidylethanolamine.

43. An ultrasound contrast agent composition of Claim
5 42, wherein the echogenic gas is a C₂₋₅ perfluorocarbon.

44. A method of imaging cancer in a patient comprising:
(1) administering, by injection or infusion, a ultrasound
contrast agent composition of Claim 41 to a patient; and
10 (2) imaging the patient using sonography.

45. A method of imaging formation of new blood vessels
in a patient comprising: (1) administering, by injection
or infusion, a ultrasound contrast agent composition of
15 of Claim 41 to a patient; (2) imaging the area of the
patient wherein the desired formation of new blood
vessels is located.

46. A method of imaging therapeutic angiogenesis in a
20 patient comprising: (1) administering, by injection or
infusion, an ultrasound contrast agent composition of
Claim 41 to a patient; (2) imaging the area of the
patient wherein the desired formation of new blood
vessels is located.

25 47. A method of imaging atherosclerosis in a patient
comprising: (1) administering, by injection or infusion,
an ultrasound contrast agent composition of Claim 41 to a
patient; (2) imaging the area of the patient wherein the
30 atherosclerosis is located.

48. A method of imaging restenosis in a patient
comprising: (1) administering, by injection or infusion,
an ultrasound contrast agent composition of Claim 41 to a
35 patient; (2) imaging the area of the patient wherein the
restenosis is located.

49. A method of imaging cardiac ischemia in a patient comprising: (1) administering, by injection or infusion, an ultrasound contrast agent composition of Claim 41 to a patient; (2) imaging the area of the myocardium wherein
5 the ischemic region is located.

50. A method of imaging myocardial reperfusion injury in a patient comprising: (1) administering, by injection or infusion, an ultrasound contrast agent composition of
10 Claim 41 to a patient; (2) imaging the area of myocardium wherein the reperfusion injury is located.

51. A therapeutic radiopharmaceutical composition, comprising:
15 (a) a therapeutic radiopharmaceutical of Claim 19; and,
(b) a parenterally acceptable carrier.

52. A diagnostic pharmaceutical composition, comprising:
20 (a) a diagnostic radiopharmaceutical, a MRI contrast agent, or a X-ray contrast agent of Claim 11; and,
(b) a parenterally acceptable carrier.

53. A method of treating restenosis in a patient
25 comprising: administering to a patient, either systemically or locally, a therapeutic radiopharmaceutical of Claim 19 capable of localizing in the restenotic area and delivering an effective dose of radiation.

30
54. A method of imaging atherosclerosis in a patient comprising: (1) administering a diagnostic radiopharmaceutical, a MRI contrast agent, or a X-ray contrast agent of Claim 11 to a patient by injection or
35 infusion; (2) imaging the area of the patient wherein the atherosclerosis is located.

55. A method of imaging restenosis in a patient comprising: (1) administering a diagnostic radiopharmaceutical, a MRI contrast agent, or a X-ray contrast agent of Claim 11 to a patient by injection or
5 infusion; (2) imaging the area of the patient wherein the restenosis is located.

56. A method of imaging cardiac ischemia in a patient comprising: (1) administering a diagnostic
10 radiopharmaceutical, a MRI contrast agent, or a X-ray contrast agent of Claim 11 to a patient by injection or infusion; (2) imaging the area of the myocardium wherein the ischemic region is located.

15 57. A method of imaging myocardial reperfusion injury in a patient comprising: (1) administering a diagnostic radiopharmaceutical, a MRI contrast agent, or a X-ray contrast agent of Claim 11 to a patient by injection or
20 infusion; (2) imaging the area of myocardium wherein the reperfusion injury is located.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 June 2000 (22.06.2000)

PCT

(10) International Publication Number
WO 00/35492 A3

- (51) International Patent Classification⁷: A61K 47/48, 49/00, 51/04, 49/04
- (74) Agent: OBRIEN, Maureen, P.; E.I. du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).
- (21) International Application Number: PCT/US99/30315
- (81) Designated States (*national*): AL, AU, BR, CA, CN, CZ, EE, HU, IL, IN, JP, KR, LT, LV, MK, MX, NO, NZ, PL, RO, SG, SI, SK, TR, UA, VN, ZA.
- (22) International Filing Date:
17 December 1999 (17.12.1999)
- (25) Filing Language: English
- (84) Designated States (*regional*): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- (26) Publication Language: English
- (30) Priority Data:
60/112,732 18 December 1998 (18.12.1998) US
- Published:
— With international search report.
- (71) Applicant: DU PONT PHARMACEUTICALS COMPANY [US/US]; Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19807 (US).
- (88) Date of publication of the international search report:
18 January 2001
- (72) Inventors: HARRIS, Thomas, David; 56 Zion Hill Road, Salem, NH 03079 (US). RAJODADHYE, Milind; 21 Honeysuckle Road, Westford, MA 01886 (US).
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 00/35492 A3

(54) Title: VITRONECTIN RECEPTOR ANTAGONIST PHARMACEUTICALS

(57) Abstract: The present invention describes novel compounds of formula: (Q)_q-L_n-C_h, useful for the diagnosis and treatment of cancer, methods of imaging tumors in a patient, and methods of treating cancer in a patient. The present invention also provides novel compounds useful for monitoring therapeutic angiogenesis treatment and destruction of new angiogenic vasculature. The present invention further provides novel compounds useful for imaging atherosclerosis, restenosis, cardiac ischemia, and myocardial reperfusion injury. The present invention still further provides novel compounds useful for the treatment of rheumatoid arthritis. The pharmaceuticals are comprised of a targeting moiety that binds to a receptor that is upregulated during angiogenesis, and optional linking group, and a therapeutically effective radioisotope or diagnostically effective imageable moiety. The imageable moiety is a gamma ray or positron emitting radioisotope, a magnetic resonance imaging contrast agent, and X-ray contrast agent, or an ultrasound contrast agent.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/30315

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K47/48 A61K49/00 A61K51/04 A61K49/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, WPI Data, PAJ, EMBASE, BIOSIS, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 16256 A (BURNHAM INST) 23 April 1998 (1998-04-23) abstract page 26; claim 1; examples 3,4 ---	1-57
E	US 6 040 311 A (DUGGAN MARK E ET AL) 21 March 2000 (2000-03-21) claims 2,27,29,32 ---	1-57
P,X	WO 99 06049 A (SMITHKLINE BEECHAM CORP ;HEERDING DIRK (US); SAMANEN JAMES MARTIN) 11 February 1999 (1999-02-11) page 15, line 33-36; claims 14,16,18 page 22 --- -/-	1-57



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

3 August 2000

Date of mailing of the international search report

14/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gonzalez Ramon, N

INTERNATIONAL SEARCH REPORT

Inter. Appl. Application No

PCT/US 99/30315

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 96 00574 A (COUSINS RUSSELL DONOVAN ;SMITHKLINE BEECHAM CORP (US); UZINSKAS IR) 11 January 1996 (1996-01-11) page 6, line 31-36; claims 12,13	1-57
Y	WO 97 08145 A (DESAI BIPINCHANDRA NANUBHAI ;LINDMARK RICHARD JOHN (US); RUSSELL M) 6 March 1997 (1997-03-06) examples 9,10,156,178,179 examples 190,191,238,239,324	1-57
Y	WO 98 14220 A (DU PONT MERCK PHARMA) 9 April 1998 (1998-04-09) abstract page 28	1-57
T	VAN WAES C. ET AL: "Effects of the novel av integrin antagonist SM256 and cis-platinum on growth of murine squamous cel carcinoma PAM LY8" INTERNATIONAL JOURNAL OF ONCOLOGY, vol. 16, 2000, pages 1189-1195, XP000925238 abstract; figure 1 see discussion page 1193	1-57
Y,P	LIU, SHUANG ET AL: "Technetium Complexes of a Hydrazinonicotinamide-Conjugated Cyclic Peptide and 2-Hydrazinopyridine: Synthesis and Characterization" INORG. CHEM. (1999), 38(6), 1326-1335 , XP000924906 abstract; figure 1	1-57
P,X	DENARDO S J ET AL: "Neovascular targeting with cyclic RGD peptide (cRGDf-ACHA) to enhance delivery of radioimmunotherapy." CANCER BIOTHER RADIOPHARM, (2000 FEB) 15 (1) 71-9. , XP000925000 abstract	1-57
P,Y	KERR J S ET AL: "Novel small molecule alpha v integrin antagonists: comparative anti-cancer efficacy with known angiogenesis inhibitors." ANTICANCER RESEARCH, (1999 MAR-APR) 19 (2A) 959-68. , XP000924849 see discussion abstract	1-57

-/-

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/30315

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
T	<p>HORTON: "The avb3 integrin "vitronectin receptor"" INT J. BIOCHEM. CELL. BIOL., vol. 29, no. 5, May 1997 (1997-05), pages 721-725, XP000923474 the whole document</p>	1-57

FURTHER INFORMATION CONTINUED FROM PCT/ISA 210

Continuation of Box I.1

Although claims 31,32,53 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
Although claims 33-36,44-50,54-57 are directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.2

Claims Nos.: 1-4,7-15,17,19-22,24,26-28,30-57

Present claims 1-57 relate to compounds/kits defined by reference to desirable characteristics or properties, namely "receptor that is upregulated during angiogenesis", "targeting moiety", "chelator", "linking groups", "a surfactant", "quinolone nonpeptide".
The claims cover all compounds/kits having these characteristics or properties, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds/kits. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compounds by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible.
Moreover claims 1-4,7-15,17,19-22,24,26-28,30-57 relate to an extremely large number of possible compounds/kits. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds/kits claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.
Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the compounds/kits prepared in the examples and specifically mentioned by name in the claims with due regard to the general idea underlying the present application.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int'l. Application No
PCT/US 99/30315

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9816256 A	23-04-1998	AU 4753997 A	11-05-1998
US 6040311 A	21-03-2000	NONE	
WO 9906049 A	11-02-1999	AU 8689798 A EP 1007051 A ZA 9806930 A	22-02-1999 14-06-2000 13-05-1999
WO 9600574 A	11-01-1996	AU 702661 B AU 3001095 A BR 9508178 A CA 2193966 A CN 1156995 A CZ 9603824 A EP 0762882 A EP 0767792 A HU 76344 A JP 10504807 T JP 10504808 T NO 965608 A NZ 290008 A PL 318199 A WO 9600730 A ZA 9505391 A	25-02-1999 25-01-1996 18-11-1997 11-01-1996 13-08-1997 17-12-1997 19-03-1997 16-04-1997 28-08-1997 12-05-1998 12-05-1998 27-02-1997 26-08-1998 26-05-1997 11-01-1996 09-02-1996
WO 9708145 A	06-03-1997	AU 702487 B AU 7103996 A BR 9610422 A CA 2230209 A CN 1201454 A CZ 9800341 A EP 0850221 A JP 11510814 T NO 980817 A NZ 318926 A PL 325312 A	25-02-1999 19-03-1997 13-07-1999 06-03-1997 09-12-1998 16-09-1998 01-07-1998 21-09-1999 24-04-1998 29-04-1999 20-07-1998
WO 9814220 A	09-04-1998	AU 4892597 A EP 0993307 A US 5888970 A ZA 9708758 A	24-04-1998 19-04-2000 30-03-1999 30-03-1999

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/28662

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :A61K 31/435

US CL :CO7D 471/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : IPC 7: CO7D 471/02.

U.S. 514/337; 546/122.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE (1907-2000) STN DATABASES File Registry, File Beilstein, File Marpat

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,599,936 A (BARREAU et al.) 04 February 1997, see cols. 1-10.	1-18 & 23

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

05 FEBRUARY 2000

Date of mailing of the international search report

20 MAR 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ALAN L. ROTMAN

Telephone No. (703) 308-0196